

Use of image analysis in the evaluation of radicular nodules in chickpeas

Utilização de análise de imagens na avaliação de nódulos radiculares em grão-de-bico

Uso del análisis de imagen en la evaluación de nódulos radiculares en garbanzos

DOI: 10.54033/cadpedv21n9-002

Originals received: 07/29/2024

Acceptance for publication: 08/19/2024

Karla Sabrina Magalhães Andrade Padilha

Doctorate Student in Plant Production

Institution: Universidade Federal de Minas Gerais (UFMG-ICA)

Address: Montes Claros, Minas Gerais, Brazil

E-mail: karlasabrina@gmail.com

Pedro Vitor de Souza Silva

Doctorate Student in Biotechnology

Institution: Universidade Estadual de Montes Claros (UNIMONTES)

Address: Montes Claros, Minas Gerais, Brazil

Email: pedrovitorsilvabio@gmail.com

Alcinei Místico Azevedo

Doctor in Plant Science

Institution: Universidade Federal de Minas Gerais (UFMG-ICA)

Address: Montes Claros, Minas Gerais, Brazil

E-mail: alcineimistico@ufmg.br

Rodinei Facco Pegoraro

Doctor in Soils and Plant Nutrition

Institution: Universidade Federal de Minas Gerais (UFMG-ICA)

Address: Montes Claros, Minas Gerais, Brazil

E-mail: rodinei@ufmg.br

Aline Martins Ferreira Barroso

Doctor Student in Plant Production

Institution: Universidade Federal de Minas Gerais (UFMG-ICA)

Address: Montes Claros, Minas Gerais, Brazil

E-mail: alinemartinsfb16@hotmail.com

Verônica Aparecida Santos Ferreira Soares

Doctor in Plant Production

Institution: Universidade Federal de Minas Gerais (UFMG-ICA)

Address: Montes Claros, Minas Gerais, Brazil

E-mail: veronicaasf@yahoo.com.br

Silvana Ferreira Bicalho

Doctorate Student in Plant Production

Institution: Universidade Federal de Minas Gerais (UFMG-ICA)

Address: Montes Claros, Minas Gerais, Brazil

E-mail: silvana.bicalho@uesb.edu.br

ABSTRACT

The use of advanced computational systems facilitates the application of a wide range of statistical techniques, available in open-source format, to perform analyses of substantial complexity in biological systems, particularly in the field of plant biology. The aim of this study is to demonstrate the application of image analysis in the evaluation of root nodules in *Cicer arietinum* plants. The research was conducted in the field, where roots were collected, subjected to rigorous cleaning protocols and subsequently photographed under precisely controlled studio conditions. This photographic process was performed using equipment configured with meticulous parameter settings. Subsequently, image analyses were performed using the statistical software package, R. Parameters of interest were quantified, including metrics such as root area, nodule area, the proportional representation of nodules in relation to the total root mass and the absolute count of nodules present. A critical comparison between the proposed analytical methodology and conventional approaches evidenced a substantial improvement in efficiency, thus highlighting the robustness and validity of this analytical framework for the outlined research objectives. Therefore, the developed methodology has significant potential to facilitate accurate and comprehensive analyses of nodules and root systems, thus making a notable contribution to fostering a deeper understanding and exploration of plant-microorganism interactions. Furthermore, this methodological advancement has the potential to contribute to reducing operational costs and optimizing the time spent on such meticulous evaluative undertakings.

Keywords: Computational Analysis. Chickpeas. Root Characteristics. High-Efficiency Phenotyping.

RESUMO

A utilização de sistemas computacionais avançados facilita a aplicação de uma ampla gama de técnicas estatísticas, disponíveis em formato de código aberto, para a realização de análises de complexidade substancial em sistemas biológicos, particularmente no domínio da biologia vegetal. O objetivo deste estudo é demonstrar a aplicação da análise de imagens na avaliação de nódulos radiculares em plantas de *Cicer arietinum*. A pesquisa foi conduzida em campo, onde as raízes foram coletadas, submetidas a protocolos rigorosos de limpeza e

posteriormente fotografados sob condições de estúdio precisamente controladas. Este processo fotográfico foi executado utilizando equipamentos configurados com ajustes de parâmetros meticulosos. Posteriormente, as análises de imagem foram realizadas utilizando o pacote de software estatístico, R. Parâmetros de interesse foram quantificados, abrangendo métricas como área da raiz, área do nódulo, a representação proporcional dos nódulos em relação à massa total da raiz e a contagem absoluta de nódulos presentes. Uma crítica comparação entre a metodologia analítica proposta e abordagens convencionais evidenciou um substancial aprimoramento na eficiência, destacando assim a robustez e validade deste arcabouço analítico para os objetivos de pesquisa delineados. Portanto, a metodologia desenvolvida possui um potencial significativo para facilitar análises precisas e abrangentes de nódulos e sistemas radiculares, conferindo assim uma contribuição notável para fomentar uma compreensão mais profunda e exploração das interações planta-microorganismo. Além disso, este avanço metodológico tem o potencial de contribuir para a redução dos custos operacionais e otimização do tempo gasto em tais empreendimentos avaliativos meticulosos.

Palavras-chave: Análise Computacional. Grão-de-Bico. Características da Raiz. Fenotipagem de Alta Eficiência.

RESUMEN

El uso de sistemas computacionales avanzados facilitó la aplicación de una amplia gama de técnicas estadísticas, disponibles en formato de código abierto, para realizar análisis de sustancial complejidad en sistemas biológicos, particularmente en el campo de la biología vegetal. El objetivo de este estudio es demostrar la aplicación del análisis de imágenes en la evaluación de nódulos radiculares en plantas de *Cicer arietinum*. La investigación se llevó a cabo en el campo, donde se recolectaron raíces, se sometieron a rigurosos protocolos de limpieza y posteriormente se fotografiaron en condiciones de estudio controladas con precisión. Este proceso fotográfico se llevó a cabo utilizando equipos configurados con meticulosos ajustes de parámetros. Posteriormente, se realizaron análisis de imágenes utilizando el paquete de software estadístico R. Se cuantificaron los parámetros de interés, abarcando métricas como el área de la raíz, el área de los nódulos, la representación proporcional de los nódulos en relación con la masa total de la raíz y el recuento absoluto de nódulos presente. Una comparación crítica entre la metodología analítica propuesta y los enfoques convencionales destacó una mejora sustancial en la eficiencia, destacando así la solidez y validez de este marco analítico para los objetivos de investigación delineados. Por lo tanto, la metodología desarrollada tiene un potencial significativo para facilitar análisis precisos y completos de nódulos y sistemas de raíces, contribuyendo así a fomentar una comprensión y exploración más profundas de las interacciones planta-microorganismo. Además, este avance metodológico tiene el potencial de contribuir a reducir los costos operativos y optimizar el tiempo dedicado a tareas de evaluación tan meticulosas.

Palabras clave: Análisis Computacional. Garbanzo. Características de la Raíz. Fenotipado de Alta Eficiencia.

1 INTRODUCTION

The use of equipment and software dedicated to analysis has proven to be of paramount importance in the pursuit of precise and expedited results. Employing strategies that enable the acquisition and analysis of data in agricultural environments can play a crucial role in optimizing current agricultural practices. This translates into increased productivity, improved quality control processes, and greater flexibility in agricultural management (Vasconez *et al.*, 2020).

Agriculture stands as one of the most prominent activities globally. With the rapid growth of the world population, the demand for food is continually rising. Consequently, new automated methods are being introduced to meet these needs, as traditional methods employed by farmers prove insufficient to address such demands (Zhang *et al.*, 2021; Dharmaraj *et al.*, 2018). In this scenario, the adoption of alternative high-efficiency phenotyping tools becomes indispensable. In this context, the use of images combined with computer vision deserves attention, as it allows for the assessment of qualitative and productivity attributes with reduced costs, lower labor demand, greater speed, efficiency, and precision (De Mesquita *et al.*, 2022).

Agriculture is currently facing significant challenges, including climate change, resource depletion, loss of biodiversity, labor shortages, and the urgent need to increase the production of nutrient-rich foods (Tariq *et al.*, 2023). In this regard, chickpea (*Cicer arietinum* L.) emerges as a legume of special interest, representing an economically accessible source of protein. Moreover, it is considered the second most consumed legume globally, playing a crucial role in regions facing a shortage of animal-origin protein (Zhang *et al.*, 2020).

The cultivation of chickpeas exhibits attractive attributes in the context of sustainable production, given its remarkable adaptability to various climatic conditions, along with significant potential in terms of both productivity and economic profitability (Grasso *et al.*, 2022). Another important aspect of chickpea cultivation lies in the ability of its roots to establish symbiosis with nitrogen-fixing bacteria, such as those of the *Rhizobium ciceri* genus, resulting in the formation of nodules capable of fixing atmospheric nitrogen in a form usable by plants.

Several research efforts have been directed towards this crop, and the application of high-efficiency phenotyping techniques through image analysis emerges as a tool of significant potential.

The use of images holds significant potential for enhancing efficiency and precision in phenotyping activities, as indicated by Haque *et al.* (2021). This approach allows for the assessment of qualitative attributes in a more economical, rapid, efficient, and precise manner. With the advancement of new technologies aiming to increase the accuracy and speed of phenotypic measurements, there is a growing interest in research in this field (Massruhá *et al.*, 2014).

An advantage of these methodologies is the ability to reproduce results even after the elimination of plant material, as the image database can be preserved. This opens the possibility of obtaining detailed information about root development through image analysis (Haque *et al.*, 2021).

Another significant benefit is methodological standardization, which assumes critical relevance by providing consistency, reliability, and a foundation for study comparison. This ensures that processes and procedures are uniformly followed. Therefore, the objective of this study is to demonstrate the application of image analysis in the evaluation of root nodules in chickpea plants.

2 METODOLOGY

2.1 STUDY AREA AND EXPERIMENTAL CONDITIONS

The field study to obtain chickpea root samples was conducted in an experimental area located in the municipality of Montes Claros – MG, at geographical coordinates 16°40'57.5"S 43°50'21.4"W, from June to September 2022, under a microsprinkler irrigation system. The seeding was carried out manually in the planting furrow with 30 seeds per linear meter. The experimental area was divided into 32 plots, each containing four cultivation rows, with one linear meter in length and 40 plants per plot.

In this study, the BRS Cristalino cultivar was used, previously inoculated with symbiotic microorganisms before seeding. Weed control was conducted manually, using hoeing as needed.

2.2 ACQUISITION AND PROCESSING OF ROOT SYSTEM IMAGES

The chickpea plants were harvested at 70 (seventy) days after planting during full bloom. During this period, ten central plants per plot, totaling 320 plants, were collected for image acquisition. After removing the plants from the soil, the roots were carefully separated from the plants, washed with tap water over a sieve (2.0mm mesh) to prevent the loss of nodules.

The dried roots were then placed in a studio for photography using a camera with ISO320, SPEED 1/1500 F1.5 M0.6, WB490K settings, under artificial lighting with a fluorescent lamp. For each sample (10 plants), three images were taken to capture the ten roots without overlap to avoid errors (a total of 96 images analyzed, with 3 images per plot).

The camera was mounted on a stand in a mini studio with dimensions of 60 x 60 x 60 cm. This strategy was implemented to standardize the image acquisition process, ensuring that all images were captured from the same height and angle. The roots were positioned against a blue background and arranged in a non-overlapping manner, ensuring there was space between them. The analysis of root characteristics was conducted using the R® software, employing the ExplImage package (Azevedo, 2023).


A computational analysis was initiated by resizing the images, originally with dimensions of 2262 x 2896 pixels, to a size of 1000 x 1279 pixels using the *"resize_image"* function. This procedure aimed to reduce the computational analysis time, as illustrated in Figure 1 (A).

Subsequently, the image was converted to grayscale, as shown in Figure 1 (B), by extracting the red band with the *"gray_scale"* function for the segmentation of the reference object (2 x 2 cm). The threshold method was then applied through the segmentation function with a value of 0.9, as depicted in Figure 1 (C).

For the segmentation of roots and nodules (Figure 1 D), the image was converted to grayscale by extracting the blue band. Subsequently, segmentation was performed considering a threshold of 0.6. Two erosions were applied to the images using the *"erode_image"* function, followed by two dilations using the *"dilate_image"* function. This strategy allowed the exclusion of pixels corresponding to small-thickness objects (roots) from the image, followed by restoring the remaining pixels to their original size. The *"mask_pixels"* function was used to highlight the objects created through a mask. The *"measure_image"* function was employed for nodule counting and area estimation (Figure 1 F). The conversion from pixel² to cm² was performed by a simple proportion considering the area and number of pixels of the reference object.

For a better visualization of the efficiency of nodule and root segmentation on the original background (Figure 1 G) and on the black background (Figure 1 H), the *"mask_pixels"* function was also used.

Figure 1 – Image Acquisition: original image and reference masks for roots and nodules, and efficiency of the analyses in R® software.



Source: From the authors, 2024.

For comparison and evaluation of the method's applicability, a manual count of the nodules present in the roots of the plants was carried out. This was done on the roots that were previously photographed after image capture.


3 RESULTS

A routine in R allows for the estimation of parameters with different scales, and thus, the functions used were efficient in estimating the following parameters of the chickpea root system: root area (cm²), nodule area (cm²), percentage of nodules in relation to roots, and number of nodules. The analyses of masks followed by segmentation and estimation of the number of nodules through the functions *erode_image*, *dilate_image*, and *measure_image* was also efficient in estimating the area of root nodules. These analyses provide higher quality in the estimation of root parameters by quickly determining living components that undergo rapid decomposition over a short period.

The time optimization in the process of obtaining root parameters stands out as one of the main benefits of using images in this study. This is because roots and root nodules quickly lose viability, dehydrate, and decompose shortly after being removed from the soil. This uniqueness justifies the unprecedented nature of this study, as the computational analysis of root nodules, especially in chickpea cultivation, lacks precedents in the scientific literature. Regarding root analyses, the limitation lies in the complexity of performing manual analyses, as such an approach requires highly specialized labor and is limited to a small number of treatments. However, the use of image analyses offers a solution to these issues, providing precise, fast, and reliable results.

The comparative results between nodule counts performed through computational analysis and manual counting highlight the method's effectiveness, as the discrepancy between the two approaches was reduced, as illustrated in Figure 2 (A), and furthermore, a coefficient of determination close to 1 was obtained (Figure 2 B). It is worth noting that manual counting has some disadvantages, such as the need for more time and the demand for highly specialized labor.

Figure 2 – Comparison between the estimated number of nodules (NNE) through image analysis and the traditional method (NNC) using a bar chart (A) and regression analysis (B).



Source: From the authors, 2024.

4 DISCUSSION

Image analysis plays a crucial role in agriculture, providing valuable information and supporting various aspects of crop management. Image analysis enables the monitoring of crops at different growth stages, helping farmers assess crop health, detect diseases, pests, and nutrient deficiencies (Anwar & Polder, 2020). In this study, these analyses allowed for the estimation of valuable root parameters for scientific activities and chickpea production.

In our scientific study, it was possible to estimate root area (cm²), nodule area (cm²), percentage of nodules in relation to roots, and number of nodules, indicating the high reliability and efficiency of the method used compared to conventional manual nodule counting methods. Works conducted by Fernandes *et al.* (2023) with sweet potato roots, using R software and the ExplImage package, also yielded satisfactory results. They concluded that image analyses combined with deep learning improve study quality, reducing subjective analyses and phenotyping time for the crop.

These analyses in agriculture offer a non-invasive and efficient means of collecting data for decision-making, allowing agricultural science professionals to optimize resource allocation, minimize inputs, mitigate risks, and improve overall agricultural productivity and sustainability (Ge *et al.*, 2018). Haque *et al.* (2021) and Fernandes *et al.* (2023), working with image analysis in sweet potatoes,

verified the high potential of this technology in phenotyping, specifically in predicting root weight. According to Fernandes *et al.* (2022), these images can be stored in a database for future queries and comparisons, enabling simultaneous processing.

Lobet *et al.* (2015) also conducted a scientific study using images to describe root system architecture through the Root System Markup Language. The aim was to enable the portability of root architecture data among different software tools in an easy and interoperable manner and to provide a standard format for root analyses. They concluded that the system accommodates a wide range of root architecture complexities at various levels of detail, storing root topology and any additional information used to describe root segments. This underscores the importance of using artificial intelligence for root analyses.

In works by Bucksch *et al.* (2014), they also present a high-throughput image-based phenotyping platform for studying roots in field crops. They demonstrate the application of image analysis techniques for characterizing root systems and conclude that the development of image processing algorithms aligned with the needs of current field research practices presents an excellent alternative for estimating root phenotypes in real soil conditions.

Similar to the results found in this study, Bucksch *et al.* (2014) report that these analyses have a low cost and can automatically classify various root components in a single image. From these studies, many phenotypic assessments can be conducted in a shorter time frame (Haque *et al.*, 2021). The use of image analysis for root system-related studies in agriculture provides opportunities to investigate other parameters and a multitude of crops with a short time frame and low cost, enabling high-throughput phenotyping.

5 CONCLUSION

The parameters obtained through image analysis encompassed root area (cm²), nodule area (cm²), the proportion of nodules to total roots, and the nodules count. Upon comparing nodule counts derived from image analysis estimates with traditional counting methods, minimal differences were observed, affirming

the efficacy of this approach for its intended purpose. The utilization of image analysis emerged as a viable strategy for precisely and swiftly estimating the quantity of root nodules. This capability holds considerable promise for research endeavors concerning chickpea cultivation, underscoring the significance of integrating image analysis methodologies.

Nevertheless, it's crucial to acknowledge the limitations of this study. For instance, variations in lighting conditions during image acquisition or the complexity of root structures may introduce some degree of error into the analysis. Furthermore, while the method demonstrated effectiveness in estimating nodule quantity, additional research is warranted to explore its applicability across different plant species and environmental conditions.

Reflecting on the societal and academic implications, the results of this research offer notable benefits. In agricultural contexts, accurate and efficient estimation of root nodules can enhance crop management practices, leading to improved yields and resource utilization. Additionally, the adoption of image analysis methodologies can streamline data collection processes, enabling researchers to conduct larger-scale studies and accelerate the pace of scientific discovery in plant biology.

Moreover, the integration of image analysis into agricultural research contributes to broader societal goals such as food security and sustainability. By facilitating more precise monitoring and management of crop health, this technology has the potential to mitigate the impact of environmental stressors and optimize agricultural production systems. Ultimately, the findings of this study not only advance scientific knowledge but also offer practical solutions to real-world challenges, benefiting both academia and society.

ACKNOWLEDGEMENTS

This work was carried out with the support of the Coordination for the Improvement of Higher Education Personnel, Brazil (CAPES) – Financing Code 001, the National Council for Scientific Development (CNPQ), and the Research Support Foundation of Minas Gerais (FAPEMIG).

REFERENCES

ANWAR, S.; POLDER, G. Use of satellite remote sensing data for crop disease management: a review. **Remote Sensing**, v. 12, n. 3, p. 464, 2020.

AZEVEDO, A. M. **ExpImage**: Tool For Analysis of Images in Experiments. CRAN package repository, 2023. Disponível em: <https://www.expstat.com/pacotes-do-r/expimage>. Acesso em: 19 jun. 2023.

BUCKSCH, A. *et al.* Image-based high-throughput field phenotyping of crop roots. **Plant Physiology**, v. 166, n. 2, p. 470-486, 2014. Disponível em: <https://academic.oup.com/plphys/article/166/2/470/6113282?login=false>. Acesso em: 06 jun. 2023.

DHARMARAJ, V.; VIJAYANAND, Chinnusamy. Artificial intelligence (AI) in agriculture. **International Journal of Current Microbiology and Applied Sciences**, v. 7, n. 12, p. 2122-2128, 2018. Disponível em: <https://doi.org/10.20546/ijcmas.2018.712.241>. Acesso em: 12 jun. 2023.

FERNANDES, A. C. G. *et al.* Convolutional neural networks in the qualitative improvement of sweet potato roots. **Scientific Reports**, v. 13, n. 1, p. 8429, 2023. Disponível em: <https://www.nature.com/articles/s41598-023-34375-6>. Acesso em: 03 jun. 2023.

FERNANDES, A. C. G. *et al.* Feasibility of computational vision in the genetic improvement of sweet potato root production. **Horticultura Brasileira**, v. 40, n. 4, p. 378-383, 2022. Disponível em: <https://www.scielo.br/j/hb/a/Gj6QyQkqQtyfn-XKDXK3MVRv/>. Acesso em: 03 jun. 2023.

GE, Y.; THOMASSON, J. A.; SUI, R.; PAUL, G. Machine learning models for assessing crop nitrogen status with high-resolution multi-spectral imagery. **Remote Sensing of Environment**, v. 216, p. 474-486, 2018.

GRASSO, N. *et al.* Chickpea protein ingredients: A review of composition, functionality, and applications. **Comprehensive reviews in food science and food safety**, v. 21, n. 1, p. 435-452, 2022. Disponível em: https://ift.onlinelibrary.wiley.com/doi/full/10.1111/1541-4337.12878?casa_token=HBh6TFvmpBcAAAAA%3AxOBr6wWiNqD2C1uL_eXuQD5L3DA1reljolBRJ6IYVsLlkrNZcJ1qe syVAtCRJcLRhjr1SWLENKeWAcUwfw. Acesso em: 04 jun. 2023.

HAQUE, S. *et al.* Computer vision approach to characterize size and shape phenotypes of horticultural crops using high-throughput imagery. **Computers and Electronics in Agriculture**, v. 182, p. 106011, 2021. Disponível em: https://www.sciencedirect.com/science/article/pii/S0168169921000296?casa_token=9zKqjw6jOakAAAAA:jXxwUiFSdnSPDymX5aFr_oh715ghheHhCDGtan4ag8URWb7iyqYL7vdxwC1cTo0HQL-HLmuGu32N. Acesso em: 04 jun. 2023.

LOBET, G. *et al.* Root system markup language: toward a unified root architecture description language. **Plant physiology**, v. 167, n. 3, p. 617-627,

2015. Disponível em: <https://academic.oup.com/plphys/article/167/3/617/6113575?login=false>. Acesso em: 3 jun. 2023.

MASSRUHÁ, S. M. F. S.; LEITE, M. A. A.; LUCHIARI JUNIOR, A.; ROMANI, L. A. S. **Tecnologias da Informação e Comunicação e suas relações com a agricultura**. Brasília, DF: Embrapa, p. 85–100, 2014.

PINHEIRO, R. de M.; GADOTTI, G. I.; BERNARDY, R.; MONTEIRO, R. D. C. M.; MOREIRA, I. B. Image processing as to important tool for artificial intelligence in the seed sector. **Revista Agrária Acadêmica**, v. 5, n. 1, 2022.

TARIQ, M. *et al.* Reconnoitering the capabilities of nodule endophytic *Pantoea dispersa* for improved nodulation and grain yield of chickpea (*Cicer arietinum* L.). **World Journal of Microbiology and Biotechnology**, v. 39, n. 3, p. 85, 2023. Disponível em: <https://link.springer.com/article/10.1007/s11274-023-03525-3>. Acesso em: 06 junh. 2023.

VASCONEZ, J. P.; DELPIANO, J.; VOUGIOUKAS, S.; CHEEIN, F. A. Comparison of convolutional neural networks in fruit detection and counting: A comprehensive evaluation. **Computers and Electronics in Agriculture**, v. 173, p. 105348, 2020. Disponível em: https://www.sciencedirect.com/science/article/pii/S016816991932232X?casa_token=gCqao0M2oA8AAAAA:JCDNA1leQxUUrzycvls1pTnEqjPvp2vjpYOmvXtHkkqATVbGIDGVbyNcrBWLawEz4inpserv43i2. Acesso em: 07 jun. 2023.

ZHANG, J. *et al.* Biogeographic distribution of chickpea rhizobia in the world. In: **Molecular Aspects of Plant Beneficial Microbes in Agriculture**. Academic press, 2020. p. 235-239. Disponível em: <https://www.sciencedirect.com/science/article/abs/pii/B9780128184691000201>. Acesso em: 08 jun. 2023.

ZHANG, P. *et al.* Nanotechnology and artificial intelligence to enable sustainable and precision agriculture. **Nature Plants**, v. 7, n. 7, p. 864-876, 2021. Disponível em: <https://www.nature.com/articles/s41477-021-00946-6>. Acesso em: jun. 2023.