

Research



**Cite this article:** Silva JL, Demolin Leite GL, de Souza Tavares W, Souza Silva FW, Sampaio RA, Azevedo AM, Serrão JE, Zanuncio JC. 2020 Diversity of arthropods on *Acacia mangium* (Fabaceae) and production of this plant with dehydrated sewage sludge in degraded area. *R. Soc. open sci.* **7**: 191196. <http://dx.doi.org/10.1098/rsos.191196>

Received: 9 July 2019

Accepted: 3 February 2020

**Subject Category:**

Organismal and evolutionary biology

**Subject Areas:**

ecology

**Keywords:**

arthropods, diversity, fertilization, litter, *Nasutitermes*, *Trigona spinipes*

**Author for correspondence:**

Germano Leão Demolin Leite  
e-mail: [gldleite@ica.ufmg.br](mailto:gldleite@ica.ufmg.br)

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.4855431>.

# Diversity of arthropods on *Acacia mangium* (Fabaceae) and production of this plant with dehydrated sewage sludge in degraded area

Júlia Leticia Silva<sup>1</sup>, Germano Leão Demolin Leite<sup>1</sup>, Wagner de Souza Tavares<sup>2</sup>, Farley William Souza Silva<sup>3</sup>, Reginaldo Arruda Sampaio<sup>1</sup>, Alcinei Místico Azevedo<sup>1</sup>, José Eduardo Serrão<sup>4</sup> and José Cola Zanuncio<sup>5</sup>

<sup>1</sup>Instituto de Ciências Agrárias, Universidade Federal de Minas Gerais, 39404-006 Montes Claros, Minas Gerais, Brasil

<sup>2</sup>Asia Pacific Resources International Ltd. (APRIL), PT. Riau Andalan Pulp and Paper (RAPP), Pangkalan Kerinci, Riau, 28300 Sumatra, Indonesia

<sup>3</sup>Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre, 69920-900 Rio Branco, Acre, Brasil

<sup>4</sup>Departamento de Biologia Geral and <sup>5</sup>Departamento de Entomologia/BIOAGRO, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brasil

JLS, 0000-0002-4394-1578; GLDL, 0000-0002-2928-3193; WdST, 0000-0002-8394-6808; FWSS, 0000-0002-2651-9375; RAS, 0000-0003-3214-6111; AMA, 0000-0001-5196-0851; JES, 0000-0002-0477-4252; JCZ, 0000-0003-2026-281X

Sewage sludge is an organic matter-rich material with abundant fractions of nitrogen and other macro and micronutrients, essential for plant growth and development such as *Acacia mangium* Willd. (Fabales: Fabaceae) used in recovering actions of degraded areas. The objective of this study was to evaluate over 24 months the abundance and diversity of chewing and pollinator insects and arthropod predators on *A. mangium* plants and the mass production and soil coverage by this plant, fertilized with dehydrated sewage sludge, in a degraded area. The experimental design was in randomized blocks with two treatments (with and without dehydrated sewage sludge) and 24 replications. The number of leaves per branch and branches per plant, defoliation percentage by chewing insects, soil cover and abundance of chewing and pollinator insects and arthropod predators were higher on *A. mangium* plants fertilized with dehydrated sewage sludge. *Nasutitermes* sp. (Blattodea: Termitidae) and *Trigona spinipes* F. (Hymenoptera:

Apidae) were the most observed insects on trunks and leaves, respectively, of *A. mangium* plants fertilized with dehydrated sewage sludge. The *A. mangium* fertilization increases the populations of different insect and spider groups on this plant.

## 1. Introduction

Sewage sludge is the residual and semi-solid material produced as a by-product during sewage treatment of industrial or municipal waste [1]. It is rich in organic matter and nutrients with potential for reuse as a fertilizer and media for seedling production after processes of its stabilization such as anaerobic digestion and composting [2]. Crops cultivated using stabilized sewage sludge include the Japanese mustard spinach, *Brassica rapa* L. var. *perviridis* (Brassicales: Brassicaceae) in Japan [3]. Sewage sludge can be re-used as fertilizer in forest plantations, degraded area under recovery process and in agriculture, reducing production costs and environmental risks [4–6]. The quality of dried and pasteurized sewage sludge is classified as class A by the USA [7]. A treated sewage sludge from the 'Estação de Tratamento de Esgoto (ETE)' in the municipality of Juramento, Minas Gerais State, Brazil had no helminth eggs and protozoan cysts, and did not increase the heavy metal contents in grains of maize, *Zea mays* L. (Poales: Poaceae), and cowpea, *Vigna unguiculata* (L.) Walp. (Fabales: Fabaceae) [8].

*Mangium*, *Acacia mangium* Willd. (Fabales: Fabaceae), is native to northeastern Queensland in Australia, the Western Province of Papua New Guinea, Papua and the eastern Maluku Islands [9]. It is a fast growing, hardy and pioneering plant with nitrifying potential, which makes it suitable for degraded area recovering [10]. This plant is used to restoring wastelands created by open-pit gold mining in Colombia [11]. The high fix atmospheric nitrogen gas fixation by this plant in symbiosis with diazotrophic archaea and bacteria increases biomass productivity and nutrient inflow via litter, favouring ecological succession [12]. These characteristics and the high *A. mangium* adaptability to acidic, infertile and flooding-prone soils increase its potential to recover degraded areas [13,14].

Insect (Insecta) diversity, with known function, population or status of these organisms, is an indicator (i.e. bioindicator) that can reveal the qualitative status of the degraded area recovery, responding rapidly to environmental changes [15]. Coleoptera, Lepidoptera and Hymenoptera, with a large number of families and described species (around 400 000; 180 000 and 150 000 respectively), are widely used as bioindicators around the world [16–18]. Plant chemical composition and development (=age) affect the diversity of phytophagous insects and their natural enemies (insects and spiders (Araneae)) and, therefore, serve as nutritional and chemical defence indexes for plants [19,20]. Sewage sludge applied as a boosting material in crops increases the soil organic matter content, besides being rich in macronutrients such as calcium, magnesium, nitrogen and phosphorus, and micronutrients such as copper and zinc [21], favouring plant growth and development and interaction with insect ecology processes.

Hypotheses tested interactions among plant, dehydrated sewage sludge, degraded area and arthropod predatory organisms: plants fertilized with dehydrated sewage sludge have larger crown and litter formation [22,23]. More complex host individuals—larger trees—(i.e. biogeographic island theory (BGI)) support a higher pest insect abundance and diversity owing to better food availability [24,25] and, consequently, more arthropod natural enemies [26]. The tree canopy is a small-scale BGI and an example to test these hypotheses [27]. BGI predicts that extinction rates are higher in smaller islands because they cannot stand high organism populations with the rarest species being more vulnerable to extinction [28]. BGI considers the history of the biological processes such as colonization, speciation and extinction to explain species distribution patterns [29]. Smaller trees are likely to support small populations.

The objective of this study was to evaluate plant biomass production, soil cover by plants and litter and diversity and abundance of chewing and pollinator insects and arthropod predators on *A. mangium* plants, over 24 months, fertilized with dehydrated sewage sludge in a degraded area. The hypotheses tested were that *A. mangium* plants fertilized with dehydrated sewage sludge have larger canopy forming more litter, helping in the degraded area recovery (i); fertilized plants were bigger (>BGI) and had greater abundance of chewing (ii) and pollinator (iii) insects, and arthropod predators were more numerous on larger plants (iv).

## 2. Material and methods

### 2.1. Experimental site

The study was carried out in a degraded area at the 'Instituto de Ciências Agrárias (ICA)' of the 'Universidade Federal de Minas Gerais (UFMG)', municipality of Montes Claros, Minas Gerais State, Brazil (latitude 16°51' S × longitude 44°55' W, altitude 943 m) from March 2015 to February 2017 (24 months; arthropod collection period). The area was defined as degraded owing to soil losses and changes in soil chemistry or hydrology [30,31]. The climate of this region is Aw: tropical savannah, with dry winter and rainy summer, according to the Köppen classification [32]. The soil type is litolic neosoil [33] with average texture, total sand = 42.0 dag Kg<sup>-1</sup>, silt = 36.0 dag Kg<sup>-1</sup>, clay = 22.0 dag Kg<sup>-1</sup>, pH-H<sub>2</sub>O = 5.0, organic matter = 4.4 dag Kg<sup>-1</sup>, P = 1.5 mg dm<sup>-3</sup>, K = 92.0 mg dm<sup>-3</sup>, Ca = 1.9 cmol<sub>c</sub> dm<sup>-3</sup>, Mg = 0.8 cmol<sub>c</sub> dm<sup>-3</sup>, Al = 2.4 cmol<sub>c</sub> dm<sup>-3</sup>, H + Al = 6.7 cmol<sub>c</sub> dm<sup>-3</sup>, cation-exchange capacity (CEC) = 5.3 cmol<sub>c</sub> dm<sup>-3</sup> and CEC at natural pH 7.0 = 9.6 cmol<sub>c</sub> dm<sup>-3</sup> after soil chemical and physical analysis carried out in 2014 in a laboratory using standard international protocols [34].

### 2.2. Experimental design

*Acacia mangium* seedlings were produced from seeds of around 5-year old trees grown at the ICA/UFMG campus. Seeds were dried, dormancy-broken and treated with recommended bactericides/fungicides before sowing following standard protocol used for *Acacia* (= *Vachellia* Wight & Arn.) *farnesiana* (L.) Willd. in Brazil [35]. Seeds were sown in 8 × 12 cm plastic polybags (a seed per plastic polybag) in a nursery with its ruff covered using black shed net, with media mixing with 30% organic compost, 30% clay soil, 30% sand and 10% of reactive natural phosphate (160 g seedling<sup>-1</sup>) in March 2014. The organic compost consisted of three parts, by volume: two parts of debris gardening pruning (≤ 5 cm) and one part of tanned nelore cattle *Bos taurus indicus* L., 1758 (Artiodactyla: Bovidae) manure. The mixture clay soil and sand was treated by a heating process at 80°C for 15 min. The soil pH of the pits was corrected with dolomitic limestone (i.e. an anhydrous carbonate mineral composed of calcium magnesium carbonate), increasing the base saturation to 50% [36]. Fritted trace elements (FTE), gypsum, micronutrients, natural phosphate and potassium chloride were added according to the soil chemical analysis for the Minas Gerais State [37]. Thirty-centimeter tall *A. mangium* seedlings were planted in pits (40 × 40 × 40 cm) spaced 2 m between them, in six parallel lines on flat terrain, spaced 2 m between lines, with four plants with and four without fertilization with dehydrated sewage sludge per line, in September 2014. These seedlings were irrigated twice a week until the beginning of the rainy season using water from a nearby river from when no additional water was provided. The plants were pruned using a razor sterilized with a solution of sodium hydroxide + sodium hypochlorite, when their branches reached 5 cm long, eliminating the additional shoots (i.e. others different from the leader shoot) and branches up to one-third of crown height, leaving only the leader shoot and lateral branches up to two-thirds of the crown height. The pruned parts of each plant (branches and shoots) were left between their respective planting lines. The experimental design was completely randomized in blocks with two treatments (20 l of dehydrated sewage sludge per pit or no dehydrated sewage sludge) and 24 replications with one plant each. The 20 l of dehydrated sewage sludge was placed in a single dose per pit at planting.

Dehydrated sewage sludge (5% moisture content) was collected at the sewage treatment plant—'Estação de Tratamento de Esgoto (ETE)' in the municipality of Juramento, Minas Gerais State, Brazil, about 40 km from the *A. mangium* experimental site. The ETE is operated by the Minas Gerais Sanitation Company S.A.—'Companhia de Saneamento de Minas Gerais S.A. (COPASA)' with capacity to treat 217 m<sup>3</sup> sewage sludge d<sup>-1</sup>. The efficiency of the system in terms of removal of organic matter is higher than 90%. The sewage sludge goes through a solarization process in coarse sand tanks for three months in the ETE reducing the thermotolerant coliform bacteria to a level accepted by the National Council for the Environment—'Conselho Nacional do Meio Ambiente (CONAMA)' (Resolution N° 375) of the Ministry of the Environment—'Ministério do Meio Ambiente' of Brazil for use in agriculture, which is less than 10<sup>3</sup> most likely number g<sup>-1</sup> of total solids. The main chemical and biological characteristics of the dehydrated sewage sludge of this company were pH-H<sub>2</sub>O = 4.40, N = 10.4 mg Kg<sup>-1</sup>, P = 2.9 mg Kg<sup>-1</sup>, K = 5.8 mg Kg<sup>-1</sup>, Cd = 0.1 µg g<sup>-1</sup>, Pb = 56.9 µg g<sup>-1</sup>, Cr = 46.7 µg g<sup>-1</sup> and faecal coliforms = 4.35 most likely number g<sup>-1</sup> after analysis carried out in a laboratory [8].

### 2.3. Plant mass production and soil coverage

Leaves per branch, branches per plant, numbers and the percentage of soil cover by litter, grass and herbaceous plants were evaluated visually per month and plot (1.0 m<sup>2</sup>), in the crown projection area of the 48 *A. mangium* plants.

### 2.4. Insects and spiders

Insects and spiders (no multiply counted) were counted by visual observation biweekly on the adaxial and abaxial surfaces of the leaves between 07.00 and 11.00 at the apical, middle and basal parts of the canopy in the northern, southern, eastern and western directions, totaling 12 leaves plant<sup>-1</sup> evaluation<sup>-1</sup> on the 48 *A. mangium* trees of six-month old for 24 months. Insects and spiders were not removed from plants during the evaluation, except those collected for identification. The total sample effort was 27 648 leaves covering the entire plant (vertical and horizontal axes), capturing as many insect and spider species as possible, especially the rarest. Insects and spiders present on the trunk (chest height) were collected, and insect defoliation was evaluated visually by the leaf area losses on a 0–100% scale with 5% increments for removed leaf area [38,39] for the 48 trees per evaluation. At least, three specimens per insect or spider species were captured per collection using aspirator, stored in glass flasks with 70% ethanol or mounted, separated into morphospecies and sent for identification.

### 2.5. Ecological indices

Averages were made by reducing the data to single trees. Ecological indices (diversity, individual abundance and species richness) were calculated for each identified species in the treatments (with or without dehydrated sewage sludge) per tree using the software BIO-DIVERSITY PROFESSIONAL, Version 2 (©1997 The Natural History Museum) [40]. The diversity was calculated using Hill's formula [41,42] and the species richness with Simpson indices [43,44]. The predator (insects and spiders) and prey ratio on *A. mangium* was calculated per tree.

### 2.6. Statistics

Data on leaves per branch, branches per plant, percentages of soil cover by litter, grass and herbaceous plants, predator per prey ratio and defoliation, diversity, abundance and richness of chewing, defoliator and pollinator insect species, and arthropod predators (see the electronic supplementary material) were submitted to the non-parametric statistical hypothesis, Wilcoxon signed-rank test ( $p < 0.05$ ) [45] using the statistical analysis program 'Sistema para Análises Estatísticas e Genéticas (SAEG)', version 9.1 [46] supplied by the 'Universidade Federal de Viçosa'.

The Spearman correlation matrix, among the most significant characteristics, was calculated. The matrices were submitted to correlation networks [47]. The edge thickness was controlled by applying a cut-of-value 0.26 (from which the Spearman correlation became significant, meaning that only edges with  $|r_{ij}| \geq 0.26$  were highlighted). These analyses were performed using the R software version 3.4.1 by R Core Team [48]. The correlation network procedure was performed using the package qgraph [47].


## 3. Results

### 3.1. Leaves per branch, branches per plant, leaves per tree, percentages of defoliation and soil cover

Leaves per branch and branches per plant, percentages of defoliation by chewing insects and soil cover (litter, grasses and herbaceous plants) were higher for *A. mangium* plants fertilized with dehydrated sewage sludge than for those without fertilization, but no effect was observed on the predator per prey ratio (table 1). The increase in the number of leaves per tree reduced the predator per prey ratio (figure 1).

### 3.2. Biodiversity and richness indexes

The biodiversity and richness indexes for chewing and pollinator insects, and spiders only were similar for *A. mangium* plants fertilized or not with dehydrated sewage sludge. On the other hand, the abundance of chewing (greater than 10 times) and pollinator (greater than 2 times) insects, and total



**Figure 1.** Estimated network structures based on Spearman correlation ( $p < 0.05$ ) generated for abundance (abund.) of chewing, pollinator and predator insects, predator spiders, species richness (SR) of chewing insects, defoliation (%), ratio predator per prey, numbers of *Nasutitermes* sp. (Blattodea: Termitidae), Oxyopidae (Araneae), Tettigoniidae (Orthoptera) and *Tropidacris collaris* (Orthoptera: Romaleidae) on *Acacia mangium* (Fabales: Fabaceae) per tree  $n = 48$ .

**Table 1.** Numbers of total leaves per branch and branches per plant, defoliation, predator per prey ratio and soil cover (mean  $\pm$  s.e.) of *Acacia mangium* (Fabales: Fabaceae) per tree with or without dehydrated sewage sludge. ( $n = 24$  per treatment, VT = value of the test.)

	sewage sludge		Wilcoxon test	
	without	with	VT	$p$
leaves per branch	22.00 $\pm$ 0.69	33.71 $\pm$ 1.10	4.9	0.00
branches per plant	17.73 $\pm$ 0.56	41.26 $\pm$ 0.90	5.7	0.00
defoliation (%)	4.35 $\pm$ 0.25	6.28 $\pm$ 0.16	5.3	0.00
predator per prey ratio	10.37 $\pm$ 2.94	6.34 $\pm$ 1.49	1.4	0.08
soil cover (%)	8.43 $\pm$ 0.54	27.47 $\pm$ 1.10	5.6	0.00

predators (greater than 1.4 times) were higher on plants fertilized with dehydrated sewage sludge. Moreover, the treatments did not affect the spider ecological indices, but plants fertilized had more biodiversity of total predators (table 2). The increase in the number of leaves per tree increased the abundance and richness of chewing insects besides the predators on *A. mangium* plants. Higher species richness of chewing insects resulted in bigger predator abundances, including spiders (figure 1).

### 3.3. Arthropods

The termite *Nasutitermes* sp. (Blattodea: Termitidae) and the Neotropical stick grasshopper *Cephalocoema* sp. (Orthoptera: Proscopiidae) numbers were higher on *A. mangium* trunks fertilized with dehydrated sewage sludge and without fertilization, respectively, while the number of all other chewing insect species was similar ( $p < 0.05$ ) between treatments. The Orthoptera chewers, the large South American grasshopper *Tropidacris collaris* Stoll, 1813 (Romaleidae) and the katydid Tettigoniidae and the Coleoptera *Lordops* sp. (Curculionidae) and *Stereoma anchoralis* Lacordaire, 1848 (Chrysomelidae) stood out in relation to the other chewing insects owing to their greater abundance on *A. mangium* plants, with or without dehydrated sewage sludge fertilization (table 3). The increase in the number of leaves per tree increased that of *Nasutitermes* sp. (figure 1).

The stingless bee, *Trigona spinipes* F., 1793 (Hymenoptera: Apidae) numbers were higher on *A. mangium* plants fertilized with dehydrated sewage sludge while those of the European honeybee, *Apis mellifera* L., 1758 and the stingless bee, *Tetragonisca angustula* Latreille, 1811 (Hymenoptera: Apidae) were similar ( $p < 0.05$ ) between treatments, yet both had lower abundance than the first species (table 4).

**Table 2.** Diversity index (DI), species richness (SR) and abundance (abund.) of chewing and pollinator insects, total predators and spiders on *Acacia mangium* (Fabales: Fabaceae) per tree (mean  $\pm$  SE) with or without dehydrated sewage sludge. ( $n = 24$  per treatment, VT = value of the test.)

	sewage sludge		Wilcoxon test	
	without	with	VT	$p$
DI chewing insects	7.42 $\pm$ 1.35	4.66 $\pm$ 0.76	1.2	0.12
SR chewing insects	2.88 $\pm$ 0.40	3.17 $\pm$ 0.33	0.9	0.19
abund. chewing insects	3.96 $\pm$ 0.59	40.75 $\pm$ 17.80	2.3	0.01
DI pollinators	3.25 $\pm$ 0.37	2.74 $\pm$ 0.24	0.7	0.26
SR pollinators	1.25 $\pm$ 0.21	1.67 $\pm$ 0.14	0.2	0.43
abund. pollinators	2.96 $\pm$ 0.58	6.42 $\pm$ 1.28	2.2	0.02
DI total predators	14.96 $\pm$ 1.46	10.42 $\pm$ 1.32	2.3	0.01
SR total predators	7.88 $\pm$ 0.52	8.33 $\pm$ 0.52	0.1	0.45
abund. total predators	56.83 $\pm$ 17.59	77.92 $\pm$ 11.52	2.1	0.02
DI spiders	4.31 $\pm$ 0.38	3.59 $\pm$ 0.49	1.6	0.06
SR spiders	2.04 $\pm$ 0.19	1.75 $\pm$ 0.20	1.4	0.08
abund. spiders	2.50 $\pm$ 0.60	3.17 $\pm$ 0.39	1.0	0.17

The wasp *Polybia* sp. (Hymenoptera: Vespidae), the jumping spider Salticidae (Araneae) and the praying mantis *Mantis religiosa* (Linnaeus, 1758) (Mantodea: Mantidae) numbers were higher ( $p < 0.05$ ) on *A. mangium* plants with or without, respectively, dehydrated sewage sludge, while those of all other predators insect and spider species was similar ( $p > 0.05$ ) between treatments (table 4). The increase in the abundance of pollinators, Tettigoniidae and *T. collaris* individuals increased that of predators as well as the species richness of chewing insects resulted in higher numbers of the lynx spider Oxyopidae (Araneae) (figure 1).

## 4. Discussion

The highest abundance of chewing, pollinator and predator insects on *A. mangium* fertilized with dehydrated sewage sludge and soil covered by litter were owing to the better development of these plants (e.g. >leaves per tree), similar to that of the flooded gum *Eucalyptus grandis* W. Hill ex Maiden (Myrtales: Myrtaceae) [22] and their higher nitrogen levels in a dehydrated sewage sludge [8] obtained from the same ETE of the current study.

The litter cover in the crown projection area of *A. mangium* plants fertilized with dehydrated sewage sludge resulted from the higher number of leaves and branches produced by these plants compared to the non-fertilized ones, important for reducing laminar erosion and increasing soil fertility [49,50], confirming the first hypothesis in which fertilized plants are better in the recovery process of degraded areas. Dehydrated sewage sludge is rich in organic matter and macronutrients such as nitrogen and phosphorus, besides micronutrients such as copper and zinc, favouring tree growth and development [51,52]. The recovery of degraded areas is slow, but the use of *A. mangium* fertilized with dehydrated sewage sludge is promising, because of its fast growth and development, efficient fix atmospheric nitrogen gas fixation, potential to improve soil quality and widespread use [24]. This agrees with the positive impact of dehydrated sewage sludge in the development of the Brazilian pine, *Araucaria angustifolia* (Bertol) Kuntze (Pinales: Araucariaceae); Argentine cedar, *Cedrela fissilis* Vell. (Sapindales: Meliaceae); *E. grandis*; *Lafloensia pacari* St.-Hil. (Myrtales: Lythraceae), and *Senna spectabilis* (DC.) Irwin & Barneby (Fabales: Fabaceae) [22,23]. Dehydrated sewage sludge sanitized with neen, *Azadirachta indica* A. Juss. (Sapindales: Meliaceae) or with lime (a calcium-containing inorganic mineral) without this plant or in composition with *Ipomoea* sp. (Solanales: Convolvulaceae) compost with or without rock phosphate were evaluated. Other treatments included dehydrated sewage sludge with rock phosphate incorporated in the soil sanitized or not with *A. indica* or lime, and this fertilization with *Ipomoea* sp. compost with rock phosphate is incorporated in the soil. The density of pathogen (i.e. helminths and

**Table 3.** Total numbers of chewing insects per *Acacia mangium* (Fabales: Fabaceae) per tree (mean  $\pm$  s.e.) with or without dehydrated sewage sludge. ( $n = 24$  per treatment, VT = value of the test.)

order: family	species	sewage sludge		Wilcoxon test	
		without	with	VT	<i>p</i>
Coleoptera:					
Buprestidae	<i>Psiloptera</i> sp.	0.04 $\pm$ 0.04	0.00 $\pm$ 0.00	1.0	0.16
Cerambycidae	non-identified	0.08 $\pm$ 0.05	0.00 $\pm$ 0.00	1.4	0.08
Chrysomelidae	<i>Alagoasa</i> sp.	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
	<i>Cerotoma</i> sp.	0.17 $\pm$ 0.07	0.21 $\pm$ 0.08	0.4	0.36
	<i>Diabrotica speciosa</i> Germar	0.08 $\pm$ 0.05	0.25 $\pm$ 0.09	1.5	0.06
	<i>Disonycha brasiliensis</i> Lima	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
	<i>Eumolpus</i> sp.	0.04 $\pm$ 0.04	0.13 $\pm$ 0.06	1.0	0.15
	<i>Lamprosoma</i> sp.	0.04 $\pm$ 0.04	0.00 $\pm$ 0.00	1.0	0.16
	<i>Parasyphraea</i> sp.	0.25 $\pm$ 0.12	0.46 $\pm$ 0.19	1.0	0.17
	<i>Stereoma anchoralis</i> Lacordaire	0.46 $\pm$ 0.21	0.21 $\pm$ 0.13	0.8	0.20
	<i>Walterianella</i> sp.	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
	<i>Wanderbiltiana</i> sp.	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
Curculionidae	<i>Lordops</i> sp.	0.00 $\pm$ 0.00	0.46 $\pm$ 0.41	1.4	0.08
	non-identified	0.04 $\pm$ 0.04	0.08 $\pm$ 0.05	0.6	0.28
Tenebrionidae	Alleculinae	0.04 $\pm$ 0.04	0.00 $\pm$ 0.00	1.0	0.16
	<i>Epitragus</i> sp.	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
Blattodea:					
Termitidae	<i>Nasutitermes</i> sp. <sup>a</sup>	0.00 $\pm$ 0.00	36.37 $\pm$ 14.73	3.3	0.00
Lepidoptera:					
non-identified	non-identified	0.33 $\pm$ 0.13	0.21 $\pm$ 0.10	0.7	0.23
Orthoptera:					
Gryllidae	non-identified	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
Tettigoniidae	non-identified	1.00 $\pm$ 0.19	0.75 $\pm$ 0.18	1.0	0.16
Proscopiidae	<i>Cephalocoema</i> sp.	0.13 $\pm$ 0.06	0.00 $\pm$ 0.00	1.8	0.04
Romaleidae	<i>Tropidacris collaris</i> Stoll	1.00 $\pm$ 0.19	1.33 $\pm$ 0.26	0.7	0.25
Phasmatodea:					
Phasmidae	<i>Phibalosoma phyllinum</i> Gray	0.00 $\pm$ 0.00	0.04 $\pm$ 0.04	1.0	0.16

<sup>a</sup>Observed on *A. mangium* trunk.

protozoans) in these treatments was low and similar with soils without fertilization or with liming and chemical fertilization [8]. These treatments did not surpass the maximum limits of annual addition and the permissible maximum levels of heavy metal concentrations in the soils, but the concentrations of lead in *Z. mays* and *V. unguiculata* grains reached values above the limits permitted for agricultural products, regardless of the addition of sewage sludge in the soil [8].

The greater abundance of chewing insects and defoliation on *A. mangium* plants fertilized with dehydrated sewage sludge is probably owing to the greater number of leaves serving as a better food source and quality for insects. This confirms the second hypothesis that the diversity and abundance of herbivorous insects and their predators are usually higher and with higher increase of chewing insects than predators (e.g. >leaves  $\leq$  predator per prey ratio) on trees with higher leaf mass [25,53,54]. These trees function as a BGI, but with a higher chance of rare species extinction on those with lower leaf mass [25,28,55]. In addition, the quantity of free amino acids and proteins is superior in plants with higher nitrogen fertilization, favouring herbivorous insects [56]. Interactions between insects and *Acacia* species plants show the potential of this plant to increase the biodiversity and recover

**Table 4.** Total numbers of spiders, and insect predators and pollinators per *Acacia mangium* (Fabales: Fabaceae) per tree (mean  $\pm$  s.e.) with or without dehydrated sewage sludge. ( $n = 24$  per treatment, VT = value of the test.)

order: family	species	sewage sludge		Wilcoxon test	
		without	with	VT	$p$
Araneae:					
Anyphaenidae	<i>Teudis</i> sp.	0.04 $\pm$ 0.04	0.00 $\pm$ 0.00	1.0	0.16
Araneidae	non-identified	1.21 $\pm$ 0.22	0.83 $\pm$ 0.19	1.3	0.10
Oxyopidae	non-identified	0.50 $\pm$ 0.17	0.75 $\pm$ 0.19	1.2	0.11
	<i>Oxyopes salticus</i> Hentz	0.17 $\pm$ 0.07	0.08 $\pm$ 0.05	0.9	0.19
Salticidae	non-identified	1.08 $\pm$ 0.33	0.54 $\pm$ 0.19	1.8	0.04
	<i>Uspachus</i> sp.	0.04 $\pm$ 0.04	0.08 $\pm$ 0.08	0.0	0.49
Sparassidae	<i>Quemedice</i> sp.	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
Tetragnathidae	<i>Leucauge</i> sp.	0.04 $\pm$ 0.04	0.08 $\pm$ 0.08	1.0	0.16
Thomisidae	<i>Aphantochilus rogersi</i> Cambridge	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
	<i>Tmarus</i> sp.	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	0.0	0.50
Hemiptera:					
Pentatomidae	<i>Podisus</i> sp.	0.04 $\pm$ 0.04	0.13 $\pm$ 0.06	1.0	0.15
Hymenoptera:					
Apidae	<i>Apis mellifera</i> L.	0.50 $\pm$ 0.19	0.38 $\pm$ 0.15	0.2	0.44
	<i>Tetragonisca angustula</i> Latreille	1.17 $\pm$ 0.26	1.38 $\pm$ 0.24	0.8	0.22
	<i>Trigona spinipes</i> F.	1.29 $\pm$ 0.39	4.67 $\pm$ 1.31	2.2	0.02
Vespidae	<i>Polybia</i> sp.	0.60 $\pm$ 0.26	3.75 $\pm$ 2.71	2.1	0.02
Mantodea:					
Mantidae	<i>Mantis religiosa</i> L.	0.25 $\pm$ 0.09	0.04 $\pm$ 0.04	2.0	0.02

degraded areas around the world [57–60]. The dehydrated sewage sludge as a biofertilizer improved macrofauna recovery, including the scarab beetles Scarabaeidae (Coleoptera) larvae and adults in degraded soils of the Cerrado (Brazilian savannah) type biome area [55].

The presence of *Nasutitermes* sp., as the most abundant insect on *A. mangium* plant trunks fertilized with dehydrated sewage sludge may be owing to the organic matter richness of this fertilizer [8,51] and the higher litter production by this plant (e.g. >leaves  $\geq$  *Nasutitermes* sp.). This insect can damage living or dead trees and processed wood, including root systems, although they caused galleries in the trunks without damaging or causing plant death [61]. Damage by *Lordops* sp., *S. anchoralis*, *T. collaris* and Tettigoniidae on *A. mangium* leaves and their greater abundance compared to that of other chewing insects is worrying. *Tropidacris collaris* damaged the swamp she-oak, *Casuarina glauca* Sieber (Fagales: Casuarinaceae) and white leadtree, *Leucaena leucocephala* (Lam.) de Wit (Fabales: Fabaceae) [62,63]. *Merocidius intermedius* Brunner Von Wattenwyl, 1895 (Orthoptera: Tettigoniidae) damaged grasses and banana *Musa* spp. fruits (Zingiberales: Musaceae) [64] and *Lordops* sp. defoliated the diesel tree, *Copaifera langsdorffii* Desf. (Fabales: Fabaceae) [65], but there is no reports of *S. anchoralis* damaging commercial plants.

The number of pollinating insects being two times higher on *A. mangium* plants fertilized with dehydrated sewage sludge is probably owing to their larger canopy size, higher number of flowers and supporting a greater insect numbers [25,54], including pollinators, confirming the third hypothesis: greater BGI greater pollinating insects. In addition, nitrogen fertilization via dehydrated sewage sludge may have increased the pollen and/or nectar production and quality (more amino acids and protein) in *A. mangium* flowers, increasing pollinator attractiveness as observed for the higher attraction of *Nicotiana* L. (Solanales: Solanaceae) species flowers with better quality of nectar sugars and amino acids by different groups of pollinators (e.g. bats (Chiroptera), hummingbirds (Trochilidae) or moths (Lepidoptera)) in Wuppertal, Germany [66] and floral pollens with better quality by bumblebees, *Bombus* Latreille, 1802 (Hymenoptera: Apidae) and honeybees, *Aphis* species in Newcastle, United Kingdom [67]. The greater *T. spinipes* pollinator abundance on *A. mangium* plants, especially on those



fertilized with dehydrated sewage sludge may be of low importance, because it can reduce pollination as reported for Cucurbitaceae (Cucurbitales) owing to insufficient pollen transportation (small body size) and/or chasing other pollinators, such as *A. mellifera* and *T. angustula*, by flying in flocks and with aggressive behaviour [68]. In addition, *T. spinipes* damages shoots and plant growth regions by removing fibres to construct their nests, as reported on *A. mangium* and *L. leucocephala*, that also had their leaves and shoots damaged [63,69].

The greater abundance of predator insects and spiders, on *A. mangium* plants fertilized with dehydrated sewage sludge, is probably owing to the higher number of chewing and pollinator insects on the plants (larger trees), that is, these predators followed their prey [70], confirming the fourth hypothesis: greater BGI greater predators. In general, the number of predator insect and spider species did not differ between *A. mangium* plants with or without dehydrated sewage sludge fertilization, but the number of spider species (30% higher on fertilized plants) and the abundance of the predatory wasp *Polybia* sp. were higher on fertilized plants. Spider predators reduced insect damage, mainly from defoliators, such as spiders in many agroecosystems in the USA [71], wolf (Araneae: Lycosidae) and sheet weaver (Araneae: Linyphiidae) spiders in winter barley, *Hordeum vulgare* L. (Poales: Poaceae) fields situated in differently structured landscapes in Uppsala, Sweden [72], wandering spiders (Araneae: Ctenidae) in agroecosystems in Italy [73] and spiders in pequi, *Caryocar brasiliense* Cambess. (Malpighiales: Caryocaraceae) trees in Minas Gerais State, Brazil [26]. Predatory wasps (Vespidae) are important natural enemies in agricultural systems such as *Brassica campestris* L. and kale, *Brassica oleracea* L. var. *acephala* DC., Arabian coffee, *Coffea arabica* L. (Gentianales: Rubiaceae) and tomato, *Solanum lycopersicon* L. (Solanales: Solanaceae), preying mainly on caterpillars and leaf miners (Lepidoptera) in several regions of Brazil [74–77]. Sewage sludge increased the ground beetle Carabidae (Coleoptera) species richness in the area of Oxford, USA [78].

## 5. Conclusion

To summarize, the larger *A. mangium* crown (>BGI) fertilized with dehydrated sewage sludge increases soil cover (e.g. litter) and the abundance of chewing (>defoliation) and pollinator insects and arthropod predators, showing that this plant is adequate for recovering degraded areas using this fertilization. The presence of *Nasutitermes* sp. on *A. mangium* plant trunks fertilized with dehydrated sewage sludge may be owing to the organic matter richness of this fertilizer and the higher litter production by this plant (e.g. >leaves  $\geq$  *Nasutitermes* sp.), but without damaging or causing plant death. On the other hand, *Lordops* sp., *T. collaris* and Tettigoniidae damaged leaves of *A. mangium* and this is worrying because these insects are pests in other economically important crops. The greater *T. spinipes* pollinator abundance on *A. mangium* plants is a problem owing to this insect damaged their shoots and plant growth regions.

**Ethics.** No specific permits are required to plant *Acacia mangium* in Brazil. The laboratory and field studies did not involve endangered or protected species.

**Data accessibility.** All data generated or analysed during this study are included in this manuscript.

**Authors' contributions.** The study was conceived and designed by J.L.S., G.L.D.L. and R.A.S., data were collected by J.L.S., W. de S.T. and F.W.S.S. with support from G.L.D.L. and R.A.S., data analysis was done by G.L.D.L. and A.M.A., figures, tables and manuscript preparation was done by J.L.S., G.L.D.L., W. de S.T., A.M.A., J.E.S. and J.C.Z. and all authors contributed to revisions and approved the final manuscript.

**Competing interests.** The authors declare that they have no conflict of interest.

**Funding.** The study was financially supported by the following Brazilian agencies 'Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)', 'Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES-Finance Code 001)', 'Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)' and 'Programa Cooperativo sobre Proteção Florestal (PROTEF)' of the 'Instituto de Pesquisas e Estudos Florestais (IPEF)'.

**Acknowledgements.** We would like to thank Dr Antônio Domingos Brescovit (Instituto Butantan, São Paulo State, Brazil—Arachnida) and Dr Ayr de Moura Bello (Fundação Oswaldo Cruz, Rio de Janeiro State, Brazil—Coleoptera) for arthropod species identification. The voucher numbers for insects are 1595/02 and 1597/02 (Centro de Estudos Faunísticos e Ambientais, Universidade Federal do Paraná, Curitiba, Paraná State, Brazil) and that of spiders is IBSP 36921–36924 (Instituto Butantan, São Paulo State, Brazil). Dr Phillip John Villani (University of Melbourne, Australia) revised and corrected the English language used in this manuscript.

## References

1. Antonkiewicz J, Baran A, Pelka R, Wisła-Świder A, Nowak E, Konieczka P. 2019 A mixture of cellulose production waste with municipal sewage as new material for an ecological management of wastes. *Ecotox. Environ. Safe.* **169**, 607–614. (doi:10.1016/j.ecoenv.2018.11.070)

2. Frišták V, Pipiška M, Soja G. 2018 Pyrolysis treatment of sewage sludge: a promising way to produce phosphorus fertilizer. *J. Clean. Prod.* **172**, 1772–1778. (doi:10.1016/j.jclepro.2017.12.015)
3. Nguyen T-B, Shima K. 2019 Composting of sewage sludge with a simple aeration method and its utilization as a soil fertilizer. *Environ. Manage.* **63**, 455–465. (doi:10.1007/s00267-017-0963-8)
4. Kimberley MO, Wang H, Wilks PJ, Fisher CR, Magesan GN. 2004 Economic analysis of growth response from a pine plantation forest applied with biosolids. *Forest Ecol. Manag.* **189**, 345–351. (doi:10.1016/j.foreco.2003.09.003)
5. Caldeira MVW, Favalessa M, de Gonçalves EO, Delarmelina WM, Santos FEV, Vieira M. 2014 Based substrates of sewage sludge for the production of seedlings of *Acacia mangium*. *Wild. Comum. Sci.* **5**, 34–43. (doi:10.14295/cs.v5i1.309)
6. Silva MI, Mackowiak C, Minogue P, Reis AF, de Moline EFV. 2016 Potential impacts of using sewage sludge biochar on the growth of plant forest seedlings. *Ciênc. Rural* **47**, e20160064. (doi:10.1590/0103-8478cr20160064)
7. Al-Gheethi AA, Efaq AN, Bala JD, Norli I, Abdel-Monem MO, Ab. Kadir MO. 2018 Removal of pathogenic bacteria from sewage-treated effluent and biosolids for agricultural purposes. *Appl. Water Sci.* **8**, Article no. 74. (doi:10.1007/s13201-018-0698-6)
8. Nogueira TAR, Sampaio RA, Fonseca IM, Ferreira CS, Santos SE, Ferreira LC, Gomes E, Fernandes LA. 2007 Heavy metals and pathogens in maize – cowpea intercropping system fertilized with sewage sludge. *Rev. Bras. Eng. Agr. Amb.* **11**, 331–338. (doi:10.1590/S1415-43662007000300014)
9. Koutika L-S, Richardson DM. 2019 *Acacia mangium* Willd.: benefits and threats associated with its increasing use around the world. *For. Ecosyst.* **6**, Article no. 2. (doi:10.1186/s40663-019-0159-1)
10. Caldeira MVW, Favalessa M, Delarmelina WM, de Gonçalves EO, dos Moura RRS. 2018 Sewage sludge assessment on growth of *Acacia mangium* seedlings by principal components analysis and orthogonal contrasts. *J. Plant Nutr.* **41**, 1303–1311. (doi:10.1080/01904167.2018.1450421)
11. Thomas E. 2014 Forest devastated by mining is reborn. *Nature* **511**, 155. (doi:10.1038/511155d)
12. Paula RR et al. 2018 Nitrogen fixation rate of *Acacia mangium* Willd at mid rotation in Brazil is higher in mixed plantations with *Eucalyptus grandis* Hill ex Maiden than in monocultures. *Ann. Forest Sci.* **75**, 14. (doi:10.1007/s13395-018-0695-9)
13. Baliero FC, Dias LE, Franco AA, Campello EFC, de Faria SM. 2004 Nutrient accumulation in the aboveground biomass, in the litter layer and phylloides decomposition of *Acacia mangium* Willd. *Ciênc. Florest.* **14**, 59–65. (doi:10.5902/198050981781)
14. Wang F, Zhu W, Zou B, Neher DA, Fu S, Xia H, Li Z. 2013 Seedling growth and soil nutrient availability in exotic and native tree species: implications for afforestation in southern China. *Plant Soil* **364**, 207–218. (doi:10.1007/s11104-012-1353-x)
15. Santos MS, Louzada JNC, Dias N, Zanetti R, Delabie JHC, Nascimento IC. 2006 Litter ants richness (Hymenoptera, Formicidae) in remnants of a semi-deciduous forest in the Atlantic rain forest, Alto do Rio Grande region, Minas Gerais, Brazil. *Iheringia Sér. Zool.* **96**, 95–101. (doi:10.1590/S0073-47212006000100017)
16. Osborn F, Goitia W, Cabrera M, Jaffé K. 1999 Ants, plants and butterflies as diversity indicators: comparison between strata at six forest sites in Venezuela. *Stud. Neotrop. Fauna E.* **34**, 59–64. (doi:10.1076/snfe.34.1.59.8918)
17. Davis AJ, Holloway JD, Huijbregts H, Kriksen J, Kirk-Springgs AH, Sutton SL. 2001 Dung beetles as indicators of change in the forests of northern Borneo. *J. Appl. Ecol.* **38**, 593–616. (doi:10.1046/j.1365-2664.2001.00619.x)
18. Burgio G et al. 2015 The influence of vegetation and landscape structural connectivity on butterflies (Lepidoptera: Papilionoidea and Hesperioidea), carabids (Coleoptera: Carabidae), syrphids (Diptera: Syrphidae), and sawflies (Hymenoptera: Symphyta) in Northern Italy farmland. *Environ. Entomol.* **44**, 1299–1307. (doi:10.1093/ee/nvv105)
19. Bowers MD, Stamp NE. 1993 Effects of plant-age, genotype, and herbivory on plantago performance and chemistry. *Ecology* **74**, 1778–1791. (doi:10.2307/1939936)
20. Coley PD, Barone JA. 1996 Herbivory and plant defenses in tropical forests. *Annu. Rev. Ecol. Syst.* **27**, 305–335. (doi:10.1146/annurev.ecolsys.27.1.305)
21. de Tavares LS, Scaramuzza WLMP, dos Weber OLS, de Valadão FA, Maas KDB. 2013 Tanning sludge and its influence on the production of seedlings of paricá (*Schizolobium amazonicum*) and on chemical properties of soil. *Ciênc. Florest.* **23**, 357–368. (doi:10.5902/1980509810547)
22. da Silva PHM, Poggiani F, Laclau JP. 2011 Applying sewage sludge to *Eucalyptus grandis* plantations: effects on biomass production and nutrient cycling through litterfall. *Appl. Environ. Soil Sci.* Article ID 710614. (doi:10.1155/2011/710614)
23. de Abreu AHM, Marzola LB, de Melo LA, dos Leles PSS, Abel ELS, Alonso JM. 2017 Urban solid waste in the production of *Lafloensia pacari* seedlings. *Rev. Bras. Eng. Agr. Amb.* **21**, 83–87. (doi:10.1590/1807-1929/agriambi.v21n2p83-87)
24. Ferreira WC, Botelho SA, Davide AC, Faria JMR. 2007 Growth of the tree stratum in restored area near Rio Grande in the Camargos hydroelectric plant, state of Minas Gerais. *Rev. Árvore* **31**, 177–185. (doi:10.1590/S0100-67622007000100020)
25. Leite GLD, dos Veloso RVS, Zanuncio JC, Azevedo AM, Silva JL, Wilcken CF, Soares MA. 2017 Architectural diversity and galling insects on *Caryocar brasiliense* trees. *Sci. Rep.* **7**, Article number 16677. (doi:10.1038/s41598-017-16954-6)
26. Leite GLD, dos Veloso RVS, Zanuncio JC, Almeida CIME, Ferreira PSF, Fernandes GW, Soares MA. 2012 Habitat complexity and *Caryocar brasiliense* herbivores (Insecta; Arachnida; Araneae). *Fla. Entomol.* **95**, 819–830. (doi:10.1653/024.095.0402)
27. Kitahara M, Fujii K. 1997 An island biogeographical approach to the analysis of butterfly community patterns in newly designed parks. *Res. Popul. Ecol.* **91**, 23–35. (doi:10.1007/BF02765247)
28. Burns KC. 2016 Native exotic richness relationships: a biogeographic approach using turnover in island plant populations. *Ecology* **97**, 2932–2938. (doi:10.1002/ecy.1579)
29. Patiño J et al. 2017 A roadmap for island biology: 50 fundamental questions after 50 years of The Theory of Island Biogeography. *J. Biogeogr.* **44**, 963–983. (doi:10.1111/jbi.12986)
30. Milton SJ, Dean WRJ, du Plessis MA, Siegfried WR. 1994 A conceptual model of arid rangeland degradation. *BioScience* **44**, 70–76. (doi:10.2307/1312204)
31. Whisenant SG. 1999 Repairing damaged wildlands: a process-oriented, landscape-scale approach. *Restor. Ecol.* **9**, 249. (doi:10.1046/j.1526-100x.2001.009002249.x)
32. Silva CL, de Roldão BC, Santos LDT, Hein PRG. 2018 Wood and bark of *Eucalyptus* and *Acacia* in pure and mixed stands. *Foresta Ambient.* **25**, e00081914. (doi:10.1590/2179-8087.081914)
33. Santana PHL, Frazão LA, Santos LDF, Fernandes LA, Sampaio RA. 2016 Soil attributes and production of *Eucalyptus* in monoculture and silvopastoral systems in the north of Minas Gerais. *Brazil. J. Agr. Sci. Technol. B* **6**, 361–370. (doi:10.17265/2161-6264/2016.06.001)
34. Pegoraro RF, Moreira CG, Dias DG, Silveira TC. 2018 Carbon and nitrogen stocks in the soil and humic substances of agricultural crops in the semi-arid region. *Rev. Ciênc. Agron.* **49**, 574–583. (doi:10.5935/1806-6690.20180065)
35. de Melo PAFR, Alves EU, dos Anjos Neto AP, de Medeiros RLS, Ferreira da Júnior DC, Mondego JM, de Leite WS, Bezerra AKD. 2016 Temperature and pre-germinative treatments for overcoming *Acacia farnesiana* (L.) Willd. (Fabaceae) seeds dormancy. *Afr. J. Agric. Res.* **11**, 3548–3553. (doi:10.5897/AJAR2016.11493)
36. Kopittke PM, Menzies NW. 2007 A review of the use of the basic cation saturation ratio and the 'ideal' soil. *Soil Sci. Soc. Am. J.* **71**, 259–265. (doi:10.2136/sssaj2006.0186)
37. Nouvellon Y, Laclau J-P, Epron D, Le Maire G, Bonnefond J-M, Gonçalves JLM, Bouillet J-P. 2012 Production and carbon allocation in monocultures and mixed-species plantations of *Eucalyptus grandis* and *Acacia mangium* in Brazil. *Tree Physiol.* **32**, 680–695. (doi:10.1093/treephys/tps041)
38. Sastawa BM, Lawan M, Maina YT. 2004 Management of insect pests of soybean: effects of sowing date and intercropping on damage and grain yield in the Nigerian Sudan savanna. *Crop Prot.* **23**, 155–161. (doi:10.1016/j.cropro.2003.07.007)
39. Mizumachi E, Mori A, Osawa N, Akiyama R, Tokuchi N. 2006 Shoot development and extension of *Quercus serrata* saplings in response to insect damage and nutrient conditions. *Ann. Bot.* **98**, 219–226. (doi:10.1093/aob/md091)
40. Krebs CJ. 1989 Bray-Curtis cluster analysis. 1. Biodiversity Pro Versão 2. See http://

- biodiversity-pro.software.informer.com/. Accessed on 2 May, 2016.
41. Hill MO. 1973 Diversity and evenness: a unifying notation and its consequences. *Ecology* **54**, 427–432. (doi:10.2307/1934352)
  42. Jost L. 2006 Entropy and diversity. *Oikos* **113**, 363–375. (doi:10.1111/j.2006.0030-1299.14714.x)
  43. Begon M, Harper JL, Townsend CR. 2007 *Ecologia: de indivíduos a ecossistemas*, 4th edn. Porto Alegre: Artmed.
  44. Lazo JA, Valdés NV, Sampaio RA, Leite GLD. 2007 Zoological diversity associated to a silvopastoral system leucaena-guinea grass with different establishment times. *Pesqui. Agropecu. Bras.* **42**, 1667–1674. (doi:10.1590/S0100-204X2007001200001)
  45. Wilcoxon F. 1945 Individual comparisons by ranking methods. *Biometrics Bull.* **1**, 80–83. (doi:10.2307/3001968)
  46. SAEG - Sistema para Análises Estatísticas. 2007 Versão 9.1: Fundação Arthur Bernardes - UFV - Viçosa, 2007. See <http://arquivo.ufv.br/saeg/>. Accessed on 30 June, 2016.
  47. Epskamp S, Cramer AOJ, Waldorp LJ, Schmittmann VD, Borsboom D. 2012 *qgraph*: network visualizations of relationships in psychometric data. *J. Stat. Softw.* **48**, 1–18. (doi:10.18637/jss.v048.i04)
  48. R Core Team. 2014 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. See <http://www.R-project.org/>. Accessed on: 30 June, 2016.
  49. Franco FS, Couto L, de Carvalho AF, Jucksch I, Fernandes Filho EI, Silva E, Meira Neto JAA. 2002 Evaluation of erosion under agroforestry and conventional systems in Zona da Mata de Minas Gerais. *Rev. Árvore* **26**, 751–760. (doi:10.1590/S0100-67622002000600011)
  50. Majid NM, Islam MM, Mathew L. 2002 Heavy metal uptake and translocation by mangium (*Acacia mangium*) from sewage sludge contaminated soil. *Aust. J. Crop Sci.* **6**, 1228–1235.
  51. Jakubus M. 2016 Estimation of phosphorus bioavailability from composted organic wastes. *Chem. Spec. Bioavailab.* **28**, 189–198. (doi:10.1080/09542299.2016.1227687)
  52. Zhang D, Luo W, Li Y, Wang G, Li G. 2018 Performance of co-composting sewage sludge and organic fraction of municipal solid waste at different proportions. *Biores. Technol.* **250**, 853–859. (doi:10.1016/j.biortech.2017.08.136)
  53. Ferrier SM, Price PW. 2004 Oviposition preference and larval performance of a rare bud-galling sawfly (Hymenoptera: Tenthredinidae) on willow in northern Arizona. *Environ. Entomol.* **33**, 700–708. (doi:10.1603/0046-225X-33.3.700)
  54. Espírito-Santo MM, de Neves FS, Andrade-Neto FR, Fernandes GW. 2007 Plant architecture and meristem dynamics as the mechanisms determining the diversity of gall-inducing insects. *Oecologia* **153**, 353–364. (doi:10.1007/s00442-007-0737-8)
  55. Kitamura AE, Alves MC, Suzuki LGAS, Gonzalez AP. 2008 Recovery of a degraded soil with green manure and sewage sludge. *Rev. Bras. Ciênc. Solo* **32**, 405–416. (doi:10.1590/S0100-06832008000100038)
  56. Gao J, Guo H, Sun Y, Ge F. 2018 Differential accumulation of leucine and methionine in red and green pea aphids leads to different fecundity in response to nitrogen fertilization. *Pest Manag. Sci.* **74**, 1779–1789. (doi:10.1002/ps.4875)
  57. Fleming PA, Hofmeyr SD, Nicolson SW. 2007 Role of insects in the pollination of *Acacia nigrescens* (Fabaceae). *S. Afr. J. Bot.* **73**, 49–55. (doi:10.1016/j.sajb.2006.06.010)
  58. McLeish MJ, Chapman TW, Schwarz MP. 2007 Host-driven diversification of gall inducing *Acacia* thrips and the aridification of Australia. *BMC Biol.* **5**, Article no. 3. (doi:10.1186/1741-7007-5-3)
  59. Palmer WA, Lockett CJ, Senaratne KADW, McLennan A. 2007 The introduction and release of *Chiasmia inconspicua* and *C. assimilis* (Lepidoptera: Geometridae) for the biological control of *Acacia nilotica* in Australia. *Biol. Control* **41**, 368–378. (doi:10.1016/j.biocontrol.2007.02.009)
  60. Silva FWS, Leite GLD, Guañabens REM, Sampaio RA, Gusmão CAG, Serrão JE, Zanuncio JC. 2015 Seasonal abundance and diversity of arthropods on *Acacia mangium* (Fabales: Fabaceae) trees as windbreaks in the Cerrado. *Fla. Entomol.* **98**, 170–174. (doi:10.1653/024.098.0129)
  61. Albuquerque AC, Matias GRRS, Couto AAVO, Oliveira MAP, Vasconcellos A. 2014 Urban termites of Recife, Northeast Brazil (Isoptera). *Sociobiology* **59**, 183–188. (doi:10.13102/sociobiology.v59i1.675)
  62. Poderoso JCM, da Costa MKM, Correia-Oliveira ME, Dantas PC, Zanuncio JC, Ribeiro GT. 2013 Occurrence of *Tropidacris collaris* (Orthoptera: Acridoidea: Romaleidae) damaging *Casuarina glauca* (Casuarinaceae) plants in the municipality of Central Bahia, Brazil. *Fla. Entomol.* **96**, 268–269. (doi:10.1653/024.096.0143)
  63. Damascena JG, Leite GLD, Silva FWS, Soares MA, Guañabens REM, Sampaio RA, Zanuncio JC. 2017 Spatial distribution of phytophagous insects, natural enemies, and pollinators on *Leucaena leucocephala* (Fabaceae) trees in the Cerrado. *Fla. Entomol.* **100**, 558–565. (doi:10.1653/024.100.0311)
  64. Zanuncio-Junior JS, Fornazier MJ, dos Martins DS, Chamorro-Rengifo J, Queiróz RB, Lazzarini AL, Ferreira PSF. 2017 *Meroncidius intermedius* (Orthoptera: Tettigoniidae): a threat to Brazilian banana. *Fla. Entomol.* **100**, 669–671. (doi:10.1653/024.100.0329)
  65. Almeida CIM, Leite GLD, Rocha SL, Machado MML, Maldonado WCH. 2006 Phenology and arthropods of *Copaifera langsdorffii* Desf. in the cerrado (Brazilian savannah). *Rev. Bras. Pl. Med.* **8**, 64–70.
  66. Tiedge K, Lohaus G. 2017 Nectar sugars and amino acids in day- and night-flowering *Nicotiana* species are more strongly shaped by pollinators' preferences than organic acids and inorganic ions. *PLoS ONE* **12**, e0176865. (doi:10.1371/journal.pone.0176865)
  67. Stabler D, Power EF, Borland AM, Barnes JD, Wright GA. 2018 A method for analysing small samples of floral pollen for free and protein-bound amino acids. *Methods Ecol. Evol.* **9**, 430–438. (doi:10.1111/2041-210X.12867)
  68. Serra BDV, de Campos LAO. 2010 Entomophilic pollination of squash, *Cucurbita moschata* (Cucurbitaceae). *Neotrop. Entomol.* **39**, 153–159. (doi:10.1590/S1519-566X2010000200002)
  69. Silva FWS, Leite GLD, Guañabens REM, Sampaio RA, Gusmão CAG, Zanuncio JC. 2014 Spatial distribution of arthropods on *Acacia mangium* (Fabales: Fabaceae) trees as windbreaks in the Cerrado. *Fla. Entomol.* **97**, 631–638. (doi:10.1653/024.097.0240)
  70. Auslander M, Nevo E, Inbar M. 2003 The effects of slope orientation on plant growth, developmental instability and susceptibility to herbivores. *J. Arid Environ.* **55**, 405–416. (doi:10.1016/S0140-1963(02)00281-1)
  71. Landis D, Wratten SD, Gurr GM. 2000 Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annu. Rev. Entomol.* **45**, 175–201. (doi:10.1146/annurev.ento.45.1.175)
  72. Öberg S, Mayr S, Dauber J. 2008 Landscape effects on recolonisation patterns of spiders in arable fields. *Agr. Ecosyst. Environ.* **23**, 211–218. (doi:10.1016/j.agee.2007.06.005)
  73. Venturino E, Isaia M, Bona F, Chatterjee S, Badino G. 2008 Biological controls of intensive agroecosystems: wanderer spiders in the *Langa astigiana*. *Ecol. Complex* **5**, 157–164. (doi:10.1016/j.ecocom.2007.10.003)
  74. Miranda MMM, Picanço M, Leite GLD, Zanuncio JC, Clercq P. 1998 Sampling and non-action levels for predators and parasitoids of virus vectors and leaf miners of tomato in Brazil. *Med. Fac. Landbouww. Univ. Gent.* **63**, 519–523. (doi:10.1016/j.ecocom.2007.10.003)
  75. Picanço M, Ribeiro LJ, Leite GLD, Gusmão MR. 1998 Selectivity of insecticides to *Polybia ignobilis* (Hymenoptera: Vespidae) predator of *Ascia monusteoriseis* (Godart) (Lepidoptera: Pieridae). *An. Soc. Entomol. Bras.* **27**, 85–90. (doi:10.1590/S0301-80591998000100011)
  76. Leite GLD, Oliveira IR, Guedes RNC, Picanço M. 2001 Comportamento de predação de *Protonectarina sylveirae* (Saussure) (Hymenoptera: Vespidae) em mostarda. *Agro-Ciência* **17**, 93–96.
  77. Picanço MC, de Oliveira IR, Fernandes FL, Martinez HEP, Bacci L, da Silva EM. 2012 Ecology of Vespidae (Hymenoptera) predators in *Coffea arabica* plantations. *Sociobiology* **59**, 1269–1280. (doi:10.13102/sociobiology.v59i4.504)
  78. Larsen KJ, Purrington FF, Brewer SR, Taylor DH. 1996 Influence of sewage sludge and fertilizer on the ground beetle (Coleoptera: Carabidae) fauna of an old-field community. *Environ. Entomol.* **25**, 452–459. (doi:10.1093/ee/25.2.452)