

Acute and chronic toxicity and antimicrobial activity of the extract of *Stryphnodendron adstringens* (Mart.) Coville¹

Anna C. Almeida^{2*}, Viviane A. Andrade³, Francine S.A. Fonseca², Auricélio A. Macêdo⁴, Renato L. Santos⁴, Keila G.F. Colen², Ernane R. Martins² and Natália A. Marcelo²

ABSTRACT.- Almeida A.C., Andrade V.A., Fonseca F.S.A., Macêdo A.A., Santos R.L., Colen K.G.F., Martins E.R. & Marcelo N.A. 2017. **Acute and chronic toxicity and antimicrobial activity of the extract *Stryphnodendron adstringens* (Mart.) Coville.** *Pesquisa Veterinária Brasileira* 37(8):840-846. Universidade Federal de Minas Gerais, Av. Universitária 1000, Montes Claros, MG 39404-547, Brazil. E-mail: aca2006@ica.ufmg.br

This study evaluated the antimicrobial activity and acute or chronic toxicity of the extract of *Stryphnodendron adstringens*. The stem bark dry extract was obtained by static maceration with ethanol. Quantification of tannins was performed by the Folin-Denis method, which indicated a total tannin content of 32.7%. The antimicrobial activity of the dry extract of *S. adstringens* was evaluated by agar-based disk diffusion assay with *Escherichia coli* (ATCC 25922) and *Staphylococcus aureus* (ATCC 25923) in the concentration of 200, 400 and 600 µL/mL. The results indicated that 600 µL/mL inhibited microbial growth, i.e. had antimicrobial activity against these species. Acute and chronic toxic effects of *S. adstringens* was evaluated in Wistar rats treated with 200, 400, 600 and 800 mg/kg of extract, administered by gavage. Liver degeneration was observed in the group of rats receiving 800 mg/kg in chronic exposure, what may indicate some degree of toxicity at this concentration. However, no systemic toxicity was observed at lower doses. Considering the broad use of *S. adstringens* as a phytotherapeutic agent for various human and animal diseases and the liver toxicity observed at high concentrations, attention should be paid to the possible adverse effect of using the extract from this plant at high concentration.

INDEX TERMS: Toxicity, antimicrobial activity, extract, *Stryphnodendron adstringens*, tannins, liver toxicity, "barbatimão", hematological parameters, biochemical parameters.

RESUMO.- [Toxicidade aguda e crônica e atividade antimicrobiana do extrato de *Stryphnodendron adstringens* (Mart.) Coville.] Avaliou-se neste estudo a atividade antimicrobiana e a toxicidades aguda e crônica do extrato da entrecasca de *Stryphnodendron adstringens*. A partir do extrato seco, obtido através da maceração estática da casca do caule em etanol de cereais, foi realizada a quantificação de taninos totais (32,7%) pelo método de Folin-Denis. A

atividade antimicrobiana do extrato seco extraído de cascas do caule de *S. adstringens* foi avaliada pela técnica de disco-difusão para os micro-organismos *Escherichia coli* (ATCC 25922) e *Staphylococcus aureus* (ATCC 25923) nas concentrações de 200, 400 e 600 µL/mL. Os testes de concentração inibitória mínima revelaram que 600 µL/mL inibiu o crescimento dos dois micro-organismos testados; o mesmo resultado foi observado para atividade bactericida na concentração de 600 µL/mL sobre essas espécies. Efeitos tóxicos sistêmicos agudos e crônicos do extrato seco de *S. adstringens* administrados por gavagem foram estudados em ratos Wistar, utilizando as doses de 400, 600 e 800 mg/kg. Foi observada degeneração hepática no grupo de animais que receberam 800 mg/kg tanto no estudo da toxicidade aguda quanto crônica, que pode indicar algum grau de toxicidade de *S. adstringens* nessa concentração. Considerando o amplo uso de *S. adstringens* como fitoterápico

¹ Received on November 27, 2015.

Accepted for publication on September 4, 2016.

² Instituto de Ciências Agrárias, Universidade Federal de Minas Gerais (UFMG), Campus Montes Claros, Av. Universitária 1000, Montes Claros, MG 39404-547, Brazil. *Corresponding author: aca2006@ica.ufmg.br

³ Instituto Federal de Educação, Ciência e Tecnologia do Norte de Minas Gerais (IFNMG), Rua Dois 300, Village do Lago I, Montes Claros, MG 39404-058, Brazil.

⁴ Departamento de Clínica e Cirurgia, Escola de Veterinária, UFMG, Av. Antônio Carlos 6627, Pampulha, Belo Horizonte, MG 31270-901, Brazil.

para humanos e animais, atenção deve ser dispensada para ingestão em altas doses mediante os efeitos tóxicos observados neste estudo.

TERMOS DE INDEXAÇÃO: Toxicidade, atividade antimicrobiana, extrato, *Stryphnodendron adstringens*, taninos, toxicidade hepática, "barbatimão", parâmetros hematológicos, parâmetros bioquímicos.

INTRODUCTION

Tannins are water-soluble polyphenols that are present in many plants. There is a series of previous studies describing the biological action of tannins including their pharmacological, and toxic activities. The dosage and type of tannins determine these effects (Macáková et al. 2014).

Stryphnodendron adstringens, popularly known as barbatimão, is a medicinal plant abundant in central Brazil, which has been traditionally used as a popular herbal medicine (Melo et al. 2007), which is endemic to the savannah, and its bark has a high tannin content (25-37%). Macáková et al. (2014) and Okuda & Ito (2011) described a wide pharmacological spectrum of activity of polyphenols including hydrolysable tannins. However, Ferreira et al. (2009) and Aguiar-Filho (2013) reported intoxication of ruminants with plants that have tannins.

Local populations in the savannah regions of Brazil use infusion or tea of leaves or stem bark from *S. adstringens* as astringent, antimicrobial, homeostatic agent, antidiarrheal, antihypertensive or blood clotting agent. In addition, it is used for treating leucorrhoea, hemorrhoids, hemorrhages, and inflammation. Various studies have suggested the use of infusions and decoctions of this plant in Brazil for the treatment of animal diseases (Confessor et al. 2009, Viu & Viu 2011, Monteiro et al. 2011).

Several *in vitro* and *in vivo* studies on the activity of *Stryphnodendron* sp. tannins showed that the crude and purified extracts have anti-inflammatory, antiulcerogenic, antiprotozoal, anthelmintic, antiviral, and antifungal activities (Martins et al. 2003, Silva et al. 2009, Cansi & Demo 2011, Moura 2011). Its use in the clinical treatment of bacterial and parasitic diseases and for skin lesions in animals have also been reported (Confessor et al. 2009, Monteiro et al. 2011, Viu & Viu 2011).

Studies on the acute and chronic toxicity of *Stryphnodendron* sp. extracts showed variable results depending on the method and extract concentration studied. Some researchers have evaluated the acute toxicity of extract of stem bark from *S. adstringens* orally ingested (Lima et al. 1998, Audi et al. 1999, Rebecca et al. 2002). They demonstrated that LD_{50%} ranged from 400 to 2699mg/kg. An initial preclinical toxicological assay conducted by Almeida et al. (2010) demonstrated that *S. adstringens* leaf extract was most toxic among the tested (leaf hydroalcoholic extracts of *Lippia sidoides*, *Myracrodruon urundeuva* and *Caryocar brasiliense*), however, limited studies have been conducted to verify its toxic potential.

The National Policy for Medicinal Plants and Herbal Medicines (PNPMF), and the Brazilian National Program for Medicinal Plants and Phytotherapies published the Brazilian National List of Medicinal Plants of Interest to the SUS (RENISUS) in January 2009. This list contains medicinal

plants that exhibit potential for generating products of interest to Brazilian Unified Health System (SUS), including *S. adstringens* (Brasil 2006). This study aimed to evaluate the antimicrobial activity and acute and chronic toxicity of hydroalcoholic extracts of stem bark from *S. adstringens* from the northern region of Minas Gerais, Brazil.

MATERIALS AND METHODS

Obtaining and preparing the *Stryphnodendron adstringens* extract. *S. adstringens* stem bark was harvested at Bela Vista farm, in Botumirim, State of Minas Gerais, Brazil (16° 57.847' S, 43° 04.308' W, elevation: 890m). The plant was identified by Rubens Teixeira de Queiroz at the Federal University of Paraíba, Brazil. To obtain the extract, stem bark of *S. adstringens* was dried in a forced air circulation chamber under an average temperature of 50°C, followed by daily weighing until the material reached a constant weight. The dry *S. adstringens* bark was ground using a Willey knife mill in a continuous system. The ground material was subjected to extraction by static soaking in cereal ethanol for eight days (bark:ethanol ratio of 1:2). The mixture was then filtered, and the solvent was evaporated from the extract in an air-circulated oven at 45°C until completely dry. The dried *S. adstringens* extract obtained was stored in an amber vial away from moisture and light.

Phenols and tannins content. Initially, 0.75g of dry extract of *S. adstringens* was dissolved in 150ml of distilled water to obtain a solution 1 which was used to determine total phenolic by the Folin-Denis method using tannic acid (0.1, 0.5, 1.0, 2.5, and 3.75mg/mL) as the standard for the calibration curve (Pansera et al. 2003). The system was maintained at agitation for 30 minutes at 90°C. Next, the system was cooled, the content was transferred to a volumetric flask and the volume was completed to 250mL with distilled water. The flask was maintained at rest for 30 minutes, followed by filtration, and the first 50mL of the filtrate was discarded. The Folin-Denis method comprised addition of the solution 1 (5mL) to 2mL of Folin-Denis reagent (2.50g sodium tungstate, 0.5g phosphomolybdic acid, 1.25mL phosphoric acid, and distilled water for a final volume of 25mL) and 50mL of 20% (w/v) sodium carbonate solution. The solution 1 also was used to calculate the content of tannin. This calculation was carried out using a reference solution containing Folin-Denis reagent, pyrogallol and sodium carbonate. The absorbance was measured at wavelength of 715nm. The parameters was used in the expressions below. Samples were analyzed using a spectrophotometer 3 minutes after adding the last reagent. The analyses were performed in triplicate, and the tannin content was expressed in percentage (w/w) according to the equation proposed by Dôres & Casali (2007):

$$A^{1\%} = \frac{A_3 \times 10}{c} \quad TT = \frac{FD \times A_1}{(m-p) \times A^{1\%}}$$

Where: A^{1%} = specific absorbance of the reference solution, A₃ = measured absorbance of the reference substance, c = concentration in mg/mL, TT = total tannins in % (m/m), FD = 50 (dilution factor of the sample), A₁ = measured absorbance of the total tannins, m = water determination, p = drug mass (g) considering the water determination.

Microbial sensibility test. The microbial sensitivity test of *S. adstringens* extract was performed using the disk-diffusion technique. The minimum-inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) were defined according to the National Committee for Clinical Laboratory Standards (CLSI 2012). *Escherichia coli* (ATCC 25922) and *Staphylococcus aureus*

(ATCC 25923) were prepared and standardized to 0.5 on the MacFarland scale. Based on results of previous research (Pinho et al. 2012). Concentrations of 200, 400, and 600 $\mu\text{L}/\text{mL}$ were used for the disk-diffusion tests, and the MIC test. Further, 600 $\mu\text{L}/\text{mL}$ was used for the MBC test. Tests were performed in duplicate for each of the extract concentrations used.

Rats (*in vivo* toxicity). The acute toxicity classification method established by the OECD 420 (2001) was adopted to evaluate acute and OECD 452 (2008) with adaptations was employed to evaluate chronic toxicity. For this procedure, female Wistar rats, weighing between 180 and 200g, were used. The experimental procedures using rats were approved by the Animal Experimentation Ethics Committee of the UFMG, under protocol no. 034/2008.

Acute toxicity. After obtaining the microbiological test results, 200, 400, and 600mg/kg concentrations were tested. These doses were also defined by considering previously published data that reported concentrations up to 600mg/kg of *S. adstringens* bark extract administered orally that did not induce animal death (Rebecca et al. 2002). For evaluation of acute systemic toxicity, a single dose of each concentration prepared in sterile saline solution was administered by gavage. The rats were divided into four groups with five rats in each group: group 1 (control with sterile saline solution), group 2 (200mg/kg), group 3 (400mg/kg), and group 4 (600mg/kg). After treatment, all rats received food and water and a 12-h light/dark circadian cycle at $22\pm 2^\circ\text{C}$. The rats were weighed every other day. At the end of the experimental period (14 days), the surviving rats were sedated with a combination of ketamine (80mg) and xylazine (15mg/kg) administered intraperitoneally and killed. Blood samples (approximately 4mL) were collected by cardiac puncture to determine the hematological (erythrocyte, hemoglobin, hematocrit (PCV), MCV, MCH, MCHC, platelet, and lymphocyte count) and biochemical parameters (ALT, AST urea and creatinine). Macroscopic analyses were performed, and the liver, heart, and kidneys were weighed to determine their relative weights. Tissue samples were collected for histopathological analyses. These tissues were fixed by immersion in 10% buffered formalin for 24 hours, followed by dehydration in increasing ethanol concentrations, diaphonization in xylene, and paraffin embedding. Five- μm thick sections were cut and stained with hematoxylin and eosin (HE). Lesions were evaluated according to the intensity, and a score was given to each lesion, where 0 = absence of lesions, 1 = mild lesion, 2 = moderate lesion, and 3 = severe lesion. The analyses were performed at the Molecular Pathology Laboratory of the Veterinary School of the UFMG.

Chronic toxicity. Rats were kept under the same conditions described for the acute toxicity tests. The experiment was divided into two phases, which had either 15 or 30 days of duration. For each of these phases, two groups of five rats each were formed. One group received sterile saline solution (control) and the other received 800mg/kg of the *S. adstringens* extract, daily, once a day. At the end of each phase the rats were killed and fragments of the kidneys, liver, and heart were collected. Animal weight, hematological and biochemical parameters, macroscopic evaluation of the organs and histopathological analyses were performed according to the procedure described for acute toxicity evaluation.

Statistical analysis. Data were analyzed using analysis of variance (ANOVA) followed by Bonferroni post test or Dunn's multiple comparison test. When appropriate, the Student t-test with the Bonferroni adjustment was used for pairwise comparisons. A value of $p < 0.05$ was used to indicate a statistically significant difference. For histological analyses, the Kruskal-Wallis non-parametric test was performed followed by Dunn's multiple comparison test. All statistical tests were performed using the software Graph Pad Prism (San Diego, CA).

RESULTS AND DISCUSSION

Through the determination of phenolic compounds, it was possible to infer the total tannin content (32.7%) of the of *S. adstringens* stem bark. The result agrees with the Brazilian Pharmacopeia recommends minimum 8.0% of total tannins (Brasil 2010) and was similar to that reported in studies conducted in the same species (20-40%) (Castro et al. 2009, Corrêa et al. 2012). Tannins are the major compounds responsible for pharmacological activity of this particular tree and they exhibit various biological functions. Antimicrobial activity of tannins is well documented; however the mechanisms of this activity, as well as their bioavailability have not been satisfactorily clarified (Mačáková et al. 2014).




Fig.1. Sensitivity test for *Escherichia coli* and *Staphylococcus aureus* to the *Stryphnodendron adstringens* extract. Data represent mean and standard deviation. * Indicates a significant difference between the treatments and the control ($p < 0.05$).




Fig.2. Evaluation of the animal and organ weights. (A) Animal weight of the control and groups treated with different concentrations of *Stryphnodendron adstringens* extract; the values show a variation in the animal weight during 14 days of treatment. (B) Relative organ weight ($p < 0.05$).

Results of the sensitivity tests using the *S. adstringens* extract disks (600µL/mL) exhibited a significant inhibition zone against *S. aureus*; however, there was no significant difference against *Escherichia coli* (Fig.1). This same concentration inhibited *Staphylococcus aureus* and *E. coli* growth in the test tube. This was confirmed by the absence of microbial growth in Petri dishes, according to the recommendations of NCCLS (2003). Costa et al. (2011) used extract obtained by methodology different from the one adopted in this study to evaluate growth of *E. coli* and *S. aureus* isolated from bovine milk, and Pinho et al. (2012) evaluated the activity of *S. adstringens* extract obtained from leaves in these same microorganisms. These studies showed that concentrations ranging from 25 to 500mg/mL of dried crude *S. adstringens* extract did not inhibit *E. coli* growth (Costa et al. 2011, Pinho et al. 2012). Concentrations higher than or equal to 300mg/mL can inhibit *S. aureus* growth (Pinho et al. 2012). Although it is not possible to make a comparison of our results with the former ones, as they used methodology and sources of different extracts, our results corroborate with those findings and suggest that *S. adstringens* extract also inhibits microbial growth.

The rat weights (Fig.2A) and relative organ weights (Fig.2B) of the groups treated with the different concentrations of *S. adstringens* extract did not statistically differ compared to the control group, in contrast to the study of Rebecca et al. (2002) and Costa et al. (2013) that observed impairment in animal development at the dosages tested.

Hematological parameters were within the reference ranges, whereas no significant changes were observed in the complete blood count analyses and biochemical parameters, indicating that the rats did not develop any clinical abnormalities throughout the experiment.

Biochemical analysis of blood taken from rats treated with *S. adstringens* showed that there were changes only in concentration of aspartate aminotransferase (AST) in rats treated with 200mg/kg of *S. adstringens*, indicating a possible hepatic alteration in this group (Table 1). However, results of the histopathological analyses of the organs collected for the acute toxicity tests (based on histopathology scores) revealed that only the livers of the group treated with 600mg/kg of *S. adstringens* extract exhibited significant microscopic changes compared with the control group

Table 1. Biochemical parameters of Wistar rats treated with a single oral dose of the *Stryphnodendron adstringens* bark extract

Parameter	Control	200 mg/mL	400 mg/mL	600 mg/mL
ALT	55.8±3.8393	64±5.4924 ^a	67.4±3.5015	64±9.3005
AST	197.2±9.0078	148.4±15.917	204.2±23.0703	197.2±21.5021
Urea	37±3.6055	52.8±2.5573	50.6±4.1545	50.6±5.6973
Creatinine	0.374±0.02502	0.52±0.0483	0.424±0.03219	0.486±0.06592

Biochemical parameters obtained from the serum of rats, treated a single dose of an oral vehicle (Control, n=5) or a single oral dose of *S. adstringens* at doses of 200, 400, and 600mg/kg (n=5). Values represent mean ± standard deviation. Statistical comparisons of the experimental rats with the control group were performed using analysis of variance (ANOVA) and Bonferroni post-tests; n = represents the number of rats in each group. a = represents a significant difference when compared with the control (p<0.001).




Fig.3. Histopathological analysis of acute toxicity of the orally ingested *Stryphnodendron adstringens* extract in rats treated with 200, 400, and 600mg/mL for 14 days. *Indicates a significant difference between the treatments compared and the control (p<0.001).

(p<0.001) (Fig.3). This alteration was characterized by mild centrilobular degeneration, similar to lesions found in chronic toxicity tests as presented below. Pathological alterations were observed in high concentration, while physiological production of AST was altered only in lower concentration. This finding suggests that the *S. adstringens* extract may exhibit some liver toxicity. *S. adstringens* extract has been shown to be toxic in other studies in rats. Rebecca et al. (2002) demonstrated that repeated 800mg/kg and 1600mg/kg doses of the extract administered orally caused chronic toxicity in rats, where an LD_{50%} of 2669mg/kg was obtained. Rebecca et al. (2003) evaluated the toxicity of the same methanol extract on mitochondria obtained from liver cells and reported altered energy metabolism. Costa et al. (2013) evaluated acute and chronic toxicity of the same orally ingested fraction and reported no acute toxicity, thereby defining an LD_{50%} of 3015mg/kg. Furthermore, toxicity was not observed at the orally ingested doses of 10, 100, and 200mg/kg during the 90 days of chronic toxicity tests. Oliveira et al. (2013) evaluated the cytotoxicity of aqueous fractions of *S. adstringens* on rat macrophages at 50mg/mL doses and did not observe any decrease in cell viability.

The values obtained in this study were lower than those previously described (Rebecca et al. 2002, Costa et al. 2013), indicating that 400mg/kg of *S. adstringens* extract orally do not result in acute toxicity and can be safe and do not cause acute toxicity. Although this dose has not shown toxicity, was defined the dose of 800mg/kg for chronic toxicity experiments.

The results of the chronic toxicity experiments revealed no changes in body weight during 30 days of the experiment (Fig.4A). No significant changes in the organ weights were observed in treated rats when compared with the control group (Fig.4B). Regarding hematological and biochemical parameters, significant variation was not observed in any of the parameters evaluated (neither at 15 days nor at 30 days) relative to the control group, suggesting that there was no chronic toxicity in the rats. After 30 days of treatment only the MCV (mean corpuscular volume) and MCH (mean corpuscular concentration) had significantly decreased (Table 2). MCH indicates the quantity




Fig.4. Evaluation of the animal weight. The animal weight of the control and groups treated with 800mg/mL of *Stryphnodendron adstringens* extract. The values show a variation in the rat weight after 30 days of treatment. (A) Relative animal weight. (B) Relative organ weight.

of hemoglobin in a sample of blood and MCHC indicates the concentration of hemoglobin; reduction in hemoglobin is associated with anemia, so these results suggested a possible anemia of rats treated with 800mg/mL of *S. adstringens*, when compared with the control group. Also a decrease in urea concentration after 30 days was observed, which can be the result of a deregulation of renal and hepatic function when rats were treated with 800mg/mL of *S. adstringens* (Table 2). Similar data were obtained by Costa et al. (2013) who evaluated chronic toxicity with repeated doses for 90 days. They did not report any toxicity following the oral administration of the proanthocyanidin fraction to mice. However, Rebecca et al. (2002) found different




Fig.5. Histopathological analysis of the chronic toxicity of the *Stryphnodendron adstringens* extract. (A) Histopathological results of rats treated with 800mg/mL for 15 days, and (B) for 30 days. *Indicates a significant difference between the treatments and the control ($p < 0.001$).

results while using 800 or 1,600mg/kg for prolonged periods. The authors reported animal weight loss, increased blood glucose levels, and increased AST levels at the end of the 30-day treatment period.

Chronic toxicity was assessed by histopathological examination, and the results did not indicate any significant microscopic changes in heart and kidneys, neither at 15 days nor at 30 days after administering 800mg/kg of *S. adstringens* extract. However, in the liver, there was significant microscopic alteration in the group treated with 800mg/kg at 30 days compared to the control group ($p < 0.05$)

Table 2. Hematological and biochemical parameters of Wistar rats treated with oral doses of 800mg/kg of the *Stryphnodendron adstringens* bark extract for 30 days

	Hematological data			Biochemical data			
	Control	15 Days	30 Days	Control	15 Days	30 Days	
Erythrocytes(mm^3)	7.23 \pm 0.03	6.90667 \pm 1.59	7.9075 \pm 0.45	ALT	79.63 \pm 72.6	60.5 \pm 5.86	77 \pm 5.18
Hemoglobin (g/dL)	15.5 \pm 0.10	13.24 \pm 2.87	14.88 \pm 0.68	AST	165.2 \pm 66.51	155.5 \pm 66.39	247.8 \pm 97.07
Packed Cell Volum (%)	44.10 \pm 0.7	38.63 \pm 8.96	43.15 \pm 2.19	Urea	57.6 \pm 4.17	44.5 \pm 1.89	42.4 \pm 3.07 ^c
VCM (fL)	61.00 \pm 1.22	56.01 \pm 2.45	54.59 \pm 0.81 ^a	Creatinine	0.32 \pm 0.018	0.52 \pm 0.29	0.44 \pm 0.01
HCM (pg)	21.44 \pm 0.23	19.26 \pm 1.27	18.83 \pm 0.26 ^b				
CHCM (g/dL)	35.14 \pm 0.32	34.37 \pm 0.96	34.49 \pm 0.39				
Thrombocytes (mm^3)	613500 \pm 95500	277933 \pm 220468	449377 \pm 304037				
Leukocyte (mm^3)	7340 \pm 240	10417 \pm 3579	5654 \pm 6532				

Hematological and biochemical parameters measured in the serum of Wistar rats treated chronically using oral vehicle (Control, n=4) or with daily doses of 120mg/kg of *S. adstringens* (n=5) for 30 days. The values represent the mean \pm standard deviation. Statistical comparisons of the experimental rats with the control group were performed using analysis of variance (ANOVA) and Bonferroni post-test; n = represents the number of rats in each group. The letters a ($p < 0.04017$), b ($p < 0.005$) and c ($p < 0,0114$) represent a significant difference when compared with the control.




Fig.6. Representative micrographs of the liver from a control rat (with no histological changes), or treated with 800mg/kg of *Stryphnodendron adstringens* extract for 15 or 30 days, with minimal or mild to moderate centrilobular degeneration (hydropic hepatocellular degeneration) respectively.

(Fig.5A,B) as demonstrated by histopathology scores. This analysis revealed that the livers of the group treated with of *S. adstringens* extract exhibited minimal or mild to moderate centrilobular degeneration (hydropic hepatocellular degeneration) (Fig.6).

Rebecca et al. (2002, 2003) observed altered plasma glucose and AST levels combined with a decline in weight of the rats. The effect of the extracts on energy metabolism in the liver was investigated and the results showed that the toxic action may be correlated with interference in the oxidative phosphorylation or electron transport chain or with inhibition ATP-synthase. The authors concluded that such injuries were associated with toxic activity of the tannin present in the extracts when administered at doses above 600mg/kg. Costa et al. (2013) found lesions indicative of toxicity, while analyzing the organs from rats subjected to acute and chronic toxicity evaluation of the proanthocyanidin fraction, where only 4000 and 5000mg/kg doses were used, thereby indicating that the fraction exhibited lower toxicity at the tested concentrations than those used by Rebecca et al. (2002, 2003) and those employed in this study.

Liver lesions have been described in ruminants after intake of mashed beans from *Stryphnodendron fissuratum* (Ferreira et al. 2009, Aguiar-Filho 2013). Fonseca & Linbradi (2008) reported that possible toxicity of tannins in dye prepared with bark of *Stryphnodendron barbatiman* occurred in mammalian cells. Regarding the cytotoxicity in other mammalian cells, Oliveira et al. (2013) found mild toxicity of the *S. barbatiman* Mart. plant extract at 50mg/mL concentration on rat macrophages and significantly reduced the production of TNF- α when compared to the control group.

CONCLUSIONS

According to the sensitivity tests, the *Stryphnodendron adstringens* extract obtained from northern Minas Gerais exhibited antimicrobial activity at 600 μ L/mL.

Liver degeneration, although minimal, was observed 30 days after ingestion of the bark extract of *S. adstringens* in the groups of rats that received 800mg/kg, what may indicate some degree of toxicity.

No systemic toxicity was observed at lower doses.

Considering the broad use of *S. adstringens* as a phytotherapeutic agent for various human and animal diseases and liver toxicity observed at high concentrations, atten-

tion should be paid to the possible adverse effects of using the extract from this tree at high concentrations.

Acknowledgements. To FAPEMIG, CNPq, CAPES, the Dean of Research of the UFMG (PRPq/UFMG), and to the Laboratory of Instrumental Chemistry-ICA/UFMG.

REFERENCES

- Aguiar-Filho C.R., Albuquerque R.F., Rocha B.P., Colodel E.M., Lemos R.A.A., Riet-Correa F., Evêncio-Neto J. & Mendonça F.S. 2013. Avaliação da toxicidade das favas de *Stryphnodendron fissuratum* (Mimosoideae) em vacas gestantes. *Pesq. Vet. Bras.* 33(5):607-612.
- Almeida A.C., Sobrinho E.M., Pinho L., Souza P.N.S., Martins E.R., Duarte E.R., Santos H.O., Brandi I.V., Cangussu A.S. & Costa J.P.R. 2010. Toxicidade aguda de extratos hidroalcoólicos de folhas de alecrim-pimenta, *Stryphnodendron adstringens*, aroeira e farelo da casca de pequi administrados intraperitonealmente. *Ciência Rural* 40(1):200-203.
- Audi E.A., Toledo D.P., Perez P.G., Kimura E., Pereira W.K., Mello J.C., Nakamura C., Alves-do-Prado W., Cuman R.K. & Bersani-Amado C.A. 1999. Gastric antitumorogenic effects of *Stryphnodendron adstringens* in rats. *Phytother. Res.* 13(3):264-266.
- Brasil 2010. Farmacopeia Brasileira. Vol.2. Agência Nacional de Vigilância Sanitária, Brasília. 904p.
- Brasil 2006. Ministério da Saúde. Agência Nacional de Vigilância Sanitária. Decreto no.5813, de 22 de junho de 2006. Aprova a Política Nacional de Plantas Medicinais e Fitoterápicos e de Outras Provisões. Ministério da Saúde < Brasília, DF. Available in <http://bvsms.saude.gov.br/bvs/publicacoes/politica_nacional_fitoterapicos.pdf> Accessed on December 10, 2013.
- Cansi E.R. & Demo C. 2011. Ocorrência de miíases em animais de companhia no Distrito Federal, Brasil. *Acta Scient. Vet.* 39(3):1-5.
- Castro A.H.F., Paiva R., Alvarenga A.A. & Vitor S.M.M. 2009. Calogênese e teores de fenóis e taninos totais em barbatimão [*Stryphnodendron adstringens* (Mart.) Coville]. *Ciênc. Agrotec.* 33(2):385-390.
- CLSI 2012. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically. Approved Standard-Ninth Edition. CLSI document M07-A9. Clinical and Laboratory Standards Institute, Wayne, PA. Available in <<http://antimicrobianos.com.ar/ATB/wp-content/uploads/2012/11/03-CLSI-M07-A9-2012.pdf>> Accessed on January 22, 2016.
- Confessor M.V.A., Mendonça L.V., Mourão J.S. & Alves R.R.N. 2009. Animals to heal animals: ethnoveterinary practices in semiarid region, northeastern Brazil. *J. Ethnobiol. Ethnomed.* 5(37). Available in <<http://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-5-37>> Accessed on December 10, 2013.
- Corrêa V.S., Cerdeira A.L., Fachin A.L., Bertoni B.W., Pereira P.S., França P.S., Momm H.G., Moraes R.M. & Pereira A.M.S. 2012. Geographical variation and quality assessment of *Stryphnodendron adstringens* (Mart.) Coville within Brazil. *Genet. Resour. Crop Evolution* 59(7):1349-1356.
- Costa J.P.R., Almeida A.C., Martins E. R., Rodrigues M.N., Santos C.A. & Menezes I.R. 2011. Atividade antimicrobiana do óleo essencial de alecrim-

- pimenta e do extrato bruto seco do barbatimão diante de bactérias isoladas do leite. *Biotemas* 24(4):1-6.
- Costa M.A., Mello J.C., Kaneshima E.N., Ueda-Nakamura T., Dias Filho B.P., Audi E.A. & Nakamura C.V. 2013. Acute and chronic toxicity of an aqueous fraction of the stem bark of *Stryphnodendron adstringens* (barbatimão) in rodents. *Evid. Based Complement. Alternat. Med.* Available in: <<http://dx.doi.org/10.1155/2013/841580>> Accessed on April 14, 2014.
- Dôres R.G.R. & Casali V.W.D. 2007. Plantas Medicinais e Aromáticas: controle de qualidade. Universidade Federal de Viçosa, Viçosa, MG, p.106-107.
- Ferreira E.V., Boabaid F.M., Arruda L.P., Lemos R.A.A., Souza M.A., Nakazato L. & Colodel E.M. 2009. Intoxicação por *Stryphnodendron fissuratum* (Mimosoideae) em bovinos. *Pesq. Vet. Bras.* 29(11):951-957.
- Fonseca P. & Librandi A.P.L. 2008. Avaliação das características físico-químicas e fitoquímicas de diferentes tinturas de barbatimão (*Stryphnodendron adstringens barbatiman*). *Revta Bras. Ciênc. Farm.* 44(2):271-277.
- Lima J.C.S., Martins D.T.O. & Souza Júnior P.T. 1998. Experimental evaluation of stem bark of *Stryphnodendron adstringens* (Mart.) Coville for anti-inflammatory activity. *Phytother. Res.* 12:212-220.
- Macáková K., Kolečkář V., Cahlíková L., Chlebek J., Hošťálková A., Kuča K., Jun D. & Opletal L. 2014. Tannins and their Influence on health, p.159-208. In: Rahman A. & Choudhary M.I. (Eds), *Recent Advances in Medicinal Chemistry: tannins_and_their_influence_on_health*. Available in <<https://www.researchgate.net/publication/282796040>> Accessed on: January 22, 2016.
- Martins P.S., Alves A.L.G., Hussni J.L., Sequeira J.L., Nicoletti J.L.M. & Thomassian A. 2003. Comparação entre fitoterápicos de uso tópico na cicatrização de pele em equinos. *Arch. Vet. Sci.* 8(2):1-7.
- Melo J.O., Endo T.H., Bersani-Amado L.E., Estivalet S.A., Silmara B., Mello J.C.P. & Bersani-Amado C.A. 2007. Effect of *Stryphnodendron adstringens* (barbatimão) bark on animal models of nociception. *Braz. J. Med. Biol. Res.* 43(3):465-469.
- Monteiro M.V.B., Bevilaqua C.M.L., Palha M.D.C., Braga R.R., Schwanke K., Rodrigues S.T. & Lameira O.A. 2011. Ethnoveterinary knowledge of the inhabitants of Marajó Island, eastern Amazonia, Brazil. *Acta Amaz.* 41(2):233-242.
- Moura M.I. 2011. Enfermidades cutâneas digitais bovina: aspectos genéticos e terapêuticos. Tese de Doutorado, Escola de Veterinária e Zootecnia, Universidade Federal de Goiás, Goiânia, Go. 111p.
- OECD 2001. Guidelines for the Testing of Chemicals, n. 420: acute oral toxicity, fixed dose procedure. Available in <<http://www.oecd-ilibrary.org/docserver/download/9742001e.pdf?expires=1455306679&id=id&acname=guest&checksum=635B12BD5729564AEC4136E85BC32D12>> Accessed on January 22, 2016.
- OECD 2008. Guidelines for the Testing of Chemicals, n. 452: chronic toxicity studies. Available in <<http://www.oecd-ilibrary.org/docserver/download/9745201e.pdf?expires=1455306478&id=id&acname=guest&checksum=417154C7DA531CDAB6D1BD24D2DDA885>> Accessed on January 22, 2016.
- Okuda T. & Ito H. 2011. Tannins of constant structure in medicinal and food plants-hydrolyzable tannins and polyphenols related to tannins. *Molecules* 16:2191-2217.
- Oliveira J.R., Castro V.C., Vilela P.G.F., Camargo S.E.A., Carvalho A.T., Jorge A.O.C. & Oliveira L.D. 2013. Cytotoxicity of Brazilian plant extracts against oral microorganisms of interest to dentistry. *BMC Complement. Altern. Med.* Available in <<http://www.biomedcentral.com/1472-6882/13/208>> Accessed on April 14, 2014.
- Pansera M.R., Santos A.C.A., Paese K., Wasum R., Rossato M., Rota L.D., Pauletti G.F. & Serafini L.A. 2003. Análise de taninos totais em plantas aromáticas e medicinais cultivadas no Nordeste do Rio Grande do Sul. *Revista Bras. Farmacogn.* 13(1):17-22.
- Pinho L., Souza P.N.S., Sobrinho Macedo E., Almeida A.C. & Martins E.R. 2012. Atividade antimicrobiana de extratos hidroalcoólicos das folhas de alecrim-pimenta, aroeira, barbatimão, erva baleeira e do farelo da casca de pequi. *Ciênc. Rural* 42(2):326-331.
- Rebecca M.A., Ishii-Iwamoto E.L., Kelmer-Bratch A.M., Caparroz-Assef S.M., Cuman R.K., Pagadigorria C.L., Mello J.C., Bracht A. & Bersani-Amado C.A. 2003. Effect of *Stryphnodendron adstringens* (barbatimão) on energy metabolism in the rat liver. *Toxicol. Lett.* 143:55-63.
- Rebecca M.A., Ishii-Iwamoto E.L., Grespan R., Cuman R.K., Caparroz-Assef S.M., Mello J.C. & Bersani-Amado C.A. 2002. Toxicological studies on *Stryphnodendron adstringens*. *J. Ethnopharmacol.* 83(1/2):101-104.
- Silva L.A.F., Moura M.I., Persiano C.B., Helou J.B., Lima C.R.O., Freitas S.L.R., Silva J.A., Goulart D.S. & Castro L.T.S. 2009. Extrato da casca do barbatimão (*Stryphnodendron barbatiman* Martius) associado ao tratamento cirúrgico e toalete dos cascos na recuperação de bovinos da raça nelore com dermatite digital. *Ciênc. Anim. Bras. (Supl.1)*:373-378.
- Viu A.F.M. & Viu M.A.O. 2011. Cerrado e etnoveterinária: o que se sabe em Jataí, GO. *Revta Bras. Agroecol.* 6(3):49-61.