We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,100
Open access books available

167,000

185M

Downloads

en access books available International authors and e

Our authors are among the

154
Countries delivered to

TOP 1%

12.2%

most cited scientists

Contributors from top 500 universitie



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Chapter

Introductory Chapter: CO₂ Sequestration

Leidivan A. Frazão, Junio C. Silva and Adriana M. Silva-Olaya

1. Introduction

The Special Report from the Intergovernmental Panel on Climate Change (IPCC) [1] revealed that recent trends in greenhouse gas (GHG) emissions and the level of international ambition indicated by nationally determined contributions, within the Paris Agreement, deviate from a track consistent with limiting warming to well below 2°C. This will require a drastic reduction in greenhouse gas emissions by 2030 and thereafter removal of carbon from the atmosphere in large quantities. The IPPC reports found that many climate models can only meet the two-degree Celsius goal when carbon removal strategies are included among the potential policy options.

There are several strategies to promote carbon dioxide (CO_2) sequestration by agriculture and industry. So, it is necessary to evaluate the methodologies that have been used and to understand the gaps to achieve more sustainable production systems.

In agriculture, the management of agricultural systems that promote soil carbon sink depends on depth, clay content and mineralogy, plant available water holding capacity, nutrient reserves, landscape position, and the antecedent SOC stock [2]. As the soil carbon fluxes vary according to environmental and anthropogenic driving factors [3], soil carbon sequestration can be a short-term solution of reducing CO₂ concentration in the atmosphere.

In addition to agronomic practices, several effective methods of carbon capture and storage (CCS) have been proposed to reduce the amount of emitted CO_2 in the atmosphere. Adsorption processes can be performed using activated carbon [4] where the adsorptive process can use adsorbents derived from low-cost agro-wastes. Another way to reduce CO_2 emission into the atmosphere is by capturing CO_2 from the flue gases and storing that in deep geological formations [5]. The CCS provides financial offsets in terms of CO_2 sequestration cost.

Therefore, this book provides a comprehensive overview of the current state of the art about the strategies that contribute to reducing GHG emissions and promote CO_2 sequestration by agricultural techniques and carbon capture and storage.

2. Opportunities and challenges for CO₂ sequestration

Several studies have indicated the storage in biomass, soils, adsorption processes, and geological formations as viable techniques for CO_2 sequestration. All these technologies have the potential to mitigate global warming and climate change [2–6].

Improving agricultural land management techniques is an efficient way to increase carbon uptake and storage. Strategies to ensure soil carbon sequestration can be obtained through the adoption of different agronomic management practices [2]. Land use with grassland species can also maintain and increase soil organic carbon storage over time [7]. Other studies have reported that land use with perennial crops can also be adopted to promote CO₂ sequestration in biomass, and soil is the main component storing the highest amount of carbon in these agroecosystems [8, 9].

Carbon removal can also be achieved through the technology of adsorption on activated carbon from low-cost raw material. Agricultural and forestry residues or biomass residue wastes could be used as suitable raw materials for the production of activated carbon [10]. Furthermore, CCS by geological sequestration is another technological form for carbon removal and can be applied to different industries [5].

3. Perspectives

As the population is growing around the world and indirectly contributes to global warming, several efforts have been made to mitigate GHG emissions. So, the adoption of CO₂ sequestration technologies in the agricultural and industrial sectors has become essential to reduce the impacts of global warming and climate change.

Author details

Leidivan A. Frazão^{1*}, Junio C. Silva¹ and Adriana M. Silva-Olaya²

- 1 Universidade Federal de Minas Gerais, Montes Claros, MG, Brazil
- 2 Universidad de la Amazonia, Florencia, Colombia

*Address all correspondence to: lafrazao@ufmg.br

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. CC) BY

References

- [1] International Panel on Climate Change (IPCC). In: Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J, Shukla PR, et al., editors. Global Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty. 2018. Available from: https://www.ipcc.ch/site/assets/ uploads/sites/2/2019/06/SR15_Full_ Report_High_Res.pdf
- [2] Lal R. Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in agroecosystems. Global Change Biology. 2018;24:3285-3301
- [3] Stockmann U, Adams M, Crawford JW, Field DJ, Henakaarchchia N, Jenkins M, et al. The knowns, known unknowns and unknowns of sequestration of soil organic carbon. Agriculture, Ecosystems and Environment. 2013;**164**:80-90
- [4] Mohammad SS, Wan Mohd Ashri WD, Amirhossein H, Ahmad S. A review on surface modification of activated carbon for carbon dioxide adsorption. Journal of Analytical and Applied Pyrolysis. 2010;89:143-151
- [5] Mosleh MH, Sedighi M, Babaei M, Turner M. Geological sequestration of carbon dioxide. In: Letcher TM, editor. Managing Global Warming: An Interface of Technology and Human Issues. London, United Kingdon: Academic Press (Elsevier); 2019. pp. 487-500
- [6] Vaughan NE, Gough N, Mander S, Littleton EW, Welfle A, Gernaat DEHJ, et al. Evaluating the use of biomass energy with carbon

- capture and storage in low emission scenarios. Environmental Research Letters. 2018;**13**:044014. DOI: 10.1088/1748-9326/aaaa02
- [7] Hungate BA, Barbier EB, Ando AW, Marks SP, Reich PB, van Gestel N, et al. The economic value of grassland species for carbon storage. Science Advances. 2017;3:e1601880. DOI: 10.1126/sciadv.1601880
- [8] Leblanc HA, Russo RO. Carbon sequestration in an oil palm crop system (Elaeis guineensis) in the Caribbean lowlands of Costa Rica. Proceedings of the Florida State Horticultural Society. 2008;**121**:52-54
- [9] Mohammed AM, Robinson JS, Midmore D, Verhoef A. Carbon storage in Ghanaian cocoa ecosystems. Carbon Balance and Management. 2016;**11**(6). DOI: 10.1186/s13021-016-0045-x
- [10] Kaghazchi T, Soleimani M. Effect of raw materials on properties of activated carbons. Chemical Engineering and Technology. 2006;29:1247-1251. DOI: 10.1002/ceat.200500298

