

References

1. Zhao G, Gu X, Lu X, Pan J, Duan Z, Zhao K, et al. Novel reassortant highly pathogenic H5N2 avian influenza viruses in poultry in China. *PLoS One*. 2012;7:e46183. <http://dx.doi.org/10.1371/journal.pone.0046183>
2. Lee YJ, Kang HM, Lee EK, Song BM, Jeong J, Kwon YK, et al. Novel reassortant influenza A(H5N8) viruses, South Korea, 2014. *Emerg Infect Dis*. 2014;20:1087–9. <http://dx.doi.org/10.3201/eid2006.140233>
3. Zhou LC, Liu J, Pei EL, Xue WJ, Lyu JM, Cai YT, et al. Novel avian influenza A(H5N8) viruses in migratory birds, China, 2013–2014. *Emerg Infect Dis*. 2016;22:1121–3. <http://dx.doi.org/10.3201/eid2206.151754>
4. Wu H, Peng X, Xu L, Jin C, Cheng L, Lu X, et al. Novel reassortant influenza A(H5N8) viruses in domestic ducks, eastern China. *Emerg Infect Dis*. 2014;20:1315–8. <http://dx.doi.org/10.3201/eid2008.140339>
5. Jeong J, Kang HM, Lee EK, Song BM, Kwon YK, Kim HR, et al. Highly pathogenic avian influenza virus (H5N8) in domestic poultry and its relationship with migratory birds in South Korea during 2014. *Vet Microbiol*. 2014;173:249–57. <http://dx.doi.org/10.1016/j.vetmic.2014.08.002>
6. Marchenko VY, Susloparov IM, Kolosova NP, Goncharova NI, Shipovalov AV, Durymanov AG, et al. Influenza A(H5N8) virus isolation in Russia, 2014. *Arch Virol*. 2015;160:2857–60. <http://dx.doi.org/10.1007/s00705-015-2570-4>
7. Lee DH, Torchetti MK, Winker K, Ip HS, Song CS, Swayne DE. Intercontinental spread of Asian-origin H5N8 to North America through Beringia by migratory birds. *J Virol*. 2015;89:6521–4. <http://dx.doi.org/10.1128/JVI.00728-15>
8. Olsen B, Munster VJ, Wallensten A, Waldenström J, Osterhaus AD, Fouchier RA. Global patterns of influenza a virus in wild birds. *Science*. 2006;312:384–8. <http://dx.doi.org/10.1126/science.1122438>
9. Florin Beltran M. Uvs Nuur Lake. In: *Biomes and ecosystems*. Amenia (NY): Salem Press; 2013. p. 1260–1
10. Sharshov K, Silko N, Sousloparov I, Zaykovskaya A, Shestopalov A, Drozdov I. Avian influenza (H5N1) outbreak among wild birds, Russia, 2009. *Emerg Infect Dis*. 2010;16:349–51. <http://dx.doi.org/10.3201/eid1602.090974>

Address for correspondence: Alexander Shestopalov, 630117, Timakova St., Bldg. 2, Room 9-155, Novosibirsk, Russia; email: shestopalov2@ngs.ru

Detection of Vaccinia Virus in Urban Domestic Cats, Brazil

Galileu Barbosa Costa, Júlia Bahia Miranda, Gregório Guilherme Almeida, Jaqueline Silva de Oliveira, Mariana Siqueira Pinheiro, Stefanne Aparecida Gonçalves, Jenner Karlisson Pimenta dos Reis, Ricardo Gonçalves, Paulo César Peregrino Ferreira, Cláudio Antônio Bonjardim, Jônatas Santos Abrahão, Erna Geessien Kroon, Giliane de Souza Trindade

Author affiliation: Universidade Federal de Minas Gerais, Belo Horizonte, Brazil

DOI: <http://dx.doi.org/10.3201/eid2302.161341>

We investigated possible vaccinia virus (VACV) in urban house cats in Brazil. Serum samples from 6 cats were positive for VACV by PCR, indicating likely VACV circulation among house cats in urban areas of Brazil. This finding highlights the importance of epidemiologic surveillance to avoid outbreaks among urban human populations.

Vaccinia virus (VACV) outbreaks, first reported in Brazil in 1999, affect dairy cattle and humans in rural areas (1). Although studies have shown evidence of VACV circulation among several mammal species (1–3), no consensus exists regarding the role of these animals in the VACV transmission chain or which animal is the natural reservoir. In fact, domestic or wild mammals could be asymptomatic hosts and also contribute to VACV transmission (3).

In contrast to VACV, cowpox virus (CPXV) circulates in urban environments in Europe but also in surrounding wild and rural areas (4). CPXV is transmitted to humans mainly by cats, which play a link between the natural reservoirs and humans in the urban environment (4,5). In cats, the clinical course of CPXV infection varies from no symptoms to widespread skin necrotic lesions and can ultimately lead to death (6). Some studies have shown serologic evidence of orthopoxvirus infection in cats from Europe and have addressed the role of these animals in orthopoxvirus transmission to humans (7,8).

Because VACV and CPXV share some epidemiologic features and cats have a prominent role in the urban CPXV transmission chain, we decided to investigate whether urban domestic cats have evidence of exposure to VACV in Brazil. This study was approved by the Animal Experiments Committee of the Universidade Federal de Minas Gerais (registration protocol 315/2014).

We performed a retrospective study of serum samples from 277 house cats, collected during September 2012–December 2014 in 5 states in Brazil (online Technical Appendix Figure 1, <https://wwwnc.cdc.gov/EID/article/23/2/16-1341-Techapp1.pdf>). The states in this study were those whose veterinary clinics agreed to submit samples. We screened serum samples for neutralizing antibodies by using a $\geq 70\%$ plaque-reduction neutralization test (9). To detect VACV DNA in serum samples, we performed real-time PCR targeting the C11R and A56R genes (9). We directly sequenced A56R fragments in both orientations and in triplicate by using the Mega-BACE sequencer (GE Healthcare, Buckinghamshire, UK). We used ClustalW (<http://www.genome.jp/tools/clustalw>) and MEGA7 soft-

ware (<http://www.megasoftware.net>) to align nucleotide sequences and construct a phylogenetic tree (neighbor-joining method with 1,000 bootstraps).

The cats' ages ranged from 3 months to 15 years; 150 (54.2%) of the cats were female. Thirteen cats (4.7%) had outdoor access, and 37 (13.4%) were admitted to the veterinary clinic for ≥ 1 night. Some cats had clinical illness inconsistent with orthopoxvirus infection, which can overlap with other common dermatologic diseases affecting cats (online Technical Appendix Table). Most (8/53 [15.1%]) seropositive cats were from the Pampulha region of the city of Belo Horizonte (Minas Gerais State) (online Technical Appendix Figure 1), followed by the eastern region of

the city. We detected neutralizing antibodies in 16 animals (5.8%), with titers ranging from 100 to 1,600 neutralizing units/mL; of these, 13 (4.7%) were positive for C11R gene and 6 for A56R gene (online Technical Appendix Table). Alignment of the A56R fragments showed high similarity to the homologous gene of VACV isolates from Brazil (online Technical Appendix Figure 1). For the phylogenetic tree, we grouped sequences with VACV group 1 and 2 isolates (Figure).

We describe evidence of VACV circulation in cats in an urban environment in Brazil. Many studies have attempted to elucidate VACV outbreaks and risk factors in rural and wild areas (1–3). Our findings reveal a seropositivity rate of




Figure. Phylogenetic tree constructed based on nucleotide sequences of orthopoxvirus A56R (hemagglutinin) genes detected in serum samples of 6 house cats house cats with neutralizing antibodies for vaccinia virus, Belo Horizonte, Brazil, September 2012–December 2014. The tree was constructed with A56R gene sequences by using the neighbor-joining method with 1,000 bootstrap replicates and the Tamura 3-parameter model in MEGA7 (<http://www.megasoftware.net>). Asterisks indicate group 1 vaccinia virus isolates (nonvirulent strains) detected in 2 cats. Black circles indicate group 2 vaccinia virus isolates (virulent strains) detected in 4 cats. Numbers along branches are bootstrap values. GenBank accession numbers are shown for reference isolates. Scale bar indicates nucleotide substitutions per site.

5.8%, which is lower than the rate observed in a previous study from Norway (8) and higher than the rate observed in a study of cats in Austria (7). Notably, the Pampulha region, where most seropositive animals were detected, corresponded to areas of relatively low elevation that feature houses with green areas, cottage houses, and ecologic parks, with forested areas making up the remaining portion of the land (online Technical Appendix Figure 1).

Recent data from our research group revealed that capybaras (*Hydrochoerus hydrochaeris*) from the Pampulha region tested positive for VACV (10). These data, corroborated by molecular detection of VACV groups 1 and 2 in house cats from Belo Horizonte, further indicate the presence of VACV in an urban environment (online Technical Appendix Figure 2). In this study, PCR-positive cats showed no clinical signs that would indicate orthopoxvirus infection at the time of sample collection (online Technical Appendix Table), unlike what was observed among cats infected with CPXV in Europe (4,5). Furthermore, cats 4, 10, and 15 (online Technical Appendix Table), in which we detected ongoing VACV DNA, had no clinical signs. Although we detected group 2 VACV (virulent strains) in 4 samples, our findings corroborate the results of Bennett et al. (6), which showed that cats infected with VACV had asymptomatic infection.

Limitations of our study include selection bias of animals; it was not possible to use a convenience sample from the 5 Brazilian states. We were also unable to obtain detailed clinical information of all animals and unable to collect additional clinical samples to better understand the clinical course of VACV infection in cats. In Brazil, no records of VACV-like detection in urban populations are available, despite the fact that VACV was recently found in urban areas (10). In fact, potential sources of infection for cat populations (e.g., small rodents) should be considered. Cats could possibly seroconvert without the onset of classical illness. Hence, VACV could be circulating in cats from urban environments. The potential role of cats in infecting humans should be investigated further to determine whether VACV can emerge in urban human populations and pose a threat to public health.

Acknowledgments

We thank Associação Bichos Gerais, who provided some cat serum samples from Belo Horizonte. We also thank colleagues from the Laboratório de Vírus (ICB-UFMG) for their excellent technical support.

Financial support was provided by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq),

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), and Pró-Reitoria de Pesquisa/UFMG (PRPq). C.A. Bonjardim, J.S. Abrahão, E.G. Kroon, J.K.P. Reis, and G.S. Trindade are researchers from CNPq.

Mr. Costa is a PhD candidate in biomedicine at the Laboratório de Vírus, Microbiology Department, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais. His research is focused on the diagnosis, epidemiology, control, and prevention of orthopoxviruses and other emerging viruses.

References

1. Kroon EG, Mota BE, Abrahão JS, da Fonseca FG, de Souza Trindade G. Zoonotic Brazilian vaccinia virus: from field to therapy. *Antiviral Res.* 2011;92:150–63. <http://dx.doi.org/10.1016/j.antiviral.2011.08.018>
2. Peres MG, Bacchiega TS, Appolinário CM, Vicente AF, Allendorf SD, Antunes JM, et al. Serological study of vaccinia virus reservoirs in areas with and without official reports of outbreaks in cattle and humans in São Paulo, Brazil. *Arch Virol.* 2013;158:2433–41. <http://dx.doi.org/10.1007/s00705-013-1740-5>
3. Peres MG, Barros CB, Appolinário CM, Antunes JM, Mioni MS, Bacchiega TS, et al. Dogs and opossums positive for vaccinia virus during outbreak affecting cattle and humans, São Paulo State, Brazil. *Emerg Infect Dis.* 2016;22:271–3. <http://dx.doi.org/10.3201/eid2202.140747>
4. Essbauer S, Pfeffer M, Meyer H. Zoonotic poxviruses. *Vet Microbiol.* 2010;140:229–36. <http://dx.doi.org/10.1016/j.vetmic.2009.08.026>
5. Carletti F, Bordini L, Castilletti C, Di Caro A, Falasca L, Gioia C, et al. Cat-to-human orthopoxvirus transmission, northeastern Italy. *Emerg Infect Dis.* 2009;15:499–500. <http://dx.doi.org/10.3201/eid1503.080813>
6. Bennett M, Gaskell RM, Gaskell CJ, Baxby D, Kelly DF. Studies on poxvirus infection in cats. *Arch Virol.* 1989;104:19–33. <http://dx.doi.org/10.1007/BF01313805>
7. Nowotny N. Serologic studies of domestic cats for potential human pathogenic virus infections from wild rodents [in German]. *Zentralbl Hyg Umweltmed.* 1996;198:452–61.
8. Tryland M, Sandvik T, Holtet L, Nilsen H, Olsvik O, Traavik T. Antibodies to orthopoxvirus in domestic cats in Norway. *Vet Rec.* 1998;143:105–9. <http://dx.doi.org/10.1136/vr.143.4.105>
9. Geessien Kroon E, Santos Abrahão J, de Souza Trindade GS, Pereira Oliveira G, Moreira Franco-Luiz AP, Barbosa Costa G, et al. Natural vaccinia virus infection: diagnosis, isolation, and characterization. *Curr Protoc Microbiol.* 2016;42:14A.5.1–14A.5.43. <http://dx.doi.org/10.1002/cpmc.13>
10. Dutra LA, de Freitas Almeida GM, Oliveira GP, Abrahão JS, Kroon EG, Trindade GS. Molecular evidence of Orthopoxvirus DNA in capybara (*Hydrochoerus hydrochaeris*) stool samples. *Arch Virol.* 2016. <http://dx.doi.org/10.1007/s00705-016-3121-3>

Address for correspondence: Giliane de Souza Trindade, 6627 Avenida Presidente Antônio Carlos, Pampulha, Belo Horizonte – MG, 31270-901, Brazil; email: giliane@icb.ufmg.br

Detection of Vaccinia Virus in Urban Domestic Cats, Brazil


Technical Appendix.

Technical Appendix Table. Diagnostic results for 16 house cats with neutralizing antibodies for vaccinia virus, Belo Horizonte, Brazil, September 2012–December 2014*

Cat	PRNT ₇₀ titer (NU/mL)	qPCR C11R	qPCR A56R†	VACV strain	Clinical signs
1	1:320 (800)	+	+	VACV-BR group 2	Urethral obstruction
2	1:40 (100)	+	–	–	FeLV+
3	1:40 (100)	+	–	–	Hepatic lipidosis
4	1:160 (400)	+	+	VACV-BR group 1	None
5	1:320 (800)	–	–	–	Hepatic lipidosis
6	1:160 (400)	+	–	–	None
7	1:320 (800)	+	+	VACV-BR group 2	Urethral obstruction
8	1:40 (100)	–	–	–	FIV+
9	1:160 (400)	+	–	–	None
10	1:640 (1,600)	+	+	VACV-BR group 2	None
11	1:80 (200)	–	–	–	Sporotrichosis
12	1:80 (200)	+	+	VACV-BR group 1	<i>Mycoplasma haemofelis</i>
13	1:40 (100)	+	–	–	None
14	1:640 (1,600)	+	–	–	<i>Dipylidium caninum</i>
15	1:640 (1,600)	+	+	VACV-BR group 2	None
16	1:40 (100)	+	–	–	Uncharacterized skin lesions

*BR, Brazil; FeLV, feline leukemia virus; FIV, feline immunodeficiency virus; NU, neutralizing units; qPCR, real-time PCR; PRNT₇₀, ≥70% plaque-reduction neutralization test; VACV, vaccinia virus; +, positive; –, negative.

†Fragment sequenced.



Technical Appendix Figure 1. Map of Brazil highlighting studied areas in black. A) The 5 states in which house cats were sampled. N indicates the number of sampled house cats. B) Map of Belo Horizonte city (located in Minas Gerais State), where most of the house cats were sampled. The map shows the regions in Belo Horizonte including Pampulha and eastern regions. The numbers indicate the number of house cats sampled. Seropositive cats indicated with parentheses. C) Characteristics of Pampulha region. D) Characteristics of eastern regions, respectively.

```

#VACV-RR (AY243312)          GTA AGT ACA TCA TCT GGA ATT GTC ACT ACT AAA TCA ACC ACC GAT GAT GCG GAT CTT TAT GAT ACG TAC AAT GAT AAT GA- --- --T ACA GTA CCA CCA ACT ACT GTA GGC GGT AGT ACA ACC
#VACV-GP1V (DQ206436)
#VACV-YBH (DQ206435)
#VACV-SPAn22 (DQ222822)
#VACV-BeAns8058 (DQ206442)
#VACV-P1V (GU183769)
#Cat_1
#Cat_7
#Cat_10
#Cat_15
#VACV-ARAV (AY523994)
#VACV-CTGV (AF229247)
#VACV-CTGV-HD07 (FJ545689)
#VACV-GP2V (DQ206437)
#VACV-MURV (DQ247770)
#VACV-MARV (GQ226040)
#VACV-P2V (GU183770)
#VACV-SV2 (EF063677)
#VACV-TD_CA (GU322359)
#Cat_4
#Cat_12
#VACV-IOC (AF225248)
#VACV-BE13906 (AF375077)
#VACV-BE191 (AE375078)
#VACV-Ankara (U944848)
#HSPV (DQ79250)
#Rabbitpox (AF37511)
#HXPV-saire (DQ011155)
#HXPV-Congo (DQ011154)
#HXPV-Sierra_Leone (AY741551)
#CPXV-Srighton_Red (AF482758.2)
#CPXV-br (NC003663)
#CPXV-cpv-667-94h (AF377883)
#CPXV (AF377886)
#CPXV-89-4 (AF375085)
#CPXV-cpv-121800 (AF377877)
#VARV_Pakistan_1369 (DQ437589)
#VARV_Somalia_1977 (DQ441438)
#VARV_India_1564 (DQ437585)

```

Technical Appendix Figure 2. Nucleotide sequence of the vaccinia virus (VACV)–positive domestic cats A56R (hemagglutinin) genes (green) compared with homologous sequences of several other orthopoxviruses, Belo Horizonte, Brazil, September 2012–December 2014. Strains had the deletion region (red) conserved and were grouped with other VACVs (group 1 and 2) isolated in Brazil. The sample from Cat_1 showed 1 polymorphism (C–A) compared with other VACV isolates.