



## NOTE

Wildlife Science

# Ocular disease caused by the trematode *Philophthalmus lachrymosus* in free-living kelp gulls (*Larus dominicanus*) of Brazil

André Tavares SOMMA<sup>1)</sup>, Adriane STEUERNAGEL<sup>2)</sup>,  
Eduardo Alberto PULIDO-MURILLO<sup>3)</sup>, Hudson Alves PINTO<sup>3)</sup>,  
Larissa REIFUR<sup>4)</sup>, Bret A. MOORE<sup>5)</sup>, Fabiano MONTIANI-FERREIRA<sup>1)\*</sup>

<sup>1)</sup>Comparative Ophthalmology Lab, Department of Veterinary Medicine, Universidade Federal do Paraná, Curitiba, Brazil

<sup>2)</sup>Unidade de Estabilização de Animais Marinhos de Penha–Universidade do Vale do Itajaí (Univali), Praia de Armação do Itapocorói, Penha, Brazil

<sup>3)</sup>Laboratório de Biologia de Trematoda, Departamento de Parasitologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil

<sup>4)</sup>Departamento de Patologia Básica, Universidade Federal do Paraná, Curitiba, Brazil

<sup>5)</sup>College of Veterinary Medicine, Department of Small Animal Clinical Sciences, University of Florida, Gainesville, FL, USA

*J. Vet. Med. Sci.*

84(10): 1447–1452, 2022

doi: 10.1292/jvms.22-0193

Received: 22 April 2022

Accepted: 23 August 2022

Advanced Epub:

1 September 2022

**ABSTRACT.** In this study, clinical, parasitological and histopathological findings of thirteen kelp gulls (*Larus dominicanus*) found infected with eye flukes in Brazil are presented. Parasites detected in the ventral conjunctival fornix were identified as *Philophthalmus lachrymosus* [mean intensity of infection: 16 (5–36) worms/bird]. Eleven birds (85%) presented signs of systemic disease, such as emaciation, dehydration and depressed consciousness. Conjunctival hyperemia was observed in 22 eyes (85%). Keratitis, corneal ulcers, corneal abscess and chemosis were also detected in some eyes (4–8%). Histopathologic lesions, likely due to the parasite attachment to the conjunctiva, were found in the eyes of one infected bird that died from unrelated causes. Philophthalmosis by *P. lachrymosus* is here reported as a clinically relevant eye disease in kelp gulls.

**KEYWORDS:** conjunctival hyperemia, eye fluke, pathology, philophthalmosis, trematodes

Trematodes of the genus *Philophthalmus* are eye-flukes with cosmopolitan distributions found in birds and mammals [5, 6, 9, 26]. Zoonotic human cases of philophthalmosis have been reported in different continents, including Europe (Yugoslavia), Asia (Israel, Thailand, Sri Lanka, Japan) and North America (Mexico, United States) [12, 18, 20, 30, 38]. Depending on factors such as worm burden, parasite stage of development, and host species, philophthalmosis can be clinically relevant, causing ocular surface disease such as conjunctivitis [5, 27–29, 36, 38], keratitis [17, 21, 36] and blepharitis [5, 17]. Histological analysis of ocular trematodosis commonly reveals inflammatory infiltrates and the presence of adult parasites attached to the conjunctiva [7, 31]. In some cases, eye disease can be associated with severe corneal lesions resulting in blindness and loss of the eye [5].

Birds are considered the primary hosts of *Philophthalmus* spp. Dozens of avian species, from different phylogenetic orders (Galliformes [2, 5], Anseriformes [24, 39], Tinamiformes [29], Struthioniformes [7, 22, 23, 32, 37], Passeriformes [19] and Charadriiformes [3, 11, 14, 31, 36]) have been documented harboring these parasites. However, most reports are based on *post-mortem* examinations of the avian hosts, with few studies addressing clinical and histological details. Data on clinical and pathological findings related to this eye disease in gulls are scarce and are limited to recent reports of gulls from the Iberian Peninsula [14] and New Zealand [3]. In this study, the occurrence of philophthalmosis in kelp gulls (*Larus dominicanus*) from the southern coast of Brazil is reported. Besides identifying the etiological agent based on morphological and molecular data, we described the clinical, parasitological and histological findings related to philophthalmosis in this group of marine birds.

A total of 13 kelp gulls (*Larus dominicanus*) (12 adults and 1 juvenile) were admitted for rehabilitation at the Marine Animals Stabilization Unit of the University of the Itajaí Valley (UNIVALI) in the city of Penha, Santa Catarina State between July 2017 and September 2018. The age of the birds was classified according to the plumage color [15, 33]. Adults were considered to be at least 4-years-old, since full adult plumage is acquired only after the fourth year [15]. The birds originated from different locations along the coast of Santa Catarina state, between the cities of Barra Velha and Bombinhas (Fig. 1).

\*Correspondence to: Montiani-Ferreira F: montiani@ufpr.br, Veterinary Medicine Department, Comparative Ophthalmology Laboratory (LABOCO), Federal University of Paraná (UFPR), Setor de Ciências Agrárias, Rua dos Funcionários, 1540, CEP 80050-035, Curitiba-PR, Brazil

©2022 The Japanese Society of Veterinary Science



This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License. (CC-BY-NC-ND 4.0: <https://creativecommons.org/licenses/by-nc-nd/4.0/>)

All clinical diagnostics, medical procedures, and data acquisition were performed at the Marine Animals Stabilization Unit of UNIVALI as part of the Beach Monitoring Program—Santos Basin in the city of Penha, Santa Catarina State, located on the southern Brazilian coast. All procedures using live birds were conducted according to the Association for Research in Vision & Ophthalmology Statement for the Use of Animals in Ophthalmic and Vision Research and with Animal Use Committee of Federal University of Paraná. All birds were previously free-ranging and were collected by the UNIVALI staff due to debilitating health conditions, including signs of ocular disease.

All birds were subjected to a clinical evaluation protocol established by veterinarians from the Marine Animals Stabilization Unit. For the ophthalmic examinations, after physical restraint, the anterior segment structures were evaluated using a slit lamp biomicroscope (PSL One, Keeler, Malvern, PA, USA). A single drop of 2% fluorescein (Drogavet; Curitiba, PR, Brazil) was instilled to investigate for potential ocular surface ulcerations. The same investigator (ATS) conducted the ophthalmic examination in all 13 birds (26 eyes). Fundus examinations were not performed. Individual trematode parasites visually found in the conjunctival sac were removed using toothless Colibri ophthalmic forceps. The worms were fixed in ethanol for morphological and molecular evaluation.

Both enucleated eyes from an infected bird, which died from unrelated causes, were immersed in 10% formalin for 14 days, then transferred to 96% ethanol for storage. After fixation, the globes were sectioned in the sagittal (at the optic axis) and coronal planes for gross anatomic evaluation, and two sagittal hemi sections were collected and embedded in paraffin. Histologic sections of 5 µm each were then stained with hematoxylin and eosin. The slides were examined microscopically using a Leica DM500 microscope with the Leica Application software suite (LAZ EZ), version 2.0 (Leica Microsystems, Wetzlar, Germany). The bird was not submitted to a full necropsy. Only the eyes were sent for analysis at the Comparative Ophthalmology Lab (LABOCO-UFPR).

For the morphological study, ethanol-fixed worms were stained with alum acetocarmine, dehydrated in an ethanol series, cleared in beechwood creosote, and mounted on slides with Canada balsam. At least one parasite specimen from each infected bird, was processed and subjected to evaluation with a light microscope. Morphological and morphometric studies were performed under an Olympus BH2 light microscope (Olympus, Tokyo, Japan). Measurements were obtained using an eyepiece micrometer. Photographs were taken with a Leica ICC50 HD digital camera coupled to a Leica DM500 microscope and analyzed in the Leica Application software suite (LAZ EZ), version 2.0 (Leica Microsystems). Identification of the parasites was performed with the aid of previous works [6, 11, 16, 25, 39]. Samples of the parasites studied were deposited in the Collection of Trematodes of the Federal University of Minas Gerais (UFMG-TRE-124).

One ethanol-fixed parasite was used for molecular analysis. DNA was extracted using the Wizard® Genomic DNA Purification kit in accordance with the manufacturer's instructions. The concentration of the extracted DNA was estimated using a microvolume spectrophotometer (NanoDrop® Lite-1000). DNA amplifications were performed by PCR in a total volume of 25 µL. In each reaction, we used Platinum Hot Start Master Mix 2×, 10 µM of each primer and ~50 ng of DNA. Partial regions of the 28S rDNA (~1,200 bp) gene were amplified using primers dig12 (5'-AAGCATATCACTAAGCGG-3') and 1500R (5'-GCTATCCTGAGGGAAACTTCG-3'), which are widely used for characterization of trematodes, including for species of *Philophthalmus* [7]. The PCR conditions were those previously published by Tkach *et al.* [35]. Amplicons were sequenced in both directions by capillary electrophoresis in the ABI3730 sequencer, with the POP7 polymer and the BigDye v3.1 sequencing kit (Applied Biosystems, Inc., Foster City, CA, USA). Consensus sequences were assembled and edited using ChromasPro version 2.0.1 (Technelysium Pty Ltd., Tewantin, Australia) and compared with data available in GenBank.

Eleven birds presented signs of systemic disease [11/13 (85%)]. Emaciation was the most common finding during clinical evaluation [(8/13 (61%)). Five birds [5/13 (38%)] were dehydrated at admission and three birds [3/13 (23%)] showed depressed consciousness. Diarrhea was present in two individuals [2/13 (15%)] and cloacal prolapse and pododermatitis were found in one individual [1/13 (8%)]. The left hindlimb was missing in one individual examined. One bird died before any clinical diagnosis of systemic disease was concluded.

Parasites were found on both eyes (OU) in eleven birds [11/13 (85%)] in the left eye (OS) only in one bird (8%), and in the right eye (OD) in another bird (8%). A total of 206 parasites were found and removed. One hundred and nine (53%) parasites were found in OD and 97 (47%) were collected from OS. The mean intensity of infection was 16 worms/bird (varying from 5 to 32). All trematodes collected in this investigation were found in the ventral conjunctival fornix (Fig. 2A). Conjunctival hyperemia (Fig. 2B) was



**Fig. 1.** Map of Brazil (left) detaching the coastline of the southern state of Santa Catarina (right), between the cities of Barra Velha (north) and Bombinhas (south) in which 13 kelp gulls (*Larus dominicanus*) were collected and diagnosed with eye flukes.

the most prevalent ocular sign, observed in 22 eyes (85%). Hyperemia was more severe in eyes heavily infected by the parasites. Corneal edema (Fig. 2C), corneal abscess, and chemosis were each detected in one eye (4%). Keratitis was noted in two eyes (8%), corneal ulcer in two eyes (Fig. 2D) (8%). Detailed information about bird collecting sites, clinical exam findings and intensity of infection can be seen in Table 1.

Supportive care was provided according to the systemic condition of each bird. The treatment protocol of infected birds with conjunctivitis consisted only of the mechanical removal of the eye flukes. Birds with corneal lesions received chloramphenicol ophthalmic ointment (Novartis Biociência S/A, São Paulo, Brazil) according to the severity of each case until complete corneal healing, which have occurred in 5 to 7 days.


Hemi-section of the globes from a single individual revealed no intraocular abnormalities. The eye trematodes were not observed histologically in any of the sections. Histopathology of the eye and adnexa confirmed conjunctivitis. Microscopic lesions consisted of varying degrees of inflammation within the conjunctival tissue, with a diffuse mononuclear inflammatory cell infiltration in the conjunctival epithelium and substantia propria (Fig. 3). Caliciform cells typical of the conjunctival fornix were observed around these sites.

The worms were morphologically identified as *Philophthalmus lachrymosus* Braun, 1902 (Fig. 4). The morphology and measurements obtained in the present study are compatible to those reported to the species by different authors [11, 16, 25]. The morphometric data are presented in Table 2. The molecular analysis of the parasite sample confirmed the eye flukes evaluated in this study as a member of the genus *Philophthalmus*. In the 28S dataset (1,161 bp), obtained lineages showed 99.89% similarity with an isolate identified as *P. lachrymosus* found in Portugal [14]. The similarity with *Philophthalmus gralli* and *Philophthalmus lucipetus* were 98.62% and 98.44%, respectively. Considering other members of the family Philophthamidae (species of the marine genera *Cloacitrema* and *Parorchis*), the divergences were greater (93.96–96.55%).

*Philophthalmus lachrymosus*, sometimes spelled *lacrymosus* [11], is a species of eye fluke that has been reported to affect different birds and mammals around the world. This species is common in gulls, having already been reported in four species; brown-hooded gull (*Chroicocephalus maculipennis*) [11], black-headed gull (*Chroicocephalus ridibundus*) [26] and kelp gulls (*L. dominicanus*) [36] from Brazil, and lesser black-backed gull (*Larus fuscus*) from Portugal [14]. The present study represents the second report of this eyefluke in *L. dominicanus* from Brazil, with the first one being in a different location (Rio de Janeiro) [36]. Most prior reports of *P. lachrymosus* in gulls have been based on necropsy findings. The only exception is a recent characterization of the clinical, morphological and molecular aspects of philophthalmosis in gulls in Europe [14]. In the present study, ophthalmic findings are described in free living, wild-caught individuals. *Philophthalmus lachrymosus* may cause severe ocular lesions, including deep corneal ulcers, even blindness and loss of the globe [11, 14, 28]. Ocular surface lesions observed in our study were most likely due to the attachment of the ventral suckers of the eye fluke [7, 26]. The most frequent ocular clinical sign observed was conjunctival hyperemia (85%), similar to what has been previously described in other works with *Philophthalmus* spp. [7, 28, 29]. Subjectively the severity of hyperemia appears to

correspond with the number of parasites found in each eye. It was not possible to determine if the corneal ulcers present in one case were primary or secondary, although it was thought to be due to self-inflicted trauma secondary to the conjunctivitis and the presence of parasites. Overall, the clinical features described in the present study were supportive of previous studies [7, 14, 19, 22, 23, 27–29].

The observed ocular lesions, such as conjunctival chemosis, hyperemia, keratitis, corneal ulcers and abscess, were very likely a direct consequence of the parasite causing chronic inflammation, irritation and consequent self-traumatization. Previously, ocular lesions have only been described in general clinical terms, such as citing the presence of red eye, conjunctivitis, blepharoconjunctivitis, lacrimation, swollen eyelids, and purulent discharge [7, 22, 37]. More severe conditions such as eye destruction and blindness were reported in tinamous with an ocular infection by *Philophthalmus gralli* [29], likely to be a consequence of self-traumatism. Regarding systemic clinical signs, emaciation was reported in cases of *Philophthalmus* spp. infection in both birds [22] and mammals [28]. There is no evidence indicating that the signs of systemic disease observed here, such as dehydration,




**Fig. 2.** Selected ophthalmic clinical signs in kelp gulls (*Larus dominicanus*) infected with *Philophthalmus lachrymosus*. **A:** Left ventral conjunctival fornix showing moderate conjunctival hyperemia and numerous eye-flukes. **B:** Left eye demonstrating conjunctival hyperemia and chemosis in the dorsal bulbar conjunctiva, near the limbus. **C:** A right eye demonstrating diffuse corneal edema. **D:** Right eye showing a corneal ulcer stained with fluorescein dye.



**Table 1.** Clinical aspects of 13 kelp gulls (*Larus dominicanus*) infected with eye-flukes in the coast of Santa Catarina state, Brazil

Case ID	Date admission	Age	City/Beach	Clinical exam	Ophthalmic exam	Number of worms	
						OD	OS
46877	07/18/2017	Adult	Piçarras	Emaciated	Conjunctival hyperemia OU	13	19
56869	08/20/2017	Adult	Barra Velha/Bacia da Vovó	Emaciated	Conjunctival hyperemia OS	0	5
47199	08/25/2017	Adult	Navegantes	Dehydrated	Conjunctival hyperemia OU	11	7
31315	10/15/2017	Adult	Penha/ Armação	-	Conjunctival hyperemia OU	2	6
75996	12/31/2017	Adult	Itajaí/ Cabeçudas	Emaciated Left hindlimb missing	Conjunctival hyperemia OD Keratitis OU	12	0
38364	04/01/2018	Adult	Porto Belo/Baxio	-	Conjunctival hyperemia Chemosis OS	7	9
38361	03/26/2018	Adult	Bombinhas/Canto Grande	Depressed consciousness Dehydrated Emaciated	Conjunctival hyperemia OU	5	4
33120	03/29/2018	Young	Navegantes	Depressed consciousness Dehydrated Emaciated	Conjunctival hyperemia OU Keratitis OS Corneal ulcer OD Corneal abscess OS	5	11
33128	03/31/2018	Adult	Piçarras	-	Conjunctival hyperemia OU Corneal abscess	13	3
74058	04/06/2018	Adult	Piçarras	Emaciated Diarrhea	Conjunctival hyperemia OU	16	8
38365	04/17/2018	Adult	Bombinhas/ Zimbros	Dehydrated Emaciated	Conjunctival hyperemia OU	12	6
90995	09/15/2018	Adult	Bombinhas/ Zimbros	Depressed consciousness Diarrhea Cloacal prolapse	Conjunctival hyperemia OU Keratitis OS	2	4
115062 <sup>+</sup>	09/18/2018	Adult	Penha/Armação	Dehydrated Pododermatitis	Conjunctival hyperemia OU	11	15

<sup>+</sup> Bird died during the period of the study. OU: both eyes, OS: left eye, OD: right eye.



**Fig. 3.** A photomicrograph showing a conjunctival lesion present in the eye of a representative kelp gull (*Larus dominicanus*), bird #115062, infected with *Philophthalmus lachrymosus*. Conjunctivitis was evidenced by the presence of a diffuse mononuclear inflammatory cell infiltration in the conjunctival epithelium and substantia propria. Stain: H&E. Scale bar: 50  $\mu$ m.



**Fig. 4.** *Philophthalmus lachrymosus* found in ventral conjunctival fornix of kelp gulls (*Larus dominicanus*). Whole view of a stained worm. Scale bar: 1 mm.

depressed consciousness and emaciation were directly caused by the presence of the parasites. Nevertheless, it is very likely that the irritation, and stress and even visual deficit may have contributed to the poor overall condition of these birds. In fact, ostriches heavily infected with *P. gralli* presented a reduced body condition [22]. Tinamous with ophthalmic infections also caused by *P. gralli* were reported presenting anorexia and debility [29]. Although a detailed study is still required, it appears that the overall health condition of kelp gulls infected with *Philophthalmus* spp. is poorer than that of uninfected gulls of the population routinely examined. However, it was not possible to determine whether the ocular and systemic conditions were caused solely by the trematode infection

**Table 2.** Morphometric data of *Philophthalmus lachrymosus* collected from 13 kelp gulls (*Larus dominicanus*) in the Coast of Santa Catarina State, Brazil, between July of 2017 and September of 2018

Host Location	N	Present study <i>Larus dominicanus</i> Brazil	
		13	
Body	L	4,282 ± 502 (3,386–5,134)	
	W	1,549 ± 196 (1,274–1,893)	
Oral sucker (OS)	L	310 ± 40 (248–397)	
	W	407 ± 49 (340–475)	
Pharynx (PH)	L	415 ± 40 (355–468)	
	W	306 ± 45 (248–425)	
Ventral sucker (VS)	L	664 ± 45 (581–716)	
	W	694 ± 26 (659–730)	
OS/VS	L	1:2.2 ± 0.3 (1.6–2.6)	
	W	1:1.7 ± 0.2 (1.4–2.1)	
OS/ PH	L	1:1.3 ± 0.1 (1.2–1.6)	
	W	1:0.8 ± 0.1 (0.6–1.1)	
Ovary	L	203 ± 28 (163–227)	
	W	254 ± 29 (206–284)	
Cirrus sac	L	948 ± 229 (610–1,361)	
	W	173 ± 47 (106–227)	
Anterior testis	L	417 ± 115 (206–624)	
	W	537 ± 163 (241–794)	
Posterior testis	L	451 ± 100 (255–610)	
	W	515 ± 133 (291–744)	
Vitellaria	%	53 ± 10 (32–70)	
	Eggs	L	75 ± 5 (63–84)
W		34 ± 3 (30–41)	

L: length; W: width. All measurements are given in µm, with the range followed by the mean and standard deviation.

philophthalmosis has not been reported in the continent thus far. Nevertheless, since the zoonotic potential cannot be discarded, studies involving the elucidation of aspects of the philophthalmosis in synanthropic birds, such as *L. dominicanus*, may help to understand this type of host-parasite environment, and thus any direct preventative and control measures.

The possible impact of philophthalmosis on the ecological traits of gulls highlights the clinical relevance of the disease shown in the present study. On the other hand, advances in the knowledge of the biology of *P. lachrymosus*, including the mollusk species that acts as a first intermediate host, are still necessary. Thus, studies focusing in the elucidation of the parasite life cycle and the process of infection of kelp gulls are necessary and encouraged. Finally, additional studies on the relationship between the systemic conditions commonly noted in infected individuals, and whether they were caused solely by the trematode infection or if ocular philophthalmosis occurred secondary to the existence of systemic disease. Results obtained in this study reveal that philophthalmosis by *P. lachrymosus* is a clinically relevant ophthalmic condition in *L. dominicanus* of Brazil.

**CONFLICTS OF INTEREST.** The authors declare no conflicts of interest.

**ACKNOWLEDGMENTS.** The authors would like to thank all the staff of the Marine Animals Stabilization Unit of the University of the Itajaí Valley (UNIVALI) in the city of Penha and CAPES, especially Programa Institucional de Internacionalização (CAPES-PrInt), for the support. This work was also supported by the National Council for Scientific and Technological Development (CNPq), Brazil (student scholarship to EAPM, and research scholarship to HAP).

## REFERENCES

- Abrams RW. 1983. Pelagic seabirds and trawl-fisheries in the southern Benguela Current region. *Mar Ecol Prog Ser* **11**: 151–156. [CrossRef]
- Assis JCA, Martins NRS, Pinto HA. 2018. Experimental avian philophthalmosis: Evaluation of diagnosis and treatment of chickens infected with *Philophthalmus gralli* (Trematoda: Philophthalmidae). *Vet Parasitol* **256**: 24–28. [Medline] [CrossRef]
- Bennett J, Presswell B. 2019. Morphology and molecules resolve the identity and life cycle of an eye trematode, *Philophthalmus attenuatus* n. sp. (Trematoda: Philophthalmidae) infecting gulls in New Zealand. *Parasitol Res* **118**: 1501–1509. [Medline] [CrossRef]

4. Branco JO, Fracasso HA, Barbieri E. 2009. Breeding biology of the kelp gull (*Larus dominicanus*) at Santa Catarina coast, Brazil. *Ornitol Neotrop* **20**: 409–419.
5. Chalkowski K, Morgan A, Lepczyk CA, Zohdy S. 2021. Spread of an avian eye fluke, *Philophthalmus gralli*, through biological invasion of an intermediate host. *J Parasitol* **107**: 336–348. [Medline] [CrossRef]
6. Ching HL. 1961. The development and morphological variation of *Philophthalmus gralli* Mathis and Leger, 1910 with a comparison of species of *Philophthalmus* Looss, 1899. *Proc Helminthol Soc Wash* **28**: 130–138.
7. Church ML, Barrett PM, Swenson J, Kinsella JM, Tkach VV. 2013. Outbreak of *Philophthalmus gralli* in four greater rheas (*Rhea americana*). *Vet Ophthalmol* **16**: 65–72. [Medline] [CrossRef]
8. Crawford RJ, Cooper J, Shelton PA. 1982. Distribution, population size, breeding and conservation of the Kelp Gull in southern Africa. *Ostrich* **53**: 164–177. [CrossRef]
9. Dailey M, Ellin R, Parás A. 2005. First report of parasites from pinnipeds in the Galapagos Islands, Ecuador, with a description of a new species of *Philophthalmus* (Digenea: Philophthalmidae). *J Parasitol* **91**: 614–617. [Medline] [CrossRef]
10. Ebert LA, Schlemper JC, Pelisser MR, Pereira BD, Silva MA, Branco JO. 2016. Pathogenic bacteria associated with kelp gull *Larus dominicanus* (Charadriiformes, Laridae) on the coast of Santa Catarina State-Brazil. *Int J Curr Microbiol Appl Sci* **5**: 458–473. [CrossRef]
11. Freitas JF. 1955. Sobre dois trematódeos parasitos de aves: *Philophthalmus lachrymosus* Braun, 1902 e *Renicola mirandaribeiroi* n. sp. *Arq Mus Nac* **42**: 585–610.
12. Gutierrez Y, Grossniklaus HE, Annable WL. 1987. Human conjunctivitis caused by the bird parasite *Philophthalmus*. *Am J Ophthalmol* **104**: 417–419. [Medline] [CrossRef]
13. Heneberg P, Casero M. 2022. Recent large-scale philophthalmosis outbreak in Portugal: inefficacy of common antihelminthic agents. *Parasit Vectors* **15**: 167. [Medline] [CrossRef]
14. Heneberg P, Casero M, Waap H, Sitko J, Azevedo F, Těšínský M, Literák I. 2018. An outbreak of philophthalmosis in *Larus michahellis* and *Larus fuscus* gulls in Iberian Peninsula. *Parasitol Int* **67**: 253–261. [Medline] [CrossRef]
15. Jiguet F, Jamarillo A, Sinclair I. 2001. Identification of kelp gull. *Bird World* **14**: 112–125.
16. Jones A, Bray RA, Gibson DI eds. 2005. Keys to the Trematoda, Vol. 2, The Natural History Museum, London, CABI Publishing, Wallingford.
17. Lamothe-Argumedo R, Diaz-Camacho SP, Nawa Y. 2003. The first human case in Mexico of conjunctivitis caused by the avian parasite, *Philophthalmus lacrimosus*. *J Parasitol* **89**: 183–185. [Medline] [CrossRef]
18. Lang Y, Weiss Y, Garzozzi H, Gold D, Lengy J. 1993. A first instance of human philophthalmosis in Israel. *J Helminthol* **67**: 107–111. [Medline] [CrossRef]
19. Literák I, Heneberg P, Sitko J, Wetzel EJ, Cardenas Callirgos JM, Čapek M, Valle Basto D, Papoušek I. 2013. Eye trematode infection in small passerines in Peru caused by *Philophthalmus lucipetus*, an agent with a zoonotic potential spread by an invasive freshwater snail. *Parasitol Int* **62**: 390–396. [Medline] [CrossRef]
20. Marković A. 1939. Der erste Fall von Philophthalmose beim Menschen. *Albrecht Von Graefes Arch Ophthalmol* **140**: 515–526. [CrossRef]
21. Mimori T, Hirai H, Kifune T, Inada K. 1982. *Philophthalmus* sp. (Trematoda) in a human eye. *Am J Trop Med Hyg* **31**: 859–861. [Medline] [CrossRef]
22. Mukaratirwa S, Hove T, Cindzi ZM, Maononga DB, Taruvinga M, Matenga E. 2005. First report of a field outbreak of the oriental eye fluke, *Philophthalmus gralli*, in commercially raised ostriches (*Struthio camelus*) in Zimbabwe. *Onderstepoort J Vet Res* **72**: 203–206. [Medline] [CrossRef]
23. Mukaratirwa S, Chimbwanda M, Matekwe N, Matenga E. 2008. A comparison of the efficacy of doramectin, closantel and levamisole in the treatment of the ‘oriental eye fluke’, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*). *J S Afr Vet Assoc* **79**: 101–103. [Medline] [CrossRef]
24. Muniz-Pereira LC, Amato SB. 1993. *Philophthalmus gralli* (Digenea: Philophthalmidae) parasite of *Anas bahamensis* and *Amazonetta brasiliensis*, from lagoons of Maricá county, Rio de Janeiro, Brazil. *Mem Inst Oswaldo Cruz* **88**: 567–569. [CrossRef]
25. Nasir P, Diaz MT. 1972. Avian flukes of Venezuela. *Riv Parassitol* **33**: 245–276.
26. Nollen PM, Kanev I. 1995. The taxonomy and biology of philophthalmid eye flukes. *Adv Parasitol* **36**: 205–269. [CrossRef]
27. Phillips BE, Pérez-Rosas D, Flowers JR, Cullen JM, Law JM, Colitz C, Deresienski D, Lohmann KJ, Lewbart GA. 2018. Evaluation of the ophthalmic disease and histopathologic effects due to the ocular trematode *Philophthalmus zalophi* on juvenile Galapagos sea lions (*Zalophus wollebaeki*). *J Zoo Wildl Med* **49**: 581–590. [Medline] [CrossRef]
28. Pinto RM, dos Santos LC, Tortelly R, Menezes RC, de Moraes W, Juvenal JC, Gomes DC. 2005. Pathology and first report of natural infections of the eye trematode *Philophthalmus lachrymosus* Braun, 1902 (Digenea, Philophthalmidae) in a non-human mammalian host. *Mem Inst Oswaldo Cruz* **100**: 579–583. [Medline] [CrossRef]
29. Rojas D, Soto C, Rojas A. 2013. Pathology and first report of natural eye infection with the trematode *Philophthalmus gralli* (Digenea, Philophthalmidae) in *Tinamus major* (Tinamiformes, Tinamidae), Costa Rica. *Parasitol Int* **62**: 571–574. [Medline] [CrossRef]
30. Sapp SGH, Alhabshan RN, Bishop HS, Fox M, Ndubuisi M, Snider CE, Bradbury RS. 2019. Ocular trematodiasis caused by the avian eye fluke *Philophthalmus* in southern Texas. *Open Forum Infect Dis* **6**: ofz265. [Medline] [CrossRef]
31. Schmidt RE, Toft JD 2nd. 1981. Ophthalmic lesions in animals from a zoologic collection. *J Wildl Dis* **17**: 267–275. [Medline] [CrossRef]
32. Schuster RK. 2011. *Philophthalmus aweerensis* n. sp. (Trematoda: Philophthalmidae) found in a rhea (*Rhea americana*) in the United Arab Emirates. *Parasitol Res* **109**: 1029–1033. [Medline] [CrossRef]
33. Sick H, Barruel P. 1984. *Ornitologia Brasileira*, Vol. 1, Editora Universidade de Brasília, Brasília.
34. Silva-Costa A, Bugoni L. 2013. Feeding ecology of kelp gulls (*Larus dominicanus*) in marine and limnetic environments. *Aquat Ecol* **47**: 211–224. [CrossRef]
35. Tkach VV, Littlewood DT, Olson PD, Kinsella JM, Swiderski Z. 2003. Molecular phylogenetic analysis of the Microphalloidea Ward, 1901 (Trematoda: Digenea). *Syst Parasitol* **56**: 1–15. [Medline] [CrossRef]
36. Travassos L, Freitas JFT, Mendonça JM, Rodrigues HO. 1960. Excursão a Cabo Frio, Estado do Rio de Janeiro. *Atas Soc Biol Rio de Janeiro* **4**: 70–71 (in Portuguese).
37. Verocai GG, Lopes LN, Burlini L, Correia TR, de Souza CP, Coumendouros K. 2009. Occurrence of *Philophthalmus gralli* (Trematoda: Philophthalmidae) in farmed ostriches in Brazil. *Trop Anim Health Prod* **41**: 1241–1242. [Medline] [CrossRef]
38. Waikagul J, Dekumyoy P, Yoonuan T, Praevanit R. 2006. Conjunctiva philophthalmosis: a case report in Thailand. *Am J Trop Med Hyg* **74**: 848–849. [Medline] [CrossRef]
39. Yamaguti S. 1971. Synopsis of Digenetic Trematodes of Vertebrates, Vols I and II, Keigaku Publishing Co., Tokyo.
40. Yorio P, Branco JO, Lenzi J, Luna-Jorquera G, Zavalaga C. 2016. Distribution and trends in kelp gull (*Larus dominicanus*) coastal breeding populations in South America. *Waterbirds* **39**: 114–135. [CrossRef]