



Original Article

What physical habitat factors determine the distribution of gastropods in neotropical headwater streams?

Marden S. Linares^{a,*}, Diego R. Macedo^b, Rodrigo L. Massara^{c,d}, Marcos Callisto^a

^a Universidade Federal de Minas Gerais, Instituto de Ciências Biológicas, Departamento de Genética, Ecologia e Evolução, Laboratório de Ecologia de Bentos, Av. Antônio Carlos 6627, Pampulha, CP 486, Belo Horizonte, MG, CEP 30270-901, Brazil

^b Universidade Federal de Minas Gerais, Instituto de Geociências, Departamento de Geografia, Av. Antônio Carlos 6627, Pampulha, Belo Horizonte, MG, CEP 30270-901, Brazil

^c Universidade Federal de Minas Gerais, Instituto de Ciências Biológicas, Departamento de Genética, Ecologia e Evolução, Laboratório de Ecologia e Conservação, Programa de Pós-graduação em Ecologia, Conservação e Manejo da Vida Silvestre, Av. Antônio Carlos 6627, Pampulha, CP 486, Belo Horizonte, MG, CEP 30270-901, Brazil

^d Instituto SerraDiCal de Pesquisa e Conservação, Rua José Hemetério de Andrade, No. 570, Burtis, Belo Horizonte, MG, CEP 30493-180, Brazil

ARTICLE INFO

Keywords:

Lotic ecosystems
Alien species
Water quality
Freshwater gastropod conservation

ABSTRACT

Freshwater gastropods play a pivotal role in the structure and functioning of freshwater ecosystems, but despite their importance, there are still gaps in their ecology. Our goal was to understand what physical habitat factors are the most important for the distribution of freshwater gastropods in headwater stream ecosystems in the Neotropical Savanna and provide a baseline for conservation and management efforts for freshwater gastropods in this biome. We identified five taxa, each with different environmental preferences. *Littoridina* presence related negatively with stream slope, *Biomphalaria* presence related positively with total dissolved solids and the proportions of cobble, fine sediment and organic matter. *Gundlachia* presence correlated positively with elevation and the proportion of pools in the site. *Physa* presence related negatively with total dissolved solids and positively with alkalinity. *Melanoides tuberculata* presence correlated positively with the proportion of coarse gravel. Our results highlight the challenge for protecting native freshwater gastropod assemblages (and managing non-native invasive species), because environmental preferences vary widely amongst taxa and most are vulnerable to common anthropogenic disturbances.

1. Introduction

Rivers, lakes, ponds and other freshwater ecosystems cover less than 1% of the planet's surface area but hold more than 10% of the known species diversity (Dudgeon et al., 2006; Reid et al., 2019). Freshwater ecosystems also provide irreplaceable ecosystem services to human beings, including drinking water, water supply for multiple human uses, and hydroelectric power generation (van Rees et al., 2021). In spite of these contributions and their importance for global biodiversity and human well-being, freshwater ecosystems are among the most threatened ecosystems worldwide (Collen et al., 2014; Darwall et al., 2018; Reid et al., 2019; Turak et al., 2016). Among these, headwater streams (1st to 3rd order; Strahler, 1957) can be singled out for their high biotic diversity and taxa richness (Meyer et al., 2007) and because they represent ~80% of the channel length in a hydrographic basin (Datry et al., 2014).

Among the taxa that constitute headwater stream biodiversity, freshwater gastropods are one of the most diverse and widespread, with more than 4000 identified species living in all continents (Lopes-Lima et al., 2021; Strong et al., 2008). Freshwater gastropods play a pivotal role in the structure and functioning of lotic ecosystems, contributing to the biodiversity of benthic macroinvertebrate assemblages, water quality, nutrient cycling, and secondary productivity, because of their roles as filter feeders, algal grazers, detritivores and food for predators (Böhm et al., 2021; Johnson et al., 2013). Invasive gastropods are one of the most successful groups of non-native invasive species in freshwater ecosystems (e.g., *Melanoides tuberculata*, an Asian species that has been recorded in most South American river systems (Fernandez et al., 2003; Santos and Eskinazi-Sant'Anna, 2010)). Such species are capable of reproducing and generating fertile offspring with a high probability of survival. They also often act as ecosystem engineers, altering the physical habitats and biological processes of the ecosystems where they become

* Corresponding author.

E-mail addresses: mslinares@ufmg.br (M.S. Linares), diegorm@ufmg.br (D.R. Macedo), rmassara@gmail.com (R.L. Massara), callistom@ufmg.br (M. Callisto).

<https://doi.org/10.1016/j.watbs.2022.100076>

Received 20 April 2022; Received in revised form 26 July 2022; Accepted 25 August 2022

Available online 31 August 2022

2772-7351/© 2022 The Authors. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

established (Jones et al., 1994; Linares et al., 2020; Simberloff et al., 2013).

Despite their importance, there are still gaps regarding our knowledge of freshwater gastropod ecology (Lopes-Lima et al., 2021). The distribution and structure of gastropod assemblages is influenced by local environmental factors and given their limited movement ability, knowledge about the environmental factors determining their distribution is an important step for conserving native species and managing non-native invasive species (Bae and Park, 2020; Brown et al., 1998; Jörger et al., 2010; Strong et al., 2008).

This lack of knowledge is especially dire in the neotropics, where most studies on these taxa focus only on a few invasive species in reservoirs, large rivers and wetlands (e.g., Linares et al., 2019; Oliveira et al., 2011; Santos and Eskinazi-Sant’Anna, 2010), neglecting the roles of gastropods in headwater streams. For the Cerrado (Brazilian Neotropical Savanna), this study will be the first focusing on freshwater gastropod assemblages in headwater streams.

Here, our goal was to understand what physical habitat factors are the most important for the distribution of freshwater gastropods in headwater stream ecosystems in the Neotropical Savanna. We aimed to provide a baseline for conservation and management efforts for freshwater gastropods in the Neotropical Savanna.

2. Methodology

2.1. Study area

We sampled 40 stream sites (1st to 3rd order, based on Strahler, 1957) within a hydrological unit 35 km upstream of the Três Marias hydroelectric power reservoir, located in the Neotropical Savanna of southeastern Brazil (Fig. 1). The Três Marias Reservoir water comes primarily from the São Francisco River and its tributaries, such as the São Vicente, Paraopeba, Sucuriú, Indaiá, Ribeirão do Boi, Ribeirão da

Extrema and Borrachudo Rivers (Esteves et al., 1985). It began operating in 1962 to control floods, improve navigation, encourage development and irrigation, and generate hydroelectric power (CEMIG - Companhia Energética de Minas Gerais, 2014). The stream sites were selected using a probability-based procedure, resulting in a spatially balanced design (Macedo et al., 2014a; Olsen and Peck, 2008). The sites were sampled during the dry season of 2010 and were 40 mean channel widths long or a minimum of 150 m.

2.2. Water quality & physical habitat assessment

To assess water quality, the following physical and chemical characteristics of the water column were measured in the field for each site: pH, conductivity, and total dissolved solids (TDS). Water samples were collected for further analysis in the laboratory for dissolved oxygen, turbidity, alkalinity, total nitrogen, and total phosphorus. These analyses were conducted following APHA (1998) methodology.

To assess physical habitat condition, we followed USEPA protocols (Peck et al., 2006) adapted for the Neotropical Savanna (Ligeiro et al., 2013). In each sampling site, 11 transects (perpendicular to the channel) were established, defining 10 longitudinal sections where physical habitat measurements were taken. At each transect, we recorded channel morphology, habitat features, riparian structure, and anthropogenic alterations. Subsequently, these data were used to calculate 288 physical habitat metrics (Kaufmann et al., 1999, 2022) (Supplementary Material S1).

From the entire pool of physical habitat metrics calculated, we eliminated those with a high number of zeros ($0 \text{ on } \geq 0.8$ of the sites) and those with low variation among sites (coefficient of variation ≤ 0.3). We then determined the Pearson correlations among the remaining metrics, eliminating those that were highly correlated ($|r| \geq 0.7$; Dormann et al., 2013) and retaining the most ecologically relevant based on previous knowledge (Esselman et al., 2013; Macedo et al., 2016; Silva et al., 2017).




Fig. 1. Sampling site locations in the Três Marias hydrological unit.

We also estimated anthropogenic disturbances at local and catchment scales, based on the methodology described by Ligeiro et al. (2013). The Local Disturbance Index (LDI) was calculated based on the amount and proximity of 11 types of anthropogenic disturbances observed in-channel or in the riparian zone of each site (Kaufmann et al., 1999, 2022). Those disturbances included walls, dikes, revetments, rip rap or dams, buildings, pavement or cleared lots, roads or railroads, inlet or outlet pipes, landfills or trash, parks or maintained lawns, row crops, pastures, rangelands, hay fields, or evidence of livestock, logging, or mining.

For calculating the Catchment Disturbance Index (CDI), we estimated the land use in the catchment of each stream site. Land uses were delimited using images obtained through a TM sensor onboard the Landsat 5 satellite, taken in 2010. Polygons for the definition and quantification of land use categories were delimited and classified (pasture, agriculture and buildings), and computed as percentages. Images obtained from Google Earth satellite images were employed as ancillary data in this assessment (Macedo et al., 2014b). CDI was estimated using the land use percentages weighted by the disturbance potential of each (Böhmer et al., 2004; Ligeiro et al., 2013; Rawer-Jost et al., 2004) and calculated as:

$$CDI = 4 \times (\% \text{ residential} + \text{urban}) + 2 \times (\% \text{ agricultural areas} + \% \text{ bare soil}) + (\% \text{ pasture} + \% \text{ Eucalyptus plantations}).$$

2.3. Freshwater Gastropoda sampling

We collected freshwater gastropods at all stream sites using a D-net (30 cm opening, 0.5 mm mesh sieve), using a kick-net method. The samples were taken following a zig-zag pattern along each stream site, resulting in eleven sub-samples (0.09 m² each), totaling a 0.99 m² multi-habitat composite sample for each stream site. We fixed the samples in the field with 10% formalin and took them to the laboratory. There, the gastropods were rinsed in tap water, sorted, and identified to the lowest possible level with the aid of specialized literature (Mugnai et al., 2010; Simone, 2006).

Because of the fixation method, the specimens' soft tissue was damaged and could not be used in the identification process, preventing us from identifying all of them to species. Instead, identifications were based on the shells (Geiger et al., 2007; Meier-Brook, 1976; Paraense, 1975). Gastropoda were identified to genus, except for *Melanooides tuberculata*, which were identified to species. Sampling sites were then classified with each identified freshwater gastropod taxon as detected (1) or non-detected (0). All specimens collected were fixed in 70% alcohol and deposited in the Reference Collection of Benthic Macroinvertebrates, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais.

2.4. Data analyses

To understand which physical habitat factors were most important for the distribution of freshwater gastropod species in headwater stream ecosystems in the Três Marias hydrologic unit, we used a model selection approach based on the Akaike information criterion adjusted for small sample size (AICc; Burnham and Anderson, 2002). We used our selected variables as predictor variables in generalized linear models (GLMs) with a binomial distribution, where the response variables were the detection (1)/non-detection (0) of each gastropod taxon. Before the analyses we standardized all variables, which consisted of subtracting each variable value by the variable mean and dividing it by the variable standard deviation. As a result, all variables were centered so that the predictors had their means at 0, which allowed the units of the regression coefficients to be the same. The variables were standardized or scaled using the "scale" function in R. We then constructed models based on all possible additive variable combinations that may have influenced the probabilities of presence for each gastropod taxon (Doherty et al., 2012). This allowed us

to calculate the cumulative AICc weights (w_{+}) for each variable and evaluate which were the most likely ($w_{+} \geq 0.50$; Berger and Barbieri, 2004) to have influenced the presence for each gastropod taxon (Burnham and Anderson, 2002). Statistical analyses were implemented using the MuMIn package (Bartón, 2019) in R. To test for overdispersion or extra-binomial variation (i.e., $\hat{c} > 1$ and $p < 0.05$) in the most parameterized model of each model set, we used a simulation-based approach available in the DHARMA package (Hartig, 2022). Because the sampling site locations may have resulted in a lack-of-independence (or spatial autocorrelation) among our sites, we performed a Moran's I test for spatial autocorrelation (Lecocq et al., 2019; Smeraldo et al., 2020). To do so, we used the residuals of our most parameterized model in each model set, using the spdep package (version 1.1.2; Bivand and Wong, 2018) in R (R Development CoreTeam, 2015).

3. Results

A total of five freshwater gastropod taxa were sampled in our stream sites (Supplementary Material S2): *Littoridina* (Hydrobiidae - Caenogastropoda; 40 individuals detected in three sites), *Biomphalaria* (Planorbidae - Panpulmonata; 96 individuals detected in nine sites), *Gundlachia* (Planorbidae - Panpulmonata; 87 individuals detected in eight sites), *Physa* (Physidae - Panpulmonata; 10 individuals detected in five sites), and *Melanooides tuberculata* (Thiaridae - Caenogastropoda; 21 individuals detected in four sites). In total, we detected freshwater Gastropods in 18 (out of 40) sites (Fig. 2).

The variable selection process resulted in a total of 27 selected variables (Table 1). We organized them into six categories: Catchment metrics (two metrics), Physio-Chemical metrics (four metrics), Channel Morphology metrics (five metrics), Substrate metrics (nine metrics), Water Flow metrics (five metrics) and Disturbance metrics (two metrics).

Different variables influenced the presence/absence of freshwater gastropods (Table 2). *Littoridina* presence correlated negatively with Mean Site Slope. *Biomphalaria* presence correlated positively with Total Dissolved Solids and the proportions of Cobble, Fine Sediment and Organic Matter. *Gundlachia* presence correlated positively with Basin Elevation Range and the Proportion of Pools in the site. *Physa* presence correlated negatively with Total Dissolved Solids and positively with Alkalinity. *M. tuberculata* presence correlated positively with the proportion of Coarse Gravel. The simulation-based approach revealed no evidence of overdispersion in the tested models (i.e., $\hat{c} \sim 1$ and $p < 0.05$ in all cases; Supplementary Material S3). The Moran's I test for spatial autocorrelation did not show any spatial autocorrelation in the tested models (Supplementary Material S4).

4. Discussion

Littoridina was negatively related to channel slope, likely indicating a preference for flat lowland streams. This preference stems from these streams being comparatively more stable habitats, that are less affected by seasonal effects such as flash floods (Chattopadhyay et al., 2021; Erba et al., 2020), and the same pattern can be applied to *Gundlachia* preference for pools. *Littoridina* preference for lowland streams can have important implications for its conservation, given that lowland freshwater ecosystems are disproportionately affected by anthropogenic disturbances when compared to highland streams (de Vries et al., 2019; Oliveira et al., 2011; Usseglio-Polatera and Beisel, 2002). Lowland streams are easily accessible to humans and more often located near urban centers or agricultural areas. This results in greater occurrence of anthropogenic stressors, including altered basin land use and riparian vegetation, eutrophication, damming, channelization, and introduced invasive species (Graf et al., 2016; Horsák et al., 2009; Shandas and Alberti, 2009).

The positive correlation of *Melanooides tuberculata* with stable hard substrate corroborates previous studies in South American reservoirs (Karatajev et al., 2007; Linares et al., 2020), where high densities of




Fig. 2. Spatial distribution of presence/absence of freshwater gastropods in the Três Marias hidrologic unit.

Table 1

Metrics used to determine which physical habitat metrics were most important for the distribution of freshwater Gastropoda in Neotropical Savana streams.

Variable Code	Variable	Metric Category
Basin_elev_range	Basin Elevation Range (m)	Catchment
Basin_slope_range	Basin Slope Range (m)	Catchment
Turbidity	Turbidity (NTU)	Physio-Chemical
Conductivity	Conductivity (µS/cm)	Physio-Chemical
TDS	Total Dissolved Solids (mg/L)	Physio-Chemical
Alkalinity	Alkalinity (µEq/L of CO2)	Physio-Chemical
XBKF_D	Mean Thalweg Depth (m)	Channel Morphology
XBFW_D_RAT	Mean Width/Thalweg Depth	Channel Morphology
XWD_RAT_P	Mean Section Width/Section Depth	Channel Morphology
XCDENMID	Mean Channel Canopy Cover (%)	Channel Morphology
XSLOPE	Mean Site Slope (m)	Channel Morphology
XEMBED	Mean Total Embeddedness (%)	Substrate
PCT_BL	Boulder Percentage (%)	Substrate
PCT_CB	Cobble Percentage (%)	Substrate
PCT_GC	Coarse Gravel Percentage (%)	Substrate
PCT_GF	Fine Gravel Percentage (%)	Substrate
PCT_SA	Sand Percentage (%)	Substrate
PCT_FN	Fine Substrate Percentage (%)	Substrate
PCT_ORG	Organic Matter Percentage (%)	Substrate
PCT_WD	Wood Percentage (%)	Substrate
XVEL	Current Velocity (m/s)	Water Flow
PCT_RI	Riffle Percentage (%)	Water Flow
PCT_GL	Glide Percentage (%)	Water Flow
PCT_PL	Pool Percentage (%)	Water Flow
DIV_FLOW	Flow Diversity	Water Flow
CDI	Catchment Disturbance Index	Disturbance
LDI	Local Disturbance Index	Disturbance

individuals of this species were found in sites with hard substrates. Substrate composition is closely linked to the quality and quantity of resources available for benthic species, such as freshwater gastropods,

because hard sediments are substrates for periphytic algae, a major gastropod food source (Aguilar et al., 2015; Linares et al., 2018). As an invasive species, *M. tuberculata* is especially abundant on anthropogenically altered hard substrates (Fernandez et al., 2003; Santos and Eskinazi-Sant’Anna, 2010; Weir and Salice, 2012), such as dams. However, it is less dense in the natural, sandy substrates that characterize most Neotropical Savanna streams (Linares et al., 2019; Miyahira et al., 2020).

Biomphalaria differed from the other Gastropoda taxa, by being the only taxon that was positively correlated with total dissolved solids, a clear link to anthropogenic disturbances. Streams with high levels of total dissolved solids are associated with anthropogenic disturbances, at both local and regional spatial extents (Firmiano et al., 2021). Spatial distributions of many freshwater invertebrates are associated with the level of total dissolved solids in streams, presumably caused by its taxon-specific effects on the maintenance of ion and water balance, which is energetically expensive and thus could affect fitness (Olson and Hawkins, 2017). The positive relation between *Biomphalaria* and total dissolved solids suggests a preference for streams with lower water quality and its resistance to anthropogenic alteration (Firmiano et al., 2021; Mangadze et al., 2019). This genus is often found in wetlands, temporary lakes and other habitats characterized by high amounts of suspended particulate organic matter. Therefore, it is likely that it can successfully colonize streams with similar conditions (Kotzian and Amaral, 2013; Weir and Salice, 2012). The positive relation with both coarse and fine sediments is probably derived from different species in the genus having different habitat preferences, a result supported by previous studies (Estrada et al., 2006; Woke and Aleye-Wokoma, 2015).

Our results showed that *Physa* preferred clear, alkaline streams. As stated before, high levels of dissolved solids often have adverse physiological effects (Olson & Hawkins, 2017). Because of their thin, fragile shells, *Physa* individuals would benefit from alkaline waters (Brown

Table 2

Cumulative AICc weights (w_+) and estimates of variable effects (β parameters) for predictor variables used to model the presence probability of freshwater gastropods taxa in Neotropical Savanna streams. Values of w_+ in bold are those considered to be more likely ($w_+ \geq 0.50$). Estimates of variable effects are based on the most parsimonious model that included that variable and are given only for variables with $w_+ \geq 0.50$.

Variables	<i>Littoridina</i>		<i>Melanoides tuberculata</i>		<i>Biomphalaria</i>		<i>Gundlachia</i>		<i>Physa</i>	
	w_+	β	w_+	β	w_+	β	w_+	β	w_+	β
Basin_elev_range	0.07	–	0.07	–	0.03	–	0.76	1.974	0.18	–
Basin_slope_range	0.04	–	0.09	–	0.05	–	0.05	–	0.02	–
Turbidity	0.05	–	0.03	–	0.06	–	0.06	–	0.04	–
Conductivity	0.06	–	0.16	–	0.03	–	0.10	–	0.08	–
TDS	0.08	–	0.17	–	0.66	9.57	0.22	–	0.69	–436.82
Alcalinity	0.07	–	0.09	–	0.04	–	0.17	–	0.67	655.04
XBKF_D	0.12	–	0.19	–	0.03	–	0.13	–	0.03	–
XBFWRAT	0.43	–	0.24	–	0.05	–	0.11	–	0.03	–
XWD_RAT_P	0.24	–	0.18	–	0.04	–	0.09	–	0.11	–
XEMBED	0.09	–	0.38	–	0.16	–	0.05	–	0.03	–
XCDENMID	0.07	–	0.04	–	0.04	–	0.05	–	0.03	–
XSLOPE	0.91	–246.16	0.05	–	0.03	–	0.09	–	0.17	–
PCT_BL	0.06	–	0.02	–	0.05	–	0.05	–	0.22	–
PCT_CB	0.11	–	0.09	–	0.57	14.76	0.09	–	0.18	–
PCT_GC	0.05	–	0.62	409.91	0.03	–	0.06	–	0.09	–
PCT_GF	0.26	–	0.03	–	0.04	–	0.09	–	0.12	–
PCT_SA	0.13	–	0.03	–	0.20	–	0.08	–	0.27	–
PCT_FN	0.12	–	0.24	–	0.70	22.36	0.09	–	0.05	–
PCT_ORG	0.15	–	0.08	–	0.60	10.73	0.06	–	0.17	–
PCT_WD	0.14	–	0.06	–	0.03	–	0.06	–	0.07	–
XVEL	0.07	–	0.39	–	0.04	–	0.09	–	0.04	–
PCT_RI	0.08	–	0.09	–	0.09	–	0.05	–	0.06	–
PCT_GL	0.09	–	0.07	–	0.03	–	0.42	–	0.06	–
PCT_PD	0.06	–	0.06	–	0.17	–	0.56	2.36	0.15	–
DIV_FLUXO	0.19	–	0.35	–	0.05	–	0.12	–	0.06	–
CDI	0.07	–	0.05	–	0.04	–	0.06	–	0.16	–
LDI	0.06	–	0.04	–	0.03	–	0.05	–	0.14	–

et al., 1998), indicating a preference for streams in karstic regions (Habdija et al., 1995). It also denotes a preference for streams with good water quality, suggesting that this genus might be successfully used as a bioindicator (de Kock and Wolmarans, 2007; Wadaan, 2007).

An important caveat to this study is that our taxonomic resolution was mostly limited to genus-level identifications. Although freshwater gastropods of the same genus may be expected to live in the same habitats and have very similar environmental requirements (Narr and Krist, 2019), more diverse genera can have species with markedly different preferences, as highlighted by our *Biomphalaria* results. Our results provide an important baseline, highlighting the environmental preferences of an often-neglected group. However, we recommended that future studies incorporate less damaging field fixation and working closely with taxonomists to obtain proper species inventories.

5. Conclusions

The freshwater gastropods that we collected had differing ecological requirements, but with an underlying tendency for preferring more stable lotic ecosystems. These results highlight the challenge of protecting freshwater gastropod assemblages because their preferred environments are easily disrupted by common anthropogenic stressors.

Authors contributions

Marden S. Linares: Conceptualization, Methodology, Writing – original draft. **Diego R. Macedo:** Methodology, Writing – review & editing. **Rodrigo L. Massara:** Methodology, Writing – review & editing. **Marcos Callisto:** Writing – review & editing.

Conflict of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This research has been continually funded by Programa Peixe Vivo/ Companhia Energética de Minas Gerais, Programa de Pesquisa e Desenvolvimento Tecnológico do Setor de Energia Elétrica-Companhia Energética de Minas Gerais (P&D Aneel-Cemig GT-487, GT-550, GT-599, and GT-611) and by Fundação de Amparo à Pesquisa de Minas Gerais (APQ-01961-15). This study was financed in part by CAPES - Finance Code 001. MC was awarded National Council for Scientific & Technological Development (CNPq) research productivity grants (304060/2020-8) and by Fundação de Amparo à Pesquisa do Estado de Minas Gerais research grant (PPM 00104-18). DRM received support from CNPq (309763/2020-7). The Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) provided grants to RLM. The team of undergraduate and graduate students of the Laboratório de Ecologia de Bentos/ ICB- UFMG supported the field activities.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.watbs.2022.100076>.

References

- Aguiar, A.C.F., Gücker, B., Brauns, M., Hille, S., Boëchat, I.G., 2015. Benthic invertebrate density, biomass, and instantaneous secondary production along a fifth-order human-impacted tropical river. *Environ. Sci. Pollut. Res.* 22, 9864–9876. <https://doi.org/10.1007/s11356-015-4170-y>.
- APHA, 1998. *Standard Methods for the Examination of Water and Wastewater*, twentieth ed. American Public Health Association, American Water Works Association, Water Environmental Federation, Washington.
- Bae, M.-J., Park, Y.-S., 2020. Key determinants of freshwater gastropod diversity and distribution: the implications for conservation and management. *Water* 12, 1908. <https://doi.org/10.3390/w12071908>.
- Bartón, K., 2019. Package ‘MuMIn’. <https://cran.r-project.org/web/packages/MuMIn/index.html>.
- Berger, J.O., Barbieri, M.M., 2004. Optimal predictive model selection. *Ann. Stat.* 32, 870–897. [s11749-018-0599-x](https://doi.org/10.1214/018-0599-x).

- Bivand, R., Wong, D.W.S., 2018. Comparing implementations of global and local indicators of spatial association. *Test* 27 (3), 716–748 <https://doi.org/10.1007/>
- Böhm, M., Dewhurst-Richman, N.I., Seddon, M., Ledger, S.E.H., Albrecht, C., Allen, D., Bogan, A.E., Cordeiro, J., Cummings, K.S., Cuttled, A., Darrigran, G., Darwall, W., Fehér, Z., Gibson, C., Graf, D.L., Köhler, F., Lopes-Lima, M., Pastorino, G., Perez, K.E., Smith, K., van Damme, D., Vinarski, M.V., von Proschwitz, T., von Rintelen, T., Aldridge, D.C., Aravind, N.A., Budha, P.B., Clavijo, C., Van Tu, D., Gargominy, O., Ghamizi, M., Haase, M., Hilton-Taylor, C., Johnson, P.D., Kebapçı, Ü., Lajtner, J., Lange, C.N., Lepitzki, D.A.W., Martínez-Ortí, A., Moorkens, E.A., Neubert, E., Pollock, C.M., Prié, V., Radea, C., Ramirez, R., Ramos, M.A., Santos, S.B., Slapnik, R., Son, M.O., Stensgaard, A.S., Collen, B., 2021. The conservation status of the world's freshwater molluscs. *Hydrobiologia* 848, 3231–3254. <https://doi.org/10.1007/s10750-020-04385-w>.
- Böhmer, J., Rawer-Jost, C., Zenker, A., Meier, C., Feld, C.K., Biss, R., Hering, D., 2004. Assessing streams in Germany with benthic invertebrates: development of a multimetric invertebrate based assessment system. *Limnologica* 34, 416–432. [https://doi.org/10.1016/S0075-9511\(04\)80010-0](https://doi.org/10.1016/S0075-9511(04)80010-0).
- Brown, K.M., Alexander, J.E., Thorp, J.H., 1998. Differences in the ecology and distribution of lotic pulmonate and prosobranch gastropods. *Am. Malacol. Bull.* 14 (2), 91–101.
- Burnham, K.P., Anderson, D.R., 2002. *Model Selection and Multimodel Inference: A Practical Information-Theoretical Approach*. Springer-Verlag, New York, New York.
- CEMIG - Companhia Energética de Minas Gerais, 2014. *Condições Ecológicas em bacias hidrográficas de empreendimentos hidrelétricos – integridade Biótica. Série Peixe Vivo – Transposição de Peixes v1 (1ed)*, 1–216.
- Chattopadhyay, S., Oglecki, P., Keller, A., Kardel, I., Mirosław-Świątek, D., Mikołaj Piniewski, M., 2021. Effect of a Summer Flood on Benthic Macroinvertebrates in a Medium-Sized, Temperate, Lowland River. *Water*. <https://doi.org/10.3390/w13070885>.
- Collen, B., Whitton, F., Dyer, E.E., Baillie, J.E.M., Cumberlidge, N., Darwall, W.R.T., Pollock, C., Richman, N.I., Soulsby, A.-M., Böhm, M., 2014. Global patterns of freshwater species diversity, threat and endemism. *Global Ecol. Biogeogr.* 23, 40–51. <https://doi.org/10.1111/geb.12096>.
- CoreTeam, R., 2015. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.Rproject.org>.
- Darwall, W., Bremerich, V., De Wever, A., Dell, A.I., Freyhof, J., Gessner, M.O., Grossart, H.P., Harrison, I., Irvine, K., Jähmig, S.C., Jeschke, J.M., Lee, J.J., Lu, C., Lewandowska, A.M., Monaghan, M.T., Nejtgaard, J.C., Patricio, H., Schmidt-Kloiber, A., Stuart, S.N., Thieme, M., Tockner, K., Turak, E., Weyl, O., 2018. The Alliance for Freshwater Life: a global call to unite efforts for freshwater biodiversity science and conservation. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 28, 1015–1022. <https://doi.org/10.1002/aqc.2958>.
- Datry, T., Larned, S.T., Tockner, K., 2014. Intermittent rivers: a challenge for freshwater ecology. *Bioscience* 64 (3), 229–235. <https://doi.org/10.1093/biosci/bit027>.
- de Kock, K., Wolmarans, C.T., 2007. Distribution and Habitats of the Alien Invader Freshwater Snail *Physa Acuta* in South Africa, vol. 33. *Water SA*. <https://doi.org/10.4314/wsa.v33i5.184093>.
- de Vries, J., Kraak, M.H.S., Verdonshot, R.C.M., Verdonshot, P.F.M., 2019. Quantifying cumulative stress acting on macroinvertebrate assemblages in lowland streams. *Sci. Total Environ.* 694, 133630. <https://doi.org/10.1016/j.scitotenv.2019.133630>.
- Doherty, P.F., White, G.C., Burnham, K.P., 2012. Comparison of model building and selection strategies. *J. Ornithol.* 152, 317–323. <https://doi.org/10.1007/s10336-010-0598-5>.
- Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B., Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E., Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36, 27–46.
- Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.-I., Knowler, D.J., Lévêque, C., Naiman, R.J., Prieur-Richard, A.-H., Soto, D., Stiassny, M.L.J., Sullivan, C.A., 2006. Freshwater biodiversity: importance, threats, status and conservation challenges. *Biol. Rev.* 81, 163. <https://doi.org/10.1017/S1464793105006950>.
- Erba, S., Cazzola, M., Belfiore, C., Buffagni, A., 2020. Macroinvertebrate metrics responses to morphological alteration in Italian rivers. *Hydrobiologia* 847 (9), 2169–2191. <https://doi.org/10.1007/s10750-020-04242-w>.
- Esselman, P.C., Infante, D.M., Wang, L., Cooper, A.R., Wiefelich, D., Tsang, Y.-P., Thornbrugh, D.J., Taylor, W.W., 2013. Regional fish community indicators of landscape disturbance to catchments of the conterminous United States. *Ecol. Indic.* 26, 163–173. <https://doi.org/10.1016/j.ecolind.2012.10.028>.
- Esteves, F.A., Amorim, J.C., Cardoso, E.L., Barbosa, F.A.R., 1985. Caracterização limnológica preliminar da represa de Três Marias (MG) com base em alguns parâmetros ambientais básicos. *Cienc. Cult. (Sao Paulo)* 37, 608–617.
- Estrada, V.E., Velásquez, L.E., Caldeira, R.L., Bejarano, E.E., Rojas, W., Carvalho, O.S., 2006. Phylogenetics of South American Biomphalaria and description of a new species (Gastropoda: Planorbidae). *J. Molluscan Stud.* 72 (3), 221–228. <https://doi.org/10.1093/mollus/eyl003>.
- Fernandez, M.A., Thiengo, S.C., Simone, L.R.L., 2003. Distribution of the Introduced Freshwater Snail *Melanooides turbeculatus* (Gastropoda: Thiaridae) in Brazil. *Nautilus, Philadelphia*.
- Firmiano, K.R., Castro, D.M.P., Linares, M.S., Callisto, M., 2021. Functional responses of aquatic invertebrates to anthropogenic stressors in riparian zones of Neotropical savanna streams. *Sci. Total Environ.* 753, 141865. <https://doi.org/10.1016/j.scitotenv.2020.141865>.
- Geiger, D.L., Marshall, B.A., Ponder, W.F., Sasaki, T., Warén, A., 2007. Techniques for collecting, handling, preparing, storing and examining small molluscan specimens. *Molluscan Res.* 27 (1), 1–50.
- Graf, W., Leitner, P., Hanetseder, I., Ittner, L.D., Dossi, F., Hauer, C., 2016. Ecological degradation of a meandering river by local channelization effects: a case study in an Austrian lowland river. *Hydrobiologia* 772, 145–160. <https://doi.org/10.1007/s10750-016-2653-6>.
- Habdija, I., Lajtner, J., Belinić, I., 1995. The Contribution of Gastropod Biomass in Macroinvertebrate Communities of a Karstic River, vol. 80, pp. 103–110. <https://doi.org/10.1002/iroh.19950800113>, 1.
- Hartig, F., 2022. Package "DHARMA: Residual Diagnostics for Hierarchical (Multi-Level/Mixed) Regression Models". <http://florianhartig.github.io/DHARMA/>.
- Horsák, M., Bojková, J., Zahrádková, S., Omesová, M., Helešic, J., 2009. Impact of reservoirs and channelization on lowland river macroinvertebrates: a case study from Central Europe. *Limnol. - Ecol. Manag. Int. Waters* 39, 140–151. <https://doi.org/10.1016/j.limno.2008.03.004>.
- Johnson, P.D., Bogan, A.E., Brown, K.M., Burkhead, N.M., Cordeiro, J.R., Garner, J.T., Hartfield, P.D., Lepitzki, D.A.W., Mackie, G.L., Pip, E., Tarpley, T.A., Tiemann, J.S., Whelan, N.V., Strong, E.E., 2013. Conservation status of freshwater gastropods of Canada and the United States. *Fisheries* 38, 247–282. <https://doi.org/10.1080/03632415.2013.785396>.
- Jones, C.G., Lawton, J.H., Shachak, M., 1994. Organisms as ecosystem engineers. *Oikos* 69, 373–386. <https://doi.org/10.2307/3545850>.
- Jörger, K.M., Stöger, I., Kano, Y., Fukuda, H., Knebelberger, T., Schrödl, M., 2010. On the origin of Acochlidia and other enigmatic euthyneuran gastropods, with implications for the systematics of Heterobranchia. *BMC Evol. Biol.* 10, 323. <https://doi.org/10.1186/1471-2148-10-323>.
- Karatayev, A.Y., Padilla, D.K., Minchin, D., Boltovskoy, D., Burlakova, L.E., 2007. Changes in global economies and trade: the potential spread of exotic freshwater bivalves. *Biol. Invasions* 9, 161–180. <https://doi.org/10.1007/s10530-006-9013-9>.
- Kaufmann, P.R.P., Levine, P., Robison, E.E.G., Seeliger, C., Peck, D.V.D., 1999. *Quantifying Physical Habitat in Wadeable Streams*. U.S. Environmental Protection Agency, Washington, DC. EPA/620/R-99/003.
- Kaufmann, P.R., Hughes, R.M., Paulsen, S.G., Peck, D.V., Seeliger, C.W., Weber, M., Mitchell, R.M., 2022. Physical habitat in conterminous US streams and rivers, part 1: geomorphic controls and anthropogenic alteration. *Ecol. Indic.* <https://doi.org/10.1016/j.ecolind.2022.109046>.
- Kotzian, C.B., Amaral, A.M.B., 2013. Diversity and distribution of mollusks along the Contas River in a tropical semiarid region (Caatinga), Northeastern Brazil. *Biota Neotropica* 13, 299–314.
- Lecocq, T., Harpke, A., Rasmont, P., Schweiger, O., 2019. Integrating intraspecific differentiation in species distribution models: consequences on projections of current and future climatically suitable areas of species. *Divers. Distrib.* 25 (7), 1–13. <https://doi.org/10.1111/ddi.12916>.
- Ligeiro, R., Hughes, R.M., Kaufmann, P.R., Macedo, D.R., Firmiano, K.R., Ferreira, W.R., Oliveira, D., Melo, A.S., Callisto, M., 2013. Defining quantitative stream disturbance gradients and the additive role of habitat variation to explain macroinvertebrate taxa richness. *Ecol. Indic.* 25, 45–57. <https://doi.org/10.1016/j.ecolind.2012.09.004>.
- Linares, M.S., Callisto, M., Marques, J.C., 2018. Thermodynamic based indicators illustrate how a run-of-river impoundment in neotropical savanna attracts invasive species and alters the benthic macroinvertebrate assemblages' complexity. *Ecol. Indic.* 88, 181–189. <https://doi.org/10.1016/j.ecolind.2018.01.040>.
- Linares, M.S., Assis, W., Castro Solar, R.R., Leitão, R.P., Hughes, R.M., Callisto, M., 2019. Small hydropower dam alters the taxonomic composition of benthic macroinvertebrate assemblages in a neotropical river. *River Res. Appl.* rra. 3442. <https://doi.org/10.1002/rra.3442>.
- Linares, M.S., Macedo, D.R., Massara, R.L., Callisto, M., 2020. Why are they here? Local variables explain the distribution of invasive mollusk species in neotropical hydropower reservoirs. *Ecol. Indic.* 117, 106674. <https://doi.org/10.1016/j.ecolind.2020.106674>.
- Lopes-Lima, M., Riccardi, N., Urbanska, M., Köhler, F., Vinarski, M., Bogan, A.E., Sousa, R., 2021. Major Shortfalls Impairing Knowledge and Conservation of Freshwater Molluscs. *Hydrobiologia*. <https://doi.org/10.1007/s10750-021-04622-w>.
- Macedo, D.R., Hughes, R.M., Ligeiro, R., Ferreira, W.R., Castro, M.A., Junqueira, N.T., Oliveira, D.R., Firmiano, K.R., Kaufmann, P.R., Pompeu, P.S., Callisto, M., 2014a. The relative influence of catchment and site variables on fish and macroinvertebrate richness in cerrado biome streams. *Landsc. Ecol.* 29, 1001–1016. <https://doi.org/10.1007/s10980-014-0036-9>.
- Macedo, D.R., Pompeu, P.S., MORAIS, L., Castro, M.A., Alves, C.B.M., França, J.S., Sanches, B.O., Agra, J.U.M., Callisto, M., 2014b. Sampling site selection, land use and cover, field reconnaissance, and sampling. In: Callisto, M., Hughes, R.M., Lopes, J.M., Castro, M.A. (Eds.), *Ecological Conditions in Hydropower Basins. Série Peixe Vivo 3*. Companhia Energética de Minas Gerais. Belo Horizonte, Brazil, pp. 61–83.
- Macedo, D.R., Hughes, R.M., Ferreira, W.R., Firmiano, K.R., Silva, D.R.O., Ligeiro, R., Kaufmann, P.R., Callisto, M., 2016. Development of a benthic macroinvertebrate multimetric index (MMI) for Neotropical Savanna headwater streams. *Ecol. Indic.* 64, 132–141. <https://doi.org/10.1016/j.ecolind.2015.12.019>.
- Mangadze, T., Wasserman, R.J., Froneman, P.W., Dalu, T., 2019. Macroinvertebrate functional feeding group alterations in response to habitat degradation of headwater Austral streams. *Sci. Total Environ.* 695, 133910. <https://doi.org/10.1016/j.scitotenv.2019.133910>.
- Meier-Brook, C., 1976. The influence of vared relaxing and fixing conditions on anatomical characters in a *Planorbis* species. *Basteria* 40, 101–106.
- Meyer, J.L., Strayer, D.L., Wallace, J.B., Eggert, S.L., Helfman, G.S., Leonard, N.E., 2007. The contribution of headwater streams to biodiversity in river networks. *J. Am. Water Resour. Assoc.* 43 (1), 86–103. <https://doi.org/10.1111/j.1752-1688.2007.00008.x>.

- Miyahira, I., Pereira, L., dos Santos, L., 2020. Non-native freshwater molluscs in the Neotropics: what can be learned from Brazilian reservoirs? *Aquat. Invasions* 15, 455–472. <https://doi.org/10.3391/ai.2020.15.3.06>.
- Mugnai, R., Nessimian, J.L., Baptista, D.F., 2010. *Manual de Identificação de Macroinvertebrados Aquáticos do Estado do Rio de Janeiro*. Technical Books Editora, Rio de Janeiro.
- Narr, C.F., Krist, A.C., 2019. Improving estimates of richness, habitat associations, and assemblage characteristics of freshwater gastropods. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 30, 131–143. <https://doi.org/10.1002/aqc.3247>, 2020.
- Oliveira, M.D., Calheiros, D.F., Jacobi, C.M., Hamilton, S.K., 2011. Abiotic factors controlling the establishment and abundance of the invasive golden mussel *Limnoperna fortunei*. *Biol. Invasions* 13, 717–729. <https://doi.org/10.1007/s10530-010-9862-0>.
- Olsen, A.R., Peck, D.V., 2008. Survey design and extent estimates for the Wadeable streams assessment. *J. North Am. Benthol. Soc.* 27, 822–836. <https://doi.org/10.1899/08-050.1>.
- Olson, J.R., Hawkins, C.P., 2017. Effects of total dissolved solids on growth and mortality predict distributions of stream macroinvertebrates. *Freshwater Biology* 62 (4), 779–791. <https://doi.org/10.1111/fwb.12901>.
- Paraense, L., 1975. Estudo atual da sistemática dos planorbídeos brasileiros. *Arquivos Do Museu Nacional Do Rio de Janeiro* 55 (i).
- Peck, D., Herlihy, A., Hill, B., Hughes, R., Kaufmann, P., Klemm, D., Lazorchak, J., McCormick, F., Peterson, S., Ringold, P., Magee, T., Cappaert, M., 2006. *Environmental Monitoring and Assessment Program-Surface Waters Western Pilot Study: Field Operations Manual for Wadeable Streams*. U.S. Environmental Protection Agency, Washington, DC. EPA/620/R-06/003.
- Rawer-Jost, C., Zenker, A., Böhmer, J., 2004. Reference conditions of German stream types analysed and revised with macroinvertebrate fauna. *Limnologia* 34, 390–397. [https://doi.org/10.1016/S0075-9511\(04\)80008-2](https://doi.org/10.1016/S0075-9511(04)80008-2).
- Reid, A.J., Carlson, A.K., Creed, I.F., Eliason, E.J., Gell, P.A., Johnson, P.T.J., Kidd, K.A., McCormack, T.J., Olden, J.D., Ormerod, S.J., Smol, J.P., Taylor, W.W., Tockner, K., Vermaire, J.C., Dudgeon, D., Cooke, S.J., 2019. Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biol. Rev.* 94, 849–873. <https://doi.org/10.1111/bvr.12480>.
- Santos, C., Eskinazi-Sant'Anna, E., 2010. The introduced snail *Melanoides tuberculatus* (Muller, 1774) (Mollusca: Thiaridae) in aquatic ecosystems of the Brazilian semiarid Northeast (Piranhas-Assu river basin, state of Rio grande do Norte). *Braz. J. Biol.* 70, 1–7. <https://doi.org/10.1590/S1519-69842010000100003>.
- Shandas, V., Alberti, M., 2009. Exploring the role of vegetation fragmentation on aquatic conditions: linking upland with riparian areas in Puget Sound lowland streams. *Landsch. Urban Plann.* 90, 66–75. <https://doi.org/10.1016/j.landurbplan.2008.10.016>.
- Silva, D.R.O., Herlihy, A.T., Hughes, R.M., Callisto, M., 2017. An improved macroinvertebrate multimetric index for the assessment of Wadeable streams in the neotropical savanna. *Ecol. Indic.* 81, 514–525. <https://doi.org/10.1016/j.ecolind.2017.06.017>.
- Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E., Vilà, M., 2013. Impacts of biological invasions: what's what and the way forward. *Trends Ecol. Evol.* 28, 58–66. <https://doi.org/10.1016/j.tree.2012.07.013>.
- Simone, L.R.L., 2006. *Land and Freshwater Molluscs of Brazil*. EGB, Fapesp, São Paulo, 400 390.
- Smeraldo, S., Bosso, L., Fraissinet, M., Bordignon, L., Brunelli, M., Ancillotto, L., Russo, D., 2020. Modelling risks posed by wind turbines and power lines to soaring birds: the black stork (*Ciconia nigra*) in Italy as a case study. *Biodivers. Conserv.* 29 (3), 1959–1976. <https://doi.org/10.1007/s10531>.
- Strahler, A.N., 1957. Quantitative analysis of watershed geomorphology. *New Haven: Transactions: American Geophysical Union* 38 (6), 913–920. <https://doi.org/10.1029/TR038i006p00913>.
- Strong, E.E., Gargominy, O., Ponder, W.F., Bouchet, P., 2008. Global diversity of gastropods (Gastropoda; Mollusca) in freshwater. *Hydrobiologia* 595, 149–166. <https://doi.org/10.1007/s10750-007-9012-6>.
- Turak, E., Harrison, I., Dudgeon, D., Abell, R., Bush, A., Darwall, W., Finlayson, C.M., Ferrier, S., Freyhof, J., Hermoso, V., Juffe-Bignoli, D., Linke, S., Nel, J., Patricio, H.C., Pittock, J., Raghavan, R., Revenga, C., Simaika, J.P., De Wever, A., 2016. Essential Biodiversity Variables for measuring change in global freshwater biodiversity. *Biol. Conserv.* 213 (B), 272–279. <https://doi.org/10.1016/j.biocon.2016.09.005>.
- Usseglio-Polatera, P., Beisel, J.N., 2002. Longitudinal changes in macroinvertebrate assemblages in the Meuse River: anthropogenic effects versus natural change. *River Res. Appl.* 18, 197–211. <https://doi.org/10.1002/rra.651>.
- van Rees, C.B., Waylen, K.A., Schmidt-Kloiber, A., Thackeray, S.J., Kalinkat, G., Martens, K., Domisch, S., Lillebø, A.I., Hermoso, V., Grossart, H., Schinegger, R., Declerck, K., Adriaens, T., Denys, L., Jarić, I., Janse, J.H., Monaghan, M.T., De Wever, A., Geijzendorffer, I., Adamescu, M.C., Jähnig, S.C., 2021. Safeguarding freshwater life beyond 2020: Recommendations for the new global biodiversity framework from the European experience. *Conserv. Lett.* 14, 1–17. <https://doi.org/10.1111/cons.12771>.
- Wadaan, M., 2007. The fresh water growing snail *Physa acuta* : a suitable bioindicator for testing cadmium Toxicity. *Saudi J. Biol. Sci.* 14.
- Weir, S.M., Salice, C.J., 2012. High tolerance to abiotic stressors and invasion success of the slow growing freshwater snail, *Melanoides tuberculatus*. *Biol. Invasions* 14, 385–394. <https://doi.org/10.1007/s10530-011-0084-x>.
- Woke, G.N., Aleleye-Wokoma, I.P., 2015. Effects of sediment input on aquatic animal communities in new Calabar river, Rivers state, Nigeria. *Global Journal of Applied Sciences* 21, 29–32. <https://doi.org/10.4314/gjas.v21i1.5>.