

Colouring the Normalized Laplacian

Gabriel Coutinho¹

*Dept. Computer Science
Federal University of Minas Gerais
Belo Horizonte, Brazil*

Rafael Grandsire²

*Computer Science major
Federal University of Minas Gerais
Belo Horizonte, Brazil*

Célio Passos³

*Computational Mathematics major
Federal University of Minas Gerais
Belo Horizonte, Brazil*

Abstract

We apply Cauchy's interlacing theorem to derive some eigenvalue bounds to the chromatic number using the normalized Laplacian matrix, including a combinatorial characterization of when equality occurs. Further, we introduce some new expansion type of parameters which generalize the Cheeger constant of a graph, and relate them to the colourings which meet our eigenvalue bound with equality. Finally, we point out a connection to the Erdős-Faber-Lovász conjecture.

Keywords: Normalized Laplacian, interlacing, eigenvalue bound, chromatic number, Cheeger constant.

1 Introduction

Let G be a graph without isolated vertices, \mathbf{A} its adjacency matrix, and \mathbf{D} the diagonal matrix that records the degrees of the vertices. The matrix

$$\mathbf{L} = \mathbf{I} - \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}$$

¹ Email: gabriel@dcc.ufmg.br.

² This author acknowledges the support of the undergrad research scholarship PRPq-UFMG from ADRC (G. Coutinho) - 11/2017.

³ Research of this author was done while he was an undergrad researcher attached to the ICV-UFMG program.

is known as the normalized Laplacian matrix of G . The standard reference for an account of the interplay between combinatorics and the spectral properties of \mathbf{L} is Fan Chung's book [6]. In this paper, we are interested in which properties of \mathbf{L} can give information about graph colourings. It is unlikely that information contained solely in the eigenvalues of \mathbf{L} could determine $\chi(G)$. However one can find some eigenvalue bounds to $\chi(G)$. An eigenvalue bound is considered relatively serious if it achieves equality for at least one non-trivial infinite class of graphs, and, moreover, if the equality implies some combinatorial structure. We present a strictly speaking new bound in Theorem 2.2. Across the paper, we study some consequences of the equality cases in this bound, and in the last section we comment on a possible interesting class of examples connected to the Erdős-Faber-Lovász conjecture. We also introduce in Section 4 new expansion-type of parameters, and compare some of their properties to the spectrum of \mathbf{L} .

2 Interlacing bound for the normalized Laplacian

We begin by re-stating the famous interlacing theorem due to Cauchy (see for instance [9, Theorem 2.1]).

Theorem 2.1 *Let \mathbf{P} be a real $n \times m$ matrix such that $\mathbf{P}^T \mathbf{P} = \mathbf{I}$ and let \mathbf{M} be a symmetric $n \times n$ matrix with eigenvalues $\theta_1 \geq \dots \geq \theta_n$. Define $\mathbf{N} = \mathbf{P}^T \mathbf{M} \mathbf{P}$ and let \mathbf{N} have eigenvalues $\mu_1 \geq \dots \geq \mu_m$ and respective eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_m$. Then the following hold.*

(i) *The eigenvalues of \mathbf{N} interlace those of \mathbf{M} , meaning that, for $i = 1, \dots, m$, we have*

$$\theta_{n-m+i} \leq \mu_i \leq \theta_i.$$

(ii) *If $\mu_i = \theta_i$ or $\mu_i = \theta_{n-m+i}$ for some $i \in [1, m]$, then \mathbf{N} has a μ_i -eigenvector \mathbf{v} so that $\mathbf{P}\mathbf{v}$ is a μ_i -eigenvector of \mathbf{M} .*

(iii) *If, for some integer ℓ , $\mu_i = \theta_i$ for $i = 1, \dots, \ell$ (or $\mu_i = \theta_{n-m+i}$ for $i = \ell, \dots, m$), then $\mathbf{P}\mathbf{v}_i$ is a μ_i -eigenvector for \mathbf{M} for $i = 1, \dots, \ell$ (respectively $i = \ell, \dots, m$).*

(iv) *If the interlacing is tight, that is, if there is an integer $k \in [0, m]$ so that*

$$\theta_i = \mu_i \quad \text{for } 1 \leq i \leq k, \quad \text{and} \quad \theta_{n-m+i} = \mu_i \quad \text{for } k+1 \leq i \leq m,$$

then

$$\mathbf{M}\mathbf{P} = \mathbf{P}\mathbf{N}.$$

□

Let G be a graph on n vertices. Let $\mathbf{A}(G) = \mathbf{A}$ be its adjacency matrix, and \mathbf{D} the diagonal matrix that records the degree sequence of G . The matrix $\mathbf{D} - \mathbf{A}$ is usually called the Laplacian matrix of G , but in this paper we are interested in its normalized version

$$\mathbf{L} = \mathbf{I} - \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}.$$

Say G contains a (proper) k -colouring, that is, its vertex set can be partitioned into k independent sets S_1, \dots, S_k . Let π be this partition and \mathbf{S} be its characteristic

matrix, that is, the 01 matrix whose rows are indexed by the vertices, columns by the colour classes, and an entry equal to 1 indicates that the vertex corresponding to the row has been coloured by the class corresponding to the column. By G/π we denote the quotient graph, meaning, the graph whose vertices are the classes of π and between any two vertices there is a weighted edge whose weight is equal to the number of edges between the corresponding classes in G . In particular, G/π has no loops because π is a (proper) colouring, and $\mathbf{S}^T \mathbf{A} \mathbf{S} = \mathbf{A}(G/\pi)$.

We define

$$\mathbf{P} = \mathbf{D}^{1/2} \mathbf{S} \mathbf{N}^{-1/2},$$

where \mathbf{N} is the diagonal matrix whose k th diagonal entry is equal to sum of the degrees of the vertices in S_k . In particular, this turns the columns of \mathbf{P} into normalized vectors.

Thus

$$\mathbf{P}^T \mathbf{P} = \mathbf{I}. \quad (1)$$

Moreover,

$$\mathbf{P}^T \mathbf{L} \mathbf{P} = \mathbf{I} - \mathbf{N}^{-1/2} \mathbf{A}(G/\pi) \mathbf{N}^{-1/2} = \mathbf{L}(G/\pi). \quad (2)$$

Observe that 0 is an eigenvalue of both \mathbf{L} and $\mathbf{L}(G/\pi)$ with respective eigenvectors $\mathbf{D}^{1/2} \mathbf{1}$ and $\mathbf{N}^{1/2} \mathbf{1}$. If $0 = \lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1}$ are the eigenvalues of \mathbf{L} , then we define

$$\sigma_k = \sum_{j=1}^k \lambda_{n-j},$$

that is, the sum of the largest k eigenvalues of \mathbf{L} .

Theorem 2.2 *If $\sigma_{k-1} < k$, then $k < \chi(G)$.*

Proof. This is a consequence of Equations (1) and (2) and Theorem 2.1. In particular, let $k \geq \chi(G)$, and consider a partition π of $V(G)$ into k independent sets. The eigenvalues of $\mathbf{L}(G/\pi)$ interlace those of \mathbf{L} , so

$$k = \text{tr } \mathbf{L}(G/\pi) \leq \sigma_k.$$

We can however ignore the least eigenvalue of $\mathbf{L}(G/\pi)$ which is 0, thus the inequality still holds for σ_{k-1} . \square

Let $\pi = \{S_1, \dots, S_k\}$ be any k -partition of $V(G)$. Then the entries of $\mathbf{A}(G/\pi)$ record the number of edges between the classes of π . We recall the usual notation $e(S, T)$ that counts the number of edges between sets S and T , and $\text{vol}(S)$ which is the sum of the degrees of the vertices in S . Thus

$$\text{tr } \mathbf{L}(G/\pi) = k - \sum_{j=1}^k \frac{e(S_j, S_j)}{\text{vol}(S_j)}.$$

Using interlacing, we reach an interesting lower bound on what any attempt of colouring a graph with k colours can yield.

Corollary 2.3 *For any k -partition $\pi = \{S_1, \dots, S_k\}$ of a graph G , it follows that*

$$\sum_{j=1}^k \frac{e(S_j, S_j)}{\text{vol}(S_j)} \geq k - \sigma_{k-1}.$$

□

We have two remarks.

First, Theorem 2.2 indicates that graphs with small chromatic number should have large normalized Laplacian eigenvalues. For instance, the well-known fact that bipartite graphs have largest eigenvalue equal to 2 follows immediately from the result above.

Second, Theorem 2.2 has been, *mutatis mutandis*, already known for a long time. Interlacing has seen countless applications in graph theory. As far as we know, first versions of this technique to study the chromatic number appeared in Haemers [8], where interlacing for the adjacency matrix was used. In fact, Theorem 2.2 is equivalent to [4, Proposition 3.6.3 (i)] for regular graphs. More recently, Bollobás and Nikiforov obtained similar results for the Laplacian matrix [2,3]. Finally, Butler used interlacing [5, Chapter 5] for the normalized Laplacian and proved almost the same result (Theorem 36) as Corollary 2.3 above. The only, but yet relevant, difference is that we have our results for σ_{k-1} , and not σ_k . This seems a small difference, but note that the results are trivially true if stated with σ_k instead.

Our main goal over the next sections is to try to provide some new useful insights about an old technique.

3 Tight interlacing and equitable colourings

Recall that G is a graph with normalized Laplacian matrix \mathbf{L} , and we denote its eigenvalues by $0 = \lambda_0 \leq \dots \leq \lambda_{n-1}$. We also defined $\sigma_k(G)$ as the sum of the k largest eigenvalues of $\mathbf{L}(G)$, and made a similar definition for G/π replacing G . Assume $\pi = \{S_1, \dots, S_k\}$ is a k -colouring with characteristic matrix \mathbf{S} . Again, \mathbf{D} is the diagonal matrix recording the degrees of G and \mathbf{N} is the diagonal matrix recording the sum of the degrees of the vertices in each S_j . We now turn our attention to the case where interlacing (see Theorem 2.1) is tight.

Lemma 3.1 *Let π be a k -colouring and $\mathbf{P} = \mathbf{D}^{1/2}\mathbf{S}\mathbf{N}^{-1/2}$. If $\sigma_{k-1} = k$, then interlacing of $\mathbf{L} = \mathbf{L}(G)$ according to \mathbf{P} is tight, and in particular, $\mathbf{L}\mathbf{P} = \mathbf{P}\mathbf{L}(G/\pi)$.*

Proof. If $\sigma_{k-1}(G) = k$, then

$$k = \sigma_{k-1}(G) \geq \sigma_{k-1}(G/\pi) = \text{tr } \mathbf{L}(G/\pi) = k.$$

So the inequality holds with equality, and therefore each eigenvalue of $\mathbf{L}(G/\pi)$ must meet the upper bound given by interlacing, except for the last. Hence interlacing is

tight, and from Theorem 2.1 item (iv), the equality $\mathbf{LP} = \mathbf{PL}(G/\pi)$ holds. \square

Consider a partition $\pi = \{S_1, \dots, S_m\}$ of the set $[n]$. Assume this partition has characteristic matrix \mathbf{Q} (which is $n \times m$). Let \mathbf{M} be a symmetric $n \times n$ matrix. The partition π is called *equitable* to \mathbf{M} if

$$\sum_{k \in S_j} \mathbf{M}_{ik}$$

is constant for all i which belongs to the same class of π , for all j , or, equivalently, if the column space of \mathbf{Q} is \mathbf{M} -invariant. If $\mathbf{M} = \mathbf{A}$, then to say that π is equitable to \mathbf{A} is equivalent to saying that the number of neighbours a vertex in class i has in class j depends only on i and j .

The theorem below is an adaptation of the theory of equitable partitions for adjacency matrices [7, Chapter 9] to the normalized Laplacians.

Lemma 3.2 *Let π be a partition of $V(G)$ with characteristic matrix \mathbf{S} , and let $\mathbf{P} = \mathbf{D}^{1/2}\mathbf{S}\mathbf{N}^{-1/2}$. Then π is equitable to $\mathbf{D}^{-1}\mathbf{A}$ if and only if $\mathbf{L}(G)\mathbf{P} = \mathbf{PL}(G/\pi)$.*

Proof. We use $e(S, T)$ to denote the number of edges between sets of vertices S and T . If $S = \{a\}$, we might simply write $e(a, T)$. First we shall note that $\mathbf{L}(G)\mathbf{P} = \mathbf{PL}(G/\pi)$ is equivalent to

$$\mathbf{D}^{-1}\mathbf{A}(G)\mathbf{S} = \mathbf{S}\mathbf{N}^{-1}\mathbf{A}(G/\pi).$$

If this equality holds, and if a is a vertex that belongs to class S_i , then

$$\frac{e(a, S_j)}{d(a)} = (\mathbf{D}^{-1}\mathbf{A}(G)\mathbf{S})_{aj} = (\mathbf{S}\mathbf{N}^{-1}\mathbf{A}(G/\pi))_{aj} = \frac{e(S_i, S_j)}{\text{vol}(S_i)}.$$

So the partition is equitable. On the other hand, if the partition is equitable, then fix S_i , say $S_i = \{a_1, \dots, a_f\}$, and for all S_j , it follows that for all $a_k, a_\ell \in S_i$, we have

$$\frac{e(a_k, S_j)}{d(a_k)} = \frac{e(a_\ell, S_j)}{d(a_\ell)}.$$

So there are constants A, B and q_1, \dots, q_f such that $e(a_k, S_j) = Aq_k$ and $d(a_k) = Bq_k$. Then it is immediate to check that for all $a \in S_i$,

$$\frac{e(S_i, S_j)}{\text{vol}(S_i)} = \frac{A \sum q_k}{B \sum q_k} = \frac{A}{B} = \frac{e(a, S_j)}{d(a)},$$

thus $\mathbf{L}(G)\mathbf{P} = \mathbf{PL}(G/\pi)$. \square

Using Theorem 1 from [10], the following lemma is straightforward. We provide a proof that follows immediately from Theorem 2.2.

Lemma 3.3 *Let $\lambda = \lambda_{n-1}$ be the largest eigenvalue of the normalized Laplacian matrix \mathbf{L} of a graph G . Then*

$$\chi \geq 1 + \frac{1}{\lambda - 1}.$$

Proof. If $k = \chi$, then $\chi \leq \sigma_{\chi-1}(G) \leq (\chi - 1)\lambda$, which implies the result. \square

We can now provide a significant consequence for the case where equality in the lemma above holds.

Theorem 3.4 *Let G be a graph which admits a colouring with k colours, and let $\lambda = \lambda_{n-1}$ be its largest normalized Laplacian eigenvalue. If*

$$\lambda = \frac{k}{k-1},$$

then this colouring is equitable with respect to $\mathbf{D}^{-1}\mathbf{A}$.

Proof. If π is a k -colouring, we have $k = (k-1)\lambda \geq \sigma_{k-1}(G) \geq \sigma_{k-1}(G/\pi) = k$.

In this case, $\sigma_{k-1}(G) = k$, and by Lemma 3.1 we have a tight interlacing according to π , resulting in the equality $\mathbf{L}\mathbf{P} = \mathbf{P}\mathbf{L}(G/\pi)$. Therefore, by Lemma 3.2, π must be equitable with respect to $\mathbf{D}^{-1}\mathbf{A}$. \square

4 Regular colourings

We say that a colouring of a graph G is *regular* if between any two colour classes there is the same number of edges. For a partition $\pi = \{S_1, \dots, S_m\}$ of $V(G)$, define

$$\begin{aligned} \text{(i)} \quad \gamma(\pi) &= \frac{\min_{i \neq j} e(S_i, S_j)}{\max_i \text{vol } S_i}, \\ \text{(ii)} \quad \gamma^*(\pi) &= \frac{\max_{i \neq j} e(S_i, S_j)}{\min_i \text{vol } S_i}. \end{aligned}$$

Moreover, for reasons that will become clear quite soon, define

$$\begin{aligned} \text{(iii)} \quad \psi_k(G) &= \max_{\pi, k\text{-partition}} \gamma(\pi), \\ \text{(iv)} \quad \phi_k(G) &= \min_{\pi, k\text{-partition}} \gamma^*(\pi). \end{aligned}$$

Lemma 4.1 *If π is a k -partition of $V(G)$, then*

$$\gamma(\pi) \leq \frac{1}{k-1},$$

with equality if and only if π is a regular coloring.

Proof.

$$\begin{aligned}\gamma(\pi) &= \frac{\min_{i \neq j} e(S_i, S_j)}{\max_i \text{vol } S_i} \leq \frac{\frac{1}{\binom{k}{2}} \left(\frac{\text{vol } G}{2} \right)}{\max_i \text{vol } S_i} \\ &\leq \frac{\frac{1}{\binom{k}{2}} \left(\frac{\text{vol } G}{2} \right)}{\frac{1}{k} \text{vol } G} = \frac{1}{k-1}.\end{aligned}$$

The first inequality follows from assuming that all edges of the graph are equally distributed between the classes, and if this holds with equality, then the second inequality (which is immediate) also holds with equality. \square

Lemma 4.2 *If π is a k -colouring of G , then*

$$\gamma^*(\pi) \geq \frac{1}{k-1},$$

with equality if and only if π is a regular colouring.

Proof. The proof is very similar to that of Lemma 4.1. \square

It is fair to say that the larger $\gamma(\pi)$ is, the better π is as an attempt to regularly color G . Likewise, the smaller $\gamma^*(\pi)$ is, the more π looks like a clustering with clusters of roughly the same number of edges. These comments motivate the definitions of $\psi_k(G)$ and $\phi_k(G)$. Note that $\phi_2(G)$ is indeed the so called Cheeger constant of G .

Lemma 4.3 *Let π be a k -partition of $V(G)$, G/π be the quotient graph, and $0 = \theta_0 \leq \theta_1 \leq \dots \leq \theta_{k-1}$ be the eigenvalues of $\mathbf{L} = \mathbf{L}(G/\pi)$. Then*

$$k\gamma(\pi) \leq \theta_1 \quad \text{and} \quad \theta_{k-1} \leq k\gamma^*(\pi).$$

Proof. Recall that \mathbf{N} is the diagonal matrix recording the row sums of \mathbf{L} . From the Rayleigh quotient expression for the eigenvalues and the Courant-Fisher Theorem (see for instance [4, Chapter 2]), it follows that

$$\begin{aligned}\theta_1 &= \min_{\mathbf{v} \perp \mathbf{N} \cdot \mathbf{1}} \frac{(\mathbf{N}^{1/2} \mathbf{v})^T \mathbf{L} (\mathbf{N}^{1/2} \mathbf{v})}{(\mathbf{N}^{1/2} \mathbf{v})^T (\mathbf{N}^{1/2} \mathbf{v})} \\ &= \min_{\mathbf{v} \perp \mathbf{N} \cdot \mathbf{1}} \frac{\sum_{i,j} (\mathbf{v}_i - \mathbf{v}_j)^2 e(S_i, S_j)}{\sum_i \mathbf{v}_i^2 \text{vol } S_i} \\ &\geq \min_{\mathbf{v} \perp \mathbf{1}} \frac{\sum_{i,j} (\mathbf{v}_i - \mathbf{v}_j)^2 e(S_i, S_j)}{\sum_i \mathbf{v}_i^2 \text{vol } S_i} \\ &\geq \gamma(\pi) \min_{\mathbf{v} \perp \mathbf{1}} \frac{\sum_{i,j} (\mathbf{v}_i - \mathbf{v}_j)^2}{\sum_i \mathbf{v}_i^2} \\ &= \gamma(\pi)k,\end{aligned}$$

where the last equality follows from the fact that all non-zero eigenvalues of the conventional Laplacian matrix $(k-1)\mathbf{I} - \mathbf{A}(K_k)$ of the complete graph on k vertices are equal to k . The other inequality from the statement follows analogously. \square

Theorem 4.4 *Let G be graph. Fix k between 1 and n . Recall the definitions of $\psi_k(G)$ and $\phi_k(G)$ from the beginning of this section. Again, let $\lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1}$ be the eigenvalues of $\mathbf{L}(G)$, the normalized Laplacian matrix of G . Then*

- (i) $\psi_k(G)k \leq \lambda_{n-(k-1)}$, and
- (ii) $\phi_k(G)k \geq \lambda_{k-1}$.

Proof. This is an immediate consequence of Lemma 4.3 and interlacing. \square

Corollary 4.5 *Suppose G has a regular k -colouring, and $\sigma_{k-1} = k$. Then $\lambda_{n-1} = k/(k-1)$.*

Proof. From Lemma 4.1, it follows that $\psi_k(G) = 1/(k-1)$. From Theorem 4.4, we have $k/(k-1) \leq \lambda_{n-(k-1)}$, which, in addition to $\sigma_{k-1} = k$, leads to $\lambda_{n-1} = k/(k-1)$. \square

The concept of regular colourings also interacts nicely with the results in the previous section.

Theorem 4.6 *Suppose $\lambda_{n-1} = k/(k-1)$ and that G admits k -colouring. Then any k -colouring of G is regular.*

Proof. From Theorem 2.2, it follows that $\sigma_{k-1} \geq k$, and thus

$$\lambda_{n-(k-1)} = \dots = \lambda_{n-1} = \frac{k}{k-1}.$$

As a consequence, just like we argued in Theorem 3.4, it follows that all eigenvalues of the quotient graph are equal to $k/(k-1)$, except for the 0 eigenvalue. It is straightforward to verify that the quotient graph is a complete graph where all edges have the same edge weight. \square

Corollary 4.7 *Suppose $\lambda_{n-1} = k/(k-1)$ and that G admits k -colouring. Then any vertex a has precisely $d(a)/(k-1)$ neighbours of each colour different from its own.*

Proof. Let π be a k -colouring. From Theorem 3.4 and Theorem 4.6, we know that π is equitable with respect to $\mathbf{D}^{-1}\mathbf{A}$ and is a regular colouring. That is, there are constants α and β such that

$$\mathbf{L}(G)\mathbf{P} = \mathbf{P}(\alpha\mathbf{I} + \beta\mathbf{J}),$$

where \mathbf{J} is the all 1s matrix. From this it follows that the rows of $\mathbf{L}(G)\mathbf{P}$ are constant, with the exception of the entry in the column that corresponds to the class that contains the vertex of the row. This is equivalent to what we wanted to prove. \square

Note that Corollary 4.7 offers a quite strong integrality condition on the degrees of all vertices of a graph that satisfy Lemma 3.3 with equality.

5 Concluding remarks

A natural question at this point is whether there are examples of graphs whose colourings are equitable with respect to $\mathbf{D}^{-1}\mathbf{A}$ or regular. A *hypergraph* is a graph in which edges are allowed to have more than 2 vertices. A uniform hypergraph has all edges with the same number of vertices. A colouring of the vertex set of a hypergraph is strong if the vertices in each edge all have distinct colours. Finally, a linear hypergraph is such that any two edges intersect in at most one vertex. The standard reference is [1]. Linear hypergraphs appear naturally in connection to partial linear spaces.

Given an m -uniform linear hypergraph, one forms the underlying graph by replacing each edge by a clique on m vertices. Thus m -uniform hypergraphs correspond to the union of several copies of K_m so that two of these copies share no edge in common. Thus, strongly colouring the hypergraph or properly colouring its underlying graph amounts to the same task. The topic of colouring such graphs has received a considerable amount of attention because of the well-known Erdős-Faber-Lovász conjecture, which proposes that any m -uniform linear hypergraph on m edges is m -strongly colourable (it obviously cannot be coloured with fewer colours). Now we observe that any m -strong colouring of an m -uniform linear hypergraph on e hyperedges (or any colouring of its underlying graph G) must necessarily be

- equitable with respect $\mathbf{D}(G)^{-1}\mathbf{A}(G)$, as for any fixed colour, the number of neighbours of this colour that a vertex has is proportional to the its degree; and
- regular, as each colour class has precisely e edges in G towards any other colour class.

We checked several graphs G which are the underlying graph of an m -uniform linear hypergraph and that satisfy $\chi(G) = m$. For all of them we saw $\lambda_{n-1} = m/(m-1)$, a fact that we do not know yet how to explain. We therefore ask the question

- if G admits an m -colouring which is equitable with respect to $\mathbf{D}^{-1}\mathbf{A}$ and regular, is it true that $\lambda_{n-1} = m/(m-1)$?

Note that an affirmative answer to this question immediately implies that $\lambda_{n-1} = m/(m-1)$ is a necessary condition for the Erdős-Faber-Lovász conjecture to hold for a graph.

Another natural question is how the bound in Theorem 2.2 compares to other eigenvalue bounds. According to our computations, in most cases it tends to be better than Hoffman's bound but slightly worse than the bound in [4, Proposition 3.6.3 (i)] due to Haemers. However, it seems that there are several cases in which our bound performs better, including many of the underlying graphs of uniform linear hypergraphs.

References

- [1] Berge, C., “Graphs and Hypergraphs,” North-Holland Publishing, Amsterdam, 1973.
- [2] Bollobás, B. and V. Nikiforov, *Graphs and Hermitian matrices: eigenvalue interlacing*, Discrete Mathematics **289** (2004), pp. 119–127.
URL <https://www.sciencedirect.com/science/article/pii/S0012365X04003644>
- [3] Bollobás, B. and V. Nikiforov, *Graphs and Hermitian matrices: Exact interlacing*, Discrete Mathematics **308** (2008), pp. 4816–4821.
URL <https://www.sciencedirect.com/science/article/pii/S0012365X07006991>
- [4] Brouwer, A. E. and W. H. Haemers, “Spectra of Graphs,” Universitext, Springer, New York, 2012, xiv+250 pp.
URL <http://dx.doi.org/10.1007/978-1-4614-1939-6>
- [5] Butler, S. K., “Eigenvalues and structures of graphs,” Ph.D. thesis, UC San Diego (2008).
- [6] Chung, F., “Spectral graph theory (CBMS regional conference series in mathematics, No. 92),” American Mathematical Soc., 1996.
- [7] Godsil, C. and G. Royle, “Algebraic Graph Theory,” Springer-Verlag, New York, 2001, xx+439 pp.
- [8] Haemers, W. H., *Eigenvalue methods*, Packing and covering in combinatorics (A. Schrijver ed.), Math. Centre Tract **106** (1979), pp. 15–38.
- [9] Haemers, W. H., *Interlacing eigenvalues and graphs*, Linear Algebra and its Applications **226-228** (1995), pp. 593–616.
URL <http://linkinghub.elsevier.com/retrieve/pii/0024379595001992>
- [10] Nikiforov, V., *Chromatic number and spectral radius*, Linear Algebra and its Applications **426** (2007), pp. 810–814.
URL <http://www.sciencedirect.com/science/article/pii/S0024379507002704>