












Identification and characterization of pathogenic and multidrug-resistant bacteria in feral pigeons surrounding a veterinary hospital in Minas Gerais, Brazil

Jordana Almeida Santana¹  Carolina Pantuzza Ramos¹  Brendhal Almeida Silva¹ 
Graciela Kunrath Lima¹  Alexandra Tiso Comerlato¹  Amanda Cristina Araújo¹ 
Salene Angelini Colombo¹  Gustavo Canesso Bicalho¹  Rodrigo Otávio Silveira Silva^{1*} 

¹Faculdade de Veterinária, Universidade Federal de Minas Gerais (UFMG), 31270-901, Belo Horizonte, MG, Brasil. E-mail: rodrigo.otaviosilva@gmail.com.*Corresponding author.

ABSTRACT: Pigeons are known for their capacity to harbor and spread several zoonotic agents. Studies have suggested that pigeons are also relevant disseminators of multidrug-resistant strains. In this study, pigeons surrounding a veterinary hospital were sampled and tested for the presence of pathogenic *Escherichia coli*, *Salmonella* spp., *Staphylococcus* spp., and *Clostridioides (Clostridium) difficile*. *E. coli* isolates from 19 (40.4%) pigeons tested positive for the *E. coli* heat-stable enterotoxin 1 (EAST1)-encoding gene. The intimin-encoding gene (*eae*) of enteropathogenic *E. coli* (EPEC) was found in one isolate (2.1%). *Salmonella* spp. were found in nine (19.1%) pigeons, all from the first capture event ($P < 0.001$). *S. Typhimurium* and *S. Heidelberg* were isolated from six and three pigeons, respectively. Enterobacterial repetitive intergenic consensus (ERIC-PCR) of the *Salmonella* spp. isolates suggested that eight of the nine strains had a high genetic similarity, supporting the hypothesis of an outbreak of salmonellosis in these pigeons. Twenty (42.5%) staphylococcal isolates were recovered from 18 (38.3%) pigeons. Eight different species were detected, with *S. xylosus* being the most frequent. Two (4.3%) *C. difficile* strains were isolated. Three isolates, one each of *S. Typhimurium*, *S. aureus*, and *C. difficile*, were classified as multidrug-resistant strains. The present research suggested that pigeons residing in urban areas can act as reservoirs and disseminators of pathogenic bacteria, including nosocomial pathogens, such as diarrheagenic *E. coli* and multidrug-resistant *Staphylococcus* spp., *C. difficile*, and *Salmonella* spp.

Key words: *Salmonella*, *Clostridioides (Clostridium) difficile*, pigeons, *Staphylococcus*, synanthropes.

Identificação de bactérias patogênicas e multirresistentes a antimicrobianos em pombos urbanos no entorno de um Hospital Veterinário em Belo Horizonte, Minas Gerais, Brasil

RESUMO: Pombos urbanos são conhecidos pela sua capacidade de carrear e disseminar diversos agentes zoonóticos. Estudos tem sugerido que pombos são também relevantes na disseminação de estirpes resistentes a múltiplas drogas. No presente estudo, pombos no ambiente de um hospital veterinário foram amostrados em três diferentes períodos e testados para a presença de *Escherichia coli* patogênica, *Salmonella* spp., *Staphylococcus* spp. e *Clostridioides (Clostridium) difficile*. Isolados de *E. coli* de 19 pombos (40.4%) foram positivos para o gene codificador da toxina EAST1. O gene codificador de intimina (*eae*) do patótipo *E. coli* enteropatogênica foi encontrada em um isolado (2.1%). *Salmonella* spp. foi encontrada em nove pombos (19.1%), sendo todos isolados do primeiro período de captura ($P < 0.001$). *S. Typhimurium* foi isolado de seis animais e *S. Heidelberg* de três. A tipagem molecular de isolados de *Salmonella* spp. por ERIC-PCR demonstrou que oito estirpes possuíam alta similaridade genética entre si, sugerindo a ocorrência de um surto de salmonelose nos animais carreadores. Vinte *Staphylococcus* (42.5%) foram isolados de 18 animais (38.3%). Oito diferentes espécies foram detectadas, sendo *S. xylosus* a mais frequente. Duas estirpes de *C. difficile* não-toxigênica (4.3%) foram isoladas. Uma estirpe de *S. Typhimurium*, uma de *S. aureus* e um isolado de *C. difficile* foram classificados como resistentes a múltiplas drogas antimicrobianas. O presente estudo sugere que pombos capturados no ambiente do hospital veterinário podem atuar como reservatórios e disseminadores de bactérias patogênicas e envolvidas em infecção hospitalar, incluindo *E. coli* diarréio-gênica e *Staphylococcus* sp., *C. difficile* e *Salmonella* spp multirresistente.

Palavras-chave: *Salmonella*, *Clostridioides (Clostridium) difficile*, pestes, *Staphylococcus*, sinantrópicos.

INTRODUCTION

Urban feral pigeons (*Columba livia* f. *urbana*) are birds with a worldwide distribution commonly reported in most large cities (FERMAN et al., 2010; GARGIULO et al., 2014; SPENNEMANN & WATSON, 2017). The large population densities of

urban pigeons in cities are mainly related to the large availability of food, lack or less number of predators, and the built environment resembling the original habitat of urban pigeon ancestors (SPENNEMANN & WATSON, 2017). They are known for their capacity to harbor and spread several zoonotic agents, such as *Cryptococcus neoformans* and *Salmonella*

spp. (HAAG-WACKERNAGEL & MOCH, 2004; DUTTA et al., 2013; HAESSENDONCK et al., 2016; SPENNEMANN & WATSON, 2017). Recently, studies have suggested that pigeons can also be relevant carriers of multidrug-resistant (MDR) bacteria (BORGES et al., 2017; TORRES-MEJÍA et al., 2018; CUNHA et al., 2019). Thus, close contact with pigeons and their feces is a risk for people who frequently share the same environment with these animals (HAAG-WACKERNAGEL & MOCH, 2004; TANAKA et al., 2005; GARGIULO et al., 2014). This is the scenario for employees, students, owners, and animals in the Veterinary Hospital of the Federal University of Minas Gerais (HV-UFGM), where the pigeon population has increased markedly in the last few years.


Despite the known importance of pigeons as reservoirs of several pathogens, no studies have evaluated them in a university environment, and their possible role in the epidemiology of some relevant nosocomial infections in companion animals is still unknown. In addition, the presence of *Staphylococcus*

sp. and *Clostridioides* (previously *Clostridium*) *difficile* in pigeons has been reported in a few studies; however, none of the studies have been conducted in Brazil. Thus, this study investigated the occurrence and antimicrobial susceptibility of enteric bacteria isolated from pigeons captured in an urban area surrounding a veterinary hospital in Belo Horizonte, Minas Gerais, Brazil.

MATERIALS AND METHODS

Samples

This study was conducted at the Veterinary Hospital of the Federal University of Minas Gerais (HV-UFGM), which receives approximately 35,000 animals per year, including dogs, cats, horses, cattle, and wild animals. It is located inside the university campus and is surrounded by a green area (Figure 1). This study was motivated by the increase in pigeon population in the last few years in HV-UFGM, raising the need for a better understanding of the risks associated with these birds. Forty-seven



pigeons (*Columba livia*) were sampled after three capture campaigns (January/2019, July/2019, and January/2020), captured using mist nets (30 mm mesh size; four shelves, ap. 3 m high x 9 m length) during early mornings, in places where large animals fed. The first capture event occurred in January 2019 during the wet season, with a monthly mean precipitation of 2.3 mm and a mean temperature of 25.6 °C (INMET, 2019), with 18 birds captured. The second capture event occurred in July of the same year during the dry season, with 0 mm of precipitation and a mean temperature of 19.2 °C (INMET, 2019), with 13 birds captured. The third and last capture occurred in January 2020, with a monthly mean precipitation of 1.3 mm and a mean temperature of 21.5 °C (INMET, 2020), with 16 birds captured. The pigeons were physically restrained, given colored leg rings for identification and monitoring purposes, and left to rest in individual cages. Fresh feces were collected immediately after dropping and stored in microtubes using sterile spatulas (ROSARIO MEDINA et al., 2017). The samples were stored in a transport box with ice packs and sent to the Bacterial and Research Laboratory of the Veterinary School of UFMG for immediate processing. Each pigeon was returned to its environment once sample collection was completed.

Escherichia coli

For *E. coli* isolation, the samples were plated onto MacConkey agar (Difco, USA) and incubated at 37 °C for 24 h (RAMOS et al., 2019a). Three lactose-fermenting colonies were identified using polymerase chain reaction (PCR) and subjected to subsequent reactions for phylogenetic group characterization (A, B1, B2, C, D, E, and F) (MCDANIELS et al., 1996; CLERMONT et al., 2013). Virulence genes associated with pathogenic *E. coli*, such as enterotoxigenic *E. coli* (ETEC), enteropathogenic *E. coli* (EPEC), Shiga toxin-producing *E. coli* (STEC), enterohemorrhagic *E. coli* (EHEC), necrotoxicogenic *E. coli* (NTEC), and enteroaggregative *E. coli* (EAEC), were also identified using PCR (BLANCO et al., 1996; YAMAMOTO & NAKAZAWA, 1997; FRANCK et al., 1998). Antibiotic resistance patterns of all *E. coli* isolates that tested positive for virulence factors were evaluated using the disc diffusion method (DDM), according to the Clinical and Laboratory Standards Institute (CLSI) manual (CLSI, 2017; CLSI, 2018). The following antimicrobial agents, commonly used in human and animal clinical practice, were tested: chloramphenicol (30 µg), trimethoprim/sulfamethoxazole (25 µg), ceftriaxone

(30 µg), ceftiofur (30 µg), amoxicillin/clavulanic acid (30 µg), ampicillin (10 µg), tetracycline (30 µg), enrofloxacin (5 µg), ciprofloxacin (5 µg), gentamicin (10 µg), and amikacin (30 µg) (DME, BRA).

Salmonella spp.

For *Salmonella* spp. isolation, cloacal samples were pre-enriched in Rappaport broth (Oxoid, USA) and plated onto Hektoen enteric agar (Oxoid, USA) (RAMOS et al., 2019a). Sulfite-reducing colonies were identified as *Salmonella* spp. by genus-specific PCR, according to KUANG et al. (2015). Antigenic characterization was done according to the White-Kauffmann-Le Minor Scheme (LE MINOR & POPOFF, 1987) at the Brazilian National Reference Laboratory of Enterobacteria of the Oswaldo Cruz Foundation, followed by species, subspecies, and serotype identification (GRIMONT & WEILL, 2007). *Salmonella* spp. strains were fingerprinted using enterobacterial repetitive intergenic consensus (ERIC)-PCR and analyzed using Bionumerics 7.6 software (Applied Maths NV, Belgium) to evaluate the genetic diversity between isolates from different pigeon samples (VERSALOVIC et al., 1991; RAMOS et al., 2019b). Additionally, DDM was used to evaluate the resistance patterns of *Salmonella* spp. isolates to antimicrobial agents (CLSI, 2017; CLSI, 2018) using the following drugs: chloramphenicol (30 µg), trimethoprim/sulfamethoxazole (25 µg), ceftriaxone (30 µg), ceftiofur (30 µg), amoxicillin/clavulanic acid (30 µg), ampicillin (10 µg), tetracycline (30 µg), nalidixic acid (30 µg), enrofloxacin (5 µg), and ciprofloxacin (5 µg) (DME, BRA).

Staphylococcus spp.

For *Staphylococcus* spp. isolation, fecal samples were first suspended in 0.85% saline solution. The resultant solution (100 µL) was then streaked onto mannitol salt agar (Difco Laboratories Inc., USA), which was incubated at 37 °C for 24 h. Colonies were sub-cultured on brain heart infusion agar (Difco Laboratories Inc., USA) and identified by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS; Bruker Daltonics, Germany). A cutoff log score of 2 was used to validate identification at the species level, as recommended by the manufacturer. The strains were then subjected to DNA extraction (PITCHER et al., 1989), and methicillin-resistant staphylococci were investigated by detecting the *mecA* gene (MURAKAMI et al., 1991). In addition, isolates identified as *Staphylococcus intermedius* group (SIG) using MALDI-TOF were further confirmed by

multiplex PCR of the *nuc* gene (SASAKI et al., 2010). Non-SIG isolates with MALDI-TOF score under 2.0 were subjected to sequencing of the 16S ribosomal RNA gene as described by FOX et al. (2011). Antimicrobial susceptibility tests were performed using DDM in agar, according to the Clinical and Laboratory Standards Institute (CLSI) documents M100-S30 and VET08 (CLSI 2018; CLSI, 2020). The following antimicrobials were tested: ceftiofur (30 µg), penicillin (10 units), tetracycline (30 µg), trimethoprim/sulfamethoxazole (25 µg), chloramphenicol (30 µg), erythromycin (15 µg), clindamycin (2 µg), gentamicin (10 µg), and ciprofloxacin (5 µg) (DME, BRA). *Staphylococcus aureus* ATCC 25923 was used as the control strain. Isolates were considered MDR when resistant to three or more classes of antimicrobial agents (SWEENEY et al., 2018).

Clostridioides difficile

Samples were incubated in 96% ethanol for 30 min (1:1), and aliquots of 10 µL were plated on cycloserine-ceftiofur fructose agar supplemented with 7% horse blood and 0.1% sodium taurocholate (Sigma, USA) (SILVA et al., 2013). After incubation in an anaerobic atmosphere at 37 °C for 72 h, characteristic *C. difficile* isolates (flat, irregular, and with a ground-glass appearance) were subjected to multiplex PCR to identify the housekeeping gene (*tpi*) and virulence genes of toxin A (*tcdA*), toxin B (*tcdB*), and binary toxin (*cdtB*) (SILVA et al., 2011). The minimal inhibitory concentrations (MICs) of metronidazole, vancomycin, clindamycin, moxifloxacin, ciprofloxacin, erythromycin, rifampicin, and tetracycline were determined using Etest strips (bioMérieux Marcy l'Etoile, France) in Brucella agar (Oxoid, USA) with 5% lysed blood supplemented with hemin (Difco Laboratories, USA) and vitamin K (Sigma-Aldrich Co., USA). MIC values were interpreted according to the clinical breakpoints of the CLSI and European Committee on Antimicrobial Susceptibility Testing guidelines (PIRŠ et al., 2013; CLSI, 2015; EUCAST, 2019).

Statistical analysis

The association between capture events, presence of the tested pathogens, and frequency of each isolated microorganism was evaluated using chi-square and Fisher's exact tests. The chi-squared test for adherence was used to evaluate the distribution of variables. All statistical analyses were performed using GraphPad Prism v.8 (GraphPad Software,

San Diego, CA, USA). Differences were considered statistically significant at $P < 0.05$.

RESULTS

Escherichia coli

E. coli was identified in 42 (89.3%) pigeons, with a total of 120 isolated strains (Table 1). All seven phylogenetic groups of *E. coli* were identified (Table 2) in the sampled pigeons, with B1 being the most common phylogroup (52/120 strains, 43.3%). A total of 41 strains (34%) from 19 (40.4%) pigeons tested positive for enteroaggregative *E. coli* heat-stable enterotoxin 1 (EAST1) encoding gene. The intimin-encoding gene (*eae*) of EPEC was detected in one isolate (2.1%), which was also positive for EAST1. No resistance to the tested antimicrobials was observed in the 41 *E. coli* isolates that tested positive for virulence factors.

Salmonella spp.

Salmonella spp. were found in nine (19.1%) pigeons, all from the first capture event ($n=18$; isolation rate of 50%), indicating a strong difference between the first and the other two events ($P < 0.001$). *S. Typhimurium* and *S. Heidelberg* were isolated from six (66.7%) and three pigeons, respectively. One bird died a few days after the capture event (Figure 2; isolate "PB13"). Macroscopically, the bird had hepatomegaly, and the serosa of the small intestine was diffusely hyperemic with hemorrhagic intestinal contents. *S. Typhimurium* was isolated again from the excreta and liver of this bird. ERIC-PCR of the *Salmonella* spp. isolates suggested that eight out of the nine strains had high genetic similarity (Figure 2). These strains showed no resistance to all antimicrobials tested, while the remaining strain (*S. Typhimurium*) was resistant to trimethoprim/sulfamethoxazole, amoxicillin/clavulanic acid, ampicillin, tetracycline, enrofloxacin, ciprofloxacin, and nalidixic acid. Thus, it was classified as an MDR strain (MAGIORAKOS et al., 2012).

Staphylococcus spp.

A total of 20 (42.5%) staphylococcal isolates were recovered from 18 (38.3%) pigeons of the 47 captured pigeons. Eight different species were detected, with *S. xylosus* (30%) being the most common (Table 1). Overall, 7 (35%) isolates were resistant to at least one antimicrobial agent, whereas 13 (65%) were susceptible to all the tested compounds. Some isolates were resistant to tetracycline (35.5%), penicillin G (15%), erythromycin (5%), and clindamycin (5%). No significant differences were

Table 1 - Frequency of *Escherichia coli*, *Salmonella* spp., *Staphylococcus* spp., and *Clostridioides difficile* isolated from free-living pigeons (n=47) in three capture events in the Veterinary Hospital of Federal University of Minas Gerais (Belo Horizonte, Minas Gerais, Brazil).

Pathogen	-----Capture (%)-----			Total (n=47)
	First	Second	Third	
<i>E. coli</i>	18/18 (100)	10/13 (76.9)	14/16 (87.5)	42/47 (89.3)
EAST-1	8/18 (44.4)	6/13 (46.1)	5/16 (31.3)	19/47 (40.4)
EPEC (<i>eae</i>)	1/18 (5.5)	0/13 (0)	0/16 (0)	1/47 (2.1)
<i>Salmonella</i> spp.	9/18 (50) ^a	0/13 (0) ^b	0/16 (0) ^b	9/47 (12.8)
<i>S. Typhimurium</i>	6/18 (33.3)	0/13 (0)	0/16 (0)	6/47 (6.4)
<i>S. Heildeberg</i>	3/18 (18.7)	0/13 (0)	0/16 (0)	3/47 (6.4)
<i>Staphylococcus</i> spp.	7/18 (38.9)	6/13 (46.1)	7/16 (43.7)	20/47 (42.5)
<i>S. xylosum</i>	3/18 (16.7)	2/13 (15.4)	1/16 (6.3)	6/47 (12.8)
<i>S. sciuri</i>	2/18 (11.1)	1/13 (7.7)	2/16 (12.5)	5/47 (10.6)
<i>S. lentus</i>	0/18 (0)	0/13 (0)	3/16 (18.7)	3/47 (6.4)
<i>S. haemolyticus</i>	0/18 (0)	2/13 (15.4)	0/16 (0)	2/47 (4.2)
<i>S. aureus</i>	0/18 (0)	0/13 (0)	1/16 (6.3)	1/47 (2.1)
<i>S. intermedius</i>	1/18 (5.6)	0/13 (0)	0/16 (0)	1/47 (2.1)
<i>S. succinus</i>	0/18 (0)	1/13 (7.7)	0/16 (0)	1/47 (2.1)
<i>S. schleiferi</i>	1/18 (5.6)	0/13 (0)	0/16 (0)	1/47 (2.1)
<i>C. difficile</i> (Non-toxicogenic)	2/18 (11.1)	0/13 (0)	0/16 (0)	2/47 (4.2)

Different lower-case letters indicate a significant difference ($P < 0.05$).

Legend: EPEC - Enteropathogenic *Escherichia coli*; EAST-1 - Enteropathogenic *Escherichia coli*.

reported in resistance to these antimicrobials. One *S. aureus* isolate showed resistance to penicillin G, erythromycin, and clindamycin; and therefore, was classified as MDR. All isolates were susceptible to cefoxitin, chloramphenicol, gentamicin, ciprofloxacin, and trimethoprim-sulfamethoxazole and were negative for *mecA*.

Clostridioides difficile

Two (4.3%) *C. difficile* isolates were recovered from the pigeons. Both were nontoxicogenic (A-B-CDT-).

One isolate was classified as MDR due to resistance to erythromycin, rifampicin, and tetracycline, while both isolates were susceptible to all other antimicrobials tested, including metronidazole and vancomycin.

DISCUSSION


Despite the known importance of pigeons as reservoirs of several pathogens, no studies have evaluated them in a university environment and their role in the epidemiology of nosocomial bacteria.

Table 2 - Phylogroups of *Escherichia coli* isolates (n=120) from pigeons (n=47) captured in the Veterinary Hospital of Federal University of Minas Gerais (Belo Horizonte, Minas Gerais, Brazil).

<i>E. coli</i>	-----Phylogenetic groups-----									Total
	A	B1	B2	C	D	E	F	Clade ²	Unassignable ¹	
Isolates (%)	11 (9.2)	52 (43.3)	12 (10) ^a	18 (15)	3 (2.5)	16 (13.3) ^a	1 (0.8)	4 (3.3)	3 (2.5)	120 (100)

¹Identified as *E. coli* but not corresponding to any of the phylogroups according to CLERMONT et al. (2013).

²Clade 1, 2, 3, 4, or 5.



This study revealed the presence of zoonotic MDR pathogens in pigeons captured in the surroundings of a veterinary hospital, suggesting that pigeons can act as reservoirs and disseminators of diarrheagenic *E. coli* and MDR *Staphylococcus* spp., *C. difficile*, and *Salmonella* spp.

The high isolation rate of *E. coli* from phylogroup B1 was similar to that reported in previous studies on urban pigeons (GORDON & COWLING, 2003; SILVA et al., 2009; GHANBARPOUR & DANESHDOOST, 2012). Interestingly, we identified two virulence factors in the *E. coli* isolates. *E. coli* isolates positive for the EAST-1 encoding gene were detected in more than 40% of the birds, whereas the *eae* encoding gene from EPEC was detected in a single isolate. The high frequency of pigeons positive for EAST-1 was surprising, since this virulence factor, previously associated with outbreaks of diarrhea in humans (ZHOU et al., 2002; SUKKUA et al., 2017), has never been reported in *E. coli* isolates from pigeons. Further, pigeons are known reservoirs of EPEC (SILVA et al., 2009; GHANBARPOUR & DANESHDOOST, 2012; SACRISTÁN et al., 2014; BORGES et al., 2017; TORRES-MEJÍA et al., 2018), a major cause of childhood diarrhea worldwide (CROXEN et al., 2013; TORRES-MEJÍA et al., 2018). Together, these results reinforce

the role of pigeons as potential reservoirs of zoonotic *E. coli* pathotypes.

We tested all 42 *E. coli* isolates that were positive for virulence factors for antimicrobial resistance. Results revealed that all isolates were susceptible to all antimicrobials tested, surprisingly showing that despite living in a heavily anthropized environment, the sampled pigeons had *E. coli* isolates with no drug resistance. Our results are in contrast with previous studies that described the occurrence of MDR *E. coli* strains in pigeons in several countries (GHANBARPOUR & DANESHDOOST, 2012; BORGES et al., 2017; KARIM et al., 2020).

The fecal shedding of *Salmonella* spp. in the sampled pigeons (19.1%) was higher than that found in several previous studies, which often reported frequencies of up to 10% (DOVC et al., 2004; TANAKA et al., 2005; PEDERSEN et al., 2006; GARGIULO et al., 2014; HAESSENDONCK et al., 2016; CARVALHO et al., 2020; KACZOREK-LUKOWSKA et al., 2020). Interestingly, the isolation of *Salmonella* spp. was significantly associated with the first capture event ($P < 000.1$) conducted during the rainy season. This result raised the hypothesis of an outbreak of salmonellosis in the pigeons during this sampling period, which was reinforced after the re-isolation of *Salmonella* from

the intestinal content and liver of a pigeon that died a few days after the capture event. Furthermore, the post-mortem alterations observed in this pigeon are commonly reported in pigeons and other species with salmonellosis (SAWA & HIRAI, 1981; OLIVEIRA et al., 2019; RAMOS et al., 2021).

To better understand this possible outbreak, one *Salmonella* isolate from each animal was subjected to ERIC-PCR, a method with high discriminatory power already in use for *Salmonella* typing in outbreaks (PURIGHALLA et al., 2017; RAMOS et al., 2019b). The high similarity among the eight isolates, as suggested by ERIC-PCR, strongly suggests the occurrence of an outbreak of salmonellosis in these pigeons. Previous studies have indicated that, although, pigeons are commonly asymptomatic carriers of *Salmonella* spp., salmonellosis may occur in urban pigeons, which could significantly increase the isolation rate of this agent in the affected pigeons (SAWA & HIRAI, 1981; TIZARD, 2004; HOELZER et al., 2011; DUTTA et al., 2013; ROCHA-E-SILVA et al., 2014). Additionally, salmonellosis outbreaks in animals seem to be more common during the rainy season, which may be a risk factor for bacterial spread (PANGLOLI et al., 2008; RAMOS et al., 2019b).

Notably, we detected two well-known zoonotic *Salmonella* spp. serovars, *S. Typhimurium* and *S. Heidelberg*, in our study. Previous studies have demonstrated that *Salmonella enterica* isolates from urban pigeons commonly belong to serotype *Typhimurium* (DUTTA et al., 2013; OSMAN et al., 2013; ROCHA-E-SILVA et al., 2014; HAESSENDONCK et al., 2016; TORRES-MEJÍA et al., 2018; CARVALHO et al., 2020; KACZOREK-LUKOWSKA et al., 2020). It is also known that *S. Typhimurium* and *S. Heidelberg* are important foodborne pathogens (CDC, 2014), which can also infect humans after direct contact with healthy or diseased animals (HALE et al., 2012; RAMOS et al., 2019b). Transmission of *Salmonella* spp. from pigeons to humans has already been reported; however, studies associating human salmonellosis with pigeon contact are scarce (HAAG-WACKERNAGEL & MOCH, 2004; SPENNEMANN & WATSON, 2017). Despite this, the common shedding of these zoonotic *Salmonella* serotypes by urban pigeons and their close contact with humans should be considered a potential risk to human health, especially for those most susceptible, such as children and immunocompromised individuals (HALE et al., 2012).

Most *Salmonella* spp. isolates showed no resistance to the antimicrobials tested. Interestingly,

all isolates were genetically similar according to ERIC-PCR analysis, which also contributed to the hypothesis of an outbreak. Conversely, one *S. Typhimurium* strain was resistant to seven antimicrobials and thus, was classified as MDR (MAGIORAKOS et al., 2012). According to the World Health Organization (WHO, 2019), there are specific public health concerns related to the spread of fluoroquinolone-resistant *Salmonella* spp. In addition, β -lactams and sulfonamides are commonly used antimicrobials for the treatment of *Salmonella* spp. infections in animals and humans, and resistance to both important drugs may severely reduce the treatment options (KUANG et al., 2015).

More than one-third of the captured pigeons were positive for staphylococci and eight different species were recovered (Table 1). According to other studies, pigeons seem to harbor a high diversity of staphylococcal species, whereas other animals, such as free-living reptiles and rodents, seem to show a more homogeneous colonization pattern (SCHWARZ & WERCKENTHIN, 1994; ZIGO, 2017; SANTANA et al., 2021; SANTANA et al., 2022). It may be that the close and daily contact of the pigeons with humans and animals, combined with the hospital environment, may have influenced this large number of recovered species (SCHWARZ & WERCKENTHIN, 1994; ZIGO, 2017; KAMATHEWATTA et al., 2019).

S. xylosum, the most frequent species reported in our study, has been previously isolated from pigeons and other birds, suggesting commensalism (VELA et al., 2012; MAHMMOUD, 2013; ZIGO, 2017; MATIAS et al., 2018). This coagulase-negative *Staphylococcus* (CoNS) has also been found in different mammals and reptiles (BECKER et al., 2014; RISSI et al., 2015; MATIAS et al., 2018; SANTANA et al., 2021; SANTANA et al., 2022), and despite being mostly labeled as non-pathogenic, it has caused several opportunistic infections in animals and humans (WON et al., 2002; KOKSAL et al., 2009; AKHADDAR et al., 2010; RISSI et al., 2015).

Except for *S. succinus*, all other CoNS isolates have been reported in previous studies on pigeons (SCHWARZ & WERCKENTHIN, 1994; ZIGO, 2017). Similar to *S. xylosum*, all these species can integrate into the microbiota of skin and mucous membranes of different hosts and act as opportunistic pathogens causing distinct infections (RISSI et al., 2015). Notably, *S. haemolyticus* is highly relevant to human health and is the second most frequently isolated CoNS from nosocomial infections (SIDHU et al., 2007; CZEKAJ et al., 2015).

We also isolated *S. intermedius* and coagulase-positive *S. aureus* in this study. Both species have been frequently reported in pigeons (KIZERWETTER-ŚWIDA et al., 2015; CHROBAK-CHMIEL et al., 2021). However, *S. intermedius* is more commonly reported in wild pigeons, and its isolation from domestic pigeons is scarce (KIZERWETTER-ŚWIDA et al., 2015). In contrast, *S. aureus* is widespread among humans and different animal species, facilitating bilateral transmission between distinct hosts (WEESE & VAN DUIJKEREN, 2010). This microorganism is one of the world's most frequent causes of nosocomial infections in humans. It is also a relevant pathogen in dogs, and its treatment is often hampered by resistance to multiple antimicrobials (IPPOLITO et al., 2010; WALTHER et al., 2017; TURNER et al., 2019). Interestingly, *S. aureus* isolated in our study was an MDR strain, which reinforces the hypothesis that pigeons are potential disseminators of MDR staphylococci (KUTKOWSKA et al., 2019; CHROBAK-CHMIEL et al., 2021).

More than one-third of the staphylococci isolates showed resistance to at least one of the antimicrobials tested, mainly to tetracycline and penicillin, which are widely used in human and veterinary medicine (ARGUDÍN et al., 2017; CERBO et al., 2019). It is possible that this specific environment, where the pigeons can get directly and indirectly in contact with humans and healthy and sick animal species, may have influenced the diversity of staphylococci isolated in our study, as well as the frequency of isolates resistant to antimicrobials, as previously indicated in other studies on staphylococci in different animals and settings (FUTAGAWA-SAITO et al., 2007; HAABER et al., 2017; CERBO et al., 2019; KAMATHEWATTA et al., 2019; FROSINI et al., 2020; PALMA et al., 2020). FUTAGAWA-SAITO et al., (2007) compared the resistance profiles of staphylococci isolated from pigeons and suggested that isolates from pigeons that have direct contact with humans and other animals are more associated with resistance to several antimicrobials, which supports the idea that resistance may be acquired because of the mutual coexistence of different species.

Although, *C. difficile* is recognized as an emerging pathogen causing zoonotic diseases in humans (KNIGHT & RILEY, 2019) and has previously been reported to cause nosocomial infections in dogs (WEESE & ARMSTRONG, 2003), little is known about the role of pigeons in the epidemiology of this anaerobic microorganism.

Previous studies of other avian species have suggested that *C. difficile* is either absent or present at a very low frequency (BANDELJ et al., 2011; BURT et al., 2012; BANDELJ et al., 2014). In our study, two *C. difficile* strains were isolated (4.3%), which is lower than the previously reported rate (12.5%) by ANDRÉS-LASHERAS et al. (2017) in a study on pigeons and rodents trapped within pest control programs in pig farms. Notably, the colonization of *C. difficile* in pigeons and other birds seems to be directly linked to environmental contamination (ANDRÉS-LASHERAS et al., 2017), which may explain the difference in its isolation rates. Additionally, one isolate in our study was classified as MDR due to its resistance to erythromycin, rifampicin, and tetracycline. Our results are similar to those previously reported by ANDRÉS-LASHERAS et al. (2017) and reinforce the hypothesis that pigeons may play a role in the transmission of *C. difficile*, including antimicrobial-resistant strains.

Our research suggested that pigeons captured in the surroundings of a veterinary hospital can act as reservoirs and disseminators of pathogenic and nosocomial bacteria, including diarrheagenic *E. coli* and MDR *Staphylococcus* sp., *C. difficile*, and *Salmonella* spp. Together with previous studies, our findings reinforce the importance of pigeon population control owing to their potential role in the spread of zoonotic diseases.

ACKNOWLEDGMENTS

We thank all students and professionals who contributed to pigeon capture and Professor Marcelo Pires Nogueira de Carvalho for clinical evaluation of the birds. This work was supported by funds received from Coordination for the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES – Prêmio CAPES 2015 - 0774/2017), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq - 406402/2018-3), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG - APQ-00524-17), and Pró-Reitoria de Pesquisa da Universidade Federal de Minas Gerais (PRPq/UFMG).

DECLARATION OF CONFLICT OF INTEREST

We have no conflict of interest to declare.

BIOETHICS AND BIOSECURITY COMMITTEE APPROVAL

This study was approved by the Ethical Committee on Animal Use (CEUA) of the Universidade Federal de Minas Gerais (UFMG) under protocol 361/2018 and by the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) under protocol SISBIO 66535-2.

AUTHOR'S CONTRIBUTIONS

All authors contributed equally to the conception and writing of the manuscript. All authors critically revised and approved the final version of the manuscript.

REFERENCES

- AKHADDAR, A. et al. *Staphylococcus xylosum* isolated from an otogenic brain abscess in an adolescent. **Surgical Infections**, v.11, p.559–561, 2010. Available from: <<https://pubmed.ncbi.nlm.nih.gov/20969474/>>. Accessed: Jun. 16, 2022. doi: 10.1089/sur.2010.010.
- ANDRÉS-LASHERAS, S. et al. Presence of *Clostridium difficile* in pig faecal samples and wild animal species associated with pig farms. **Journal of Applied Microbiology**, v.122, p.462–472, 2017. Available from: <<https://pubmed.ncbi.nlm.nih.gov/27990723/>>. Accessed: Jun. 16, 2022. doi:10.1111/jam.13343.
- ARGUDÍN, M. et al. Bacteria from Animals as a Pool of Antimicrobial Resistance Genes. **Antibiotics**, v.6, p.12, 2017. Available from: <<https://pubmed.ncbi.nlm.nih.gov/28587316/>>. Accessed: Jun. 16, 2022. doi: 10.3390/antibiotics6020012.
- BANDELJ, P.; et al. Prevalence and molecular characterization of *Clostridium difficile* isolated from European Barn Swallows (*Hirundo rustica*) during migration. **BMC Veterinary Research**, v.10, p.40, 2014. Available from: <<https://pubmed.ncbi.nlm.nih.gov/24507706/>>. Accessed: Jun. 16, 2022. doi: 10.1186/1746-6148-10-40.
- BANDELJ, P. et al. Zero prevalence of *Clostridium difficile* in wild passerine birds in Europe. **FEMS microbiology letters**, v.321, p.183–185, 2011. Available from: <<https://pubmed.ncbi.nlm.nih.gov/21658108/>>. Accessed: Jun. 16, 2022. doi: 10.1111/j.1574-6968.2011.02333.x
- BECKER, K. et al. Coagulase-negative staphylococci. **Clinical Microbiology Reviews**, v.27, p.870–926, 2014. Available from: <<https://journals.asm.org/doi/10.1128/CMR.00109-13>>. Accessed: Jun. 16, 2022. doi: 10.1128/CMR.00109-13.
- BLANCO, M. et al. Polymerase chain reaction for detection of *Escherichia coli* strains producing cytotoxic necrotizing factor type 1 and type 2 (CNF1 and CNF2). **Journal of Microbiological Methods**, v.26, p.95–101, 1996. Available from: <[https://doi.org/10.1016/0167-7012\(96\)00900-1](https://doi.org/10.1016/0167-7012(96)00900-1)>. Accessed: Jun. 16, 2022. doi: 10.1016/0167-7012(96)00900-1.
- BORGES, C.A. et al. Wild birds and urban pigeons as reservoirs for diarrheagenic *Escherichia coli* with zoonotic potential. **Journal of Microbiology**, v.55, p.344–348, 2017. Available from: <<https://pubmed.ncbi.nlm.nih.gov/28281197/>>. Accessed: Jun. 16, 2022. doi: 10.1007/s12275-017-6523-3.
- BURT, S.A. et al. Vermin on pig farms are vectors for *Clostridium difficile* PCR ribotypes 078 and 045. **Veterinary Microbiology**, v.160, p.256–258, 2012. Available from: <<https://pubmed.ncbi.nlm.nih.gov/22682200/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.vetmic.2012.05.014.
- CARVALHO, D. et al. Antimicrobial susceptibility and detection of virulence-associated genes of *Escherichia coli* and *Salmonella* spp. isolated from domestic pigeons (*Columba livia*) in Brazil. **Folia Microbiologica**, v.65, p.735–745, 2020. Available from: <<https://pubmed.ncbi.nlm.nih.gov/32180120/>>. Accessed: Jun. 16, 2022. doi: 10.1007/s12223-020-00781-w.
- CERBO, A. et al. Tetracyclines: Insights and Updates of their Use in Human and Animal Pathology and their Potential Toxicity. **The Open Biochemistry Journal**, v.13, 2019. Available from: <<https://openbiochemistryjournal.com/VOLUME/13/PAGE/1/ABSTRACT/>>. Accessed: Jun. 16, 2022. doi: 10.2174/1874091X01913010001.
- CDC. Foodborne Diseases Active Surveillance Network (FoodNet): FoodNet Surveillance Report for 2012 (Final Report). Atlanta, Georgia: U.S. Department of Health and Human Services, CDC. 2014.
- CHROBAK-CHMIEL, D. et al. Pigeons as Carriers of Clinically Relevant Multidrug-Resistant Pathogens—A Clinical Case Report and Literature Review. **Frontiers in Veterinary Science**, v.8, 2021. Available from: <<https://pubmed.ncbi.nlm.nih.gov/34109235/>>. Accessed: Jun. 16, 2022. doi: 10.3389/fvets.2021.664226.
- CLERMONT, O. et al. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. **Environmental Microbiology Reports**, v.5, p.58–65, 2013. Available from: <<https://pubmed.ncbi.nlm.nih.gov/23757131/>>. Accessed: Jun. 16, 2022. doi: 10.1111/1758-2229.12019.
- CLINICAL AND LABORATORY STANDARDS INSTITUTE (CLSI). Wayne, PA, USA. Performance Standards for Antimicrobial Susceptibility Testing; **Twenty-Fifth Informational Supplement**, 2015.
- CLINICAL AND LABORATORY STANDARDS INSTITUTE (CLSI). Performance Standards for Antimicrobial Susceptibility Testing. 27th ed. CLSI M100 (ISBN 1-56238-804-5; ISBN 1-56238-805-3). **Clinical and Laboratory Standards Institute**, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087 USA, 2017.
- CLINICAL AND LABORATORY STANDARDS INSTITUTE (CLSI). Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals. 4th ed. CLSI supplement VET08 (ISBN 978-1-68440-010-2 [Print]; ISBN 978-1-68440-011-9 [Electronic]). **Clinical and Laboratory Standards Institute**, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087 USA, 2018.
- CLINICAL AND LABORATORY STANDARDS INSTITUTE (CLSI). Wayne, PA, USA. Performance Standards for Antimicrobial Susceptibility Testing. **30th ed. CLSI supplement M100, 19087**, 2020.
- CROXEN, M. A. et al. Recent advances in understanding enteric pathogenic *Escherichia coli*. **Clinical Microbiology Reviews**, v.26, p.822–880, 2013. Available from: <<https://pubmed.ncbi.nlm.nih.gov/24092857/>>. Accessed: Jun. 16, 2022. doi: 10.1128/CMR.00022-13.
- CUNHA, M. P. V. et al. CTX-M-producing *Escherichia coli* Isolated from urban pigeons (*Columba livia domestica*) in Brazil. **The Journal of Infection in Developing Countries**, v.13, p.1052–1056, 2019. Available from: <<https://pubmed.ncbi.nlm.nih.gov/32087078/>>. Accessed: Jun. 16, 2022. doi: 10.3855/jidc.11441.

- CZEKAJ, T. et al. *Staphylococcus haemolyticus* - an emerging threat in the twilight of the antibiotics age. **Microbiology (Reading, England)**, v.161, p.2061–2068, 2015. Available from: <<https://pubmed.ncbi.nlm.nih.gov/26363644/>>. Accessed: Jun. 16, 2022. doi: 10.1099/mic.0.000178.
- DOVC, A. et al. Health status of free-living pigeons (*Columba livia domestica*) in the city of Ljubljana. **Acta Veterinaria Hungarica**, v.52, p.219–226, 2004. Available from: <<https://pubmed.ncbi.nlm.nih.gov/15168753/>>. Accessed: Jun. 16, 2022. doi: 10.1556/AVet.52.2004.2.10.
- DUTTA, P. et al. Isolation of *Salmonella* Typhimurium from pigeons (*Columba livia*) in Greater Guwahati, its histopathological impact and antibiogram. **Comparative Clinical Pathology**, v.22, p.147–150, 2013. Available from: <<https://link.springer.com/article/10.1007/s00580-012-1614-3>>. Accessed: Jun. 16, 2022. doi: 10.1007/s00580-012-1614-3.
- EUCAST - The European Committee on Antimicrobial Susceptibility Testing (EUCAST). Breakpoint tables for interpretation of MICs and zone diameters. In: **European Society of Clinical Microbiology and Infectious Diseases Basel**, 2019.
- FERMAN, L. M. et al. A study of feral pigeon *Columba livia* var. in urban and suburban areas in the city of Jena, Germany. **Arxius de Miscel·lànea Zoològica**, 8, 1–8, 2010. Available from: <<http://www.bcn.cat/mciencies/publicacions/AMZ/vol8.1/AMZ-1001.htm>>. Accessed: Jun. 16, 2022.
- FOX, J. G.; GE et al. *Helicobacter hepaticus* infection in mice: models for understanding lower bowel inflammation and cancer. **Mucosal Immunology**, v.4, p.22–30, 2011. Available from: <<https://pubmed.ncbi.nlm.nih.gov/20944559/>>. Accessed: Jun. 16, 2022. doi: 10.1038/mi.2010.61.
- FRANCK, S. M. et al. Multiplex PCR for Enterotoxigenic, Attaching and Effacing, and Shiga Toxin-Producing *Escherichia coli* Strains from Calves. **Journal of Clinical Microbiology**, v.36, p.1795–1797, 1998. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC104926/>>. Accessed: Jun. 16, 2022. doi: 10.1128/jcm.36.6.1795-1797.1998.
- FROSINI, S. M. et al. A. Genes on the Move: In Vitro Transduction of Antimicrobial Resistance Genes between Human and Canine Staphylococcal Pathogens. **Microorganisms**, v. 8, p.2031, 2020. Available from: <<https://pubmed.ncbi.nlm.nih.gov/33353175/>>. Accessed: Jun. 16, 2022. doi: 10.3390/microorganisms8122031.
- FUTAGAWA-SAITO, K. et al. High occurrence of multi-antimicrobial resistance in *Staphylococcus intermedius* isolates from healthy and diseased dogs and domesticated pigeons. **Research in Veterinary Science**, v.83, p.336–339, 2007. Available from: <<https://pubmed.ncbi.nlm.nih.gov/17368493/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.rvsc.2007.01.013.
- GARGIULO, A. et al. Occurrence of Enteropathogenic Bacteria in Urban Pigeons (*Columba livia*) in Italy. **Vector-Borne and Zoonotic Diseases**, v.14, p.251–255, 2014. Available from: <<https://www.liebertpub.com/doi/10.1089/vbz.2011.0943>>. Accessed: Jun. 16, 2022. doi: 10.1089/vbz.2011.0943.
- GHANBARPOUR, R.; DANESHDOOST, S. Identification of shiga toxin and intimin coding genes in *Escherichia coli* isolates from pigeons (*Columba livia*) in relation to phylotypes and antibiotic resistance patterns. **Tropical Animal Health and Production**, v.44, p.307–312, 2012. Available from: <<https://pubmed.ncbi.nlm.nih.gov/22105907/>>. Accessed: Jun. 16, 2022. doi: 10.1007/s11250-011-0021-0.
- GORDON, D. M.; COWLING, A. The distribution and genetic structure of *Escherichia coli* in Australian vertebrates: host and geographic effects. **Microbiology (Reading, England)**, v.149, p.3575–3586, 2003. Available from: <<https://pubmed.ncbi.nlm.nih.gov/14663089/>>. Accessed: Jun. 16, 2022. doi: 10.1099/mic.0.26486-0.
- GRIMONT, P.; WEILL, F. X. Antigenic Formulae of the *Salmonella* serovars, (9th ed.) Paris: WHO Collaborating Centre for Reference and Research on *Salmonella*. **Institute Pasteur.**, p.1–166, 2007. Available from: <https://www.pasteur.fr/sites/default/files/veng_0.pdf>. Accessed: Jun. 16, 2022.
- HAABER, J. et al. Transfer of Antibiotic Resistance in *Staphylococcus aureus*. **Trends in Microbiology**, v.25, p.893–905, 2017. Available from: <<https://pubmed.ncbi.nlm.nih.gov/28641931/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.tim.2017.05.011.
- HAAG-WACKERNAGEL, D.; MOCH, H. Health hazards posed by feral pigeons. **The Journal of Infection**, v.48, p.307–313, 2004. Available from: <<https://pubmed.ncbi.nlm.nih.gov/15066331/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.jinf.2003.11.001.
- HAESSENDONCK, R. et al. Feral pigeons: A reservoir of zoonotic *Salmonella* Enteritidis strains? **Veterinary Microbiology**, v.195, p.101–103, 2016. Available from: <<https://pubmed.ncbi.nlm.nih.gov/27771054/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.vetmic.2016.09.017.
- HALE, C. R. et al. Estimates of enteric illness attributable to contact with animals and their environments in the United States. **Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America**, v.54 Suppl 5, p.S472–479, 2012. Available from: <<https://pubmed.ncbi.nlm.nih.gov/22572672/>>. Accessed: Jun. 16, 2022. doi: 10.1093/cid/cis051.
- HOELZER, K. et al. Animal contact as a source of human nontyphoidal salmonellosis. **Veterinary Research**, v.42, p.34, 2011. Available from: <<https://pubmed.ncbi.nlm.nih.gov/21324103/>>. Accessed: Jun. 16, 2022. doi: 10.1186/1297-9716-42-34.
- INMET - Instituto Nacional de Meteorologia. Available from: <<http://sisdagro.inmet.gov.br/sisdagro/app/monitoramento/bhs/mapaperiodoacum>>. Accessed: Jun. 16, 2022.
- INMET - Instituto Nacional de Meteorologia. Available from: <<http://sisdagro.inmet.gov.br/sisdagro/app/monitoramento/bhs/mapaperiodoacum>>. Accessed: Jun. 16, 2022.
- IPPOLITO, G. LE et al. Methicillin-resistant *Staphylococcus aureus*: the superbug. **International journal of infectious diseases: IJID: official publication of the International Society for Infectious Diseases**, v.14 Suppl 4, p.S7–11, 2010. Available from: <<https://pubmed.ncbi.nlm.nih.gov/20851011/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.ijid.2010.05.003.
- KACZOREK-LUKOWSKA, E. et al. Can domestic pigeon be a potential carrier of zoonotic *Salmonella*? **Transboundary and Emerging Diseases**, v.68, 2020. Available from: <<https://doi.org/10.1111/tbed.13891>>. Accessed: Jun. 16, 2022. doi: 10.1111/tbed.13891.

- KAMATHEWATTA, K. I. et al. Exploration of antibiotic resistance risks in a veterinary teaching hospital with Oxford Nanopore long read sequencing. **PLoS ONE**, v.14, p.e0217600, 2019. Available from: <<https://pubmed.ncbi.nlm.nih.gov/31145757/>>. Accessed: Jun. 16, 2022. doi: 10.1371/journal.pone.0217600.
- KARIM, S. J. I. et al. Multidrug-resistant *Escherichia coli* and *Salmonella* spp. isolated from pigeons. **Veterinary World**, v.13, p.2156–2165, 2020. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7704316/>>. Accessed: Jun. 16, 2022. doi: 10.14202/vetworld.2020.2156-2165.
- KIZERWETTER-ŚWIDA, M. et al. Genetic characterization of coagulase-positive staphylococci isolated from healthy pigeons. **Polish Journal of Veterinary Sciences**, v.18, p.627–634, 2015. Available from: <<https://pubmed.ncbi.nlm.nih.gov/26618597/>>. Accessed: Jun. 16, 2022. doi: 10.1515/pjvs-2015-0081.
- KNIGHT, D. R.; RILEY, T. V. Genomic delineation of zoonotic origins of *Clostridium difficile*. **Frontiers in Public Health**, v.7, p.164, 2019. Available from: <<https://pubmed.ncbi.nlm.nih.gov/31281807/>>. Accessed: Jun. 16, 2022. doi: 10.3389/fpubh.2019.00164.
- KOKSAL, F. et al. Antibiotic resistance patterns of coagulase-negative staphylococcus strains isolated from blood cultures of septicemic patients in Turkey. **Microbiological Research**, v.164, p.404–410, 2009. Available from: <<https://pubmed.ncbi.nlm.nih.gov/17475456/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.micres.2007.03.004.
- KUANG, X. et al. Serotypes and antimicrobial susceptibility of *Salmonella* spp. isolated from farm animals in China. **Frontiers in Microbiology**, v.6, p.602, 2015. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4476277/>>. Accessed: Jun. 16, 2022. doi: 10.3389/fmicb.2015.00602.
- KUTKOWSKA, J. et al. Methicillin-resistant *Staphylococcus aureus* and glycopeptide-resistant enterococci in fecal samples of birds from South-Eastern Poland. **BMC Veterinary Research**, v.15, p.472, 2019. Available from: <<https://pubmed.ncbi.nlm.nih.gov/31888629/>>. Accessed: Jun. 16, 2022. doi: 10.1186/s12917-019-2221-1.
- LE MINOR, L.; POPOFF, M. Y. Y. 1987. Designation of *Salmonella enterica* sp. nov., nom. rev., as the Type and Only Species of the Genus *Salmonella*: Request for an Opinion. **International Journal of Systematic and Evolutionary Microbiology**, v.37, p.465–468, [s.d.].
- MAGIORAKOS, A. P. et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. **Clinical Microbiology and Infection**, v.18, p.268–281, 2012. Available from: <<https://pubmed.ncbi.nlm.nih.gov/21793988/>>. Accessed: Jun. 16, 2022. doi: 10.1111/j.1469-0691.2011.03570.x.
- MATIAS, C. A. R. et al. *Staphylococcus* spp. isolated from wild birds apprehended in the local illegal trade in Rio de Janeiro, Brazil, and relevance in public health. **Letters in Applied Microbiology**, v.67, p.292–298, 2018. Available from: <<https://pubmed.ncbi.nlm.nih.gov/29924392/>>. Accessed: Jun. 16, 2022. doi: 10.1111/lam.13035.
- MCDANIELS, A. E. et al. Confirmational identification of *Escherichia coli*, a comparison of genotypic and phenotypic assays for glutamate decarboxylase and beta-D-glucuronidase. **Applied and Environmental Microbiology**, v.62, p.3350–3354, 1996. Available from: <<https://pubmed.ncbi.nlm.nih.gov/8795225/>>. Accessed: Jun. 16, 2022. doi: 10.1128/aem.62.9.3350-3354.1996.
- MURAKAMI, K. et al. Identification of methicillin-resistant strains of staphylococci by polymerase chain reaction. **Journal of Clinical Microbiology**, v.29, p.2240–2244, 1991. Available from: <<https://pubmed.ncbi.nlm.nih.gov/1939577/>>. Accessed: Jun. 16, 2022. doi: 10.1128/JCM.29.10.2240-2244.1991.
- MAHMMOUD, E. N. Isolation and Identification of Bacteria from Genital Organs of Pigeons. **Rafidain Journal of Science**, v.24, p.1–15, 2013. Available from: <https://rsci.mosuljournals.com/article_78799.html>. Accessed: Jun. 16, 2022. doi: 10.33899/rjs.2013.78799.
- OLIVEIRA, J. G. et al. *Salmonella* Typhimurium - associated meningoencephalomyelitis in a foal. **Ciência Rural**, v.49, 2019. Available from: <<https://doi.org/10.1590/0103-8478cr20190008>>. Accessed: Jun. 16, 2022. doi: 10.1590/0103-8478cr20190008.
- OSMAN, K. M. et al. *Salmonella enterica* isolated from pigeon (*Columba livia*) in Egypt. **Foodborne Pathogens and Disease**, v.10, p.481–483, 2013. Available from: <<https://pubmed.ncbi.nlm.nih.gov/23531124/>>. Accessed: Jun. 16, 2022. doi: 10.1089/fpd.2012.1347.
- PALMA, E.; TILOCCA, B.; RONCADA, P. Antimicrobial Resistance in Veterinary Medicine: An Overview. **International Journal of Molecular Sciences**, v.21, p.1914, 2020. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7139321/>>. Accessed: Jun. 16, 2022. doi: 10.3390/ijms21061914.
- PANGLOLI, P. et al. Seasonal incidence and molecular characterization of *Salmonella* from dairy cows, calves, and farm environment. **Foodborne Pathogens and Disease**, v.5, p.87–96, 2008. Available from: <<https://pubmed.ncbi.nlm.nih.gov/18260819/>>. Accessed: Jun. 16, 2022. doi: 10.1089/fpd.2008.0048.
- PEDERSEN, K. et al. Prevalence of shiga toxin-producing *Escherichia coli* and *Salmonella enterica* in rock pigeons captured in Fort Collins, Colorado. **Journal of Wildlife Diseases**, v.42, p.46–55, 2006. Available from: <<https://pubmed.ncbi.nlm.nih.gov/16699148/>>. Accessed: Jun. 16, 2022. doi: 10.7589/0090-3558-42.1.46.
- PIRŠ, T. et al. Antimicrobial susceptibility of animal and human isolates of *Clostridium difficile* by broth microdilution. **Journal of Medical Microbiology**, v.62, p.1478–1485, 2013. Available from: <<https://pubmed.ncbi.nlm.nih.gov/23861298/>>. Accessed: Jun. 16, 2022. doi: 10.1099/jmm.0.058875-0.
- PITCHER, D. G. et al. Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. **Letters in Applied Microbiology**, v.8, p.151–156, 1989. Available from: <<https://doi.org/10.1111/j.1472-765X.1989.tb00262.x>>. Accessed: Jun. 16, 2022. doi: 10.1111/j.1472-765X.1989.tb00262.x.
- PURIGHALLA, S. et al. Investigation into a community outbreak of *Salmonella* Typhi in Bengaluru, India. **The Indian Journal of Medical Research**, v.146, p.S15–S22, 2017. Available from: <<https://pubmed.ncbi.nlm.nih.gov/29205191/>>. Accessed: Jun. 16, 2022. doi: 10.4103/ijmr.IJMR_1201_16.
- RAMOS, C. P. et al. Enteric Organisms Detected in Feces of Dogs with Bloody Diarrhea: 45 Cases. **Topics in Companion Animal**

- Medicine**, v.45, p.100549, 2021. Available from: <<https://pubmed.ncbi.nlm.nih.gov/34044173/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.tcam.2021.100549.
- RAMOS, C. P. et al. Identification and Characterization of *Escherichia coli*, *Salmonella* Spp., *Clostridium perfringens*, and *C. difficile* Isolates from Reptiles in Brazil. **BioMed Research International**, v.2019, p.1–9, 2019a. Available from: <<https://pubmed.ncbi.nlm.nih.gov/31263711/>>. Accessed: Jun. 16, 2022. doi: 10.1155/2019/9530732.
- RAMOS, C. P. et al. Outbreak of multidrug-resistant Salmonella Typhimurium in calves at a veterinary hospital in Brazil. **Ciência Rural**, v.49, 2019b. Available from: <<https://doi.org/10.1590/0103-8478cr20180788>>. Accessed: Jun. 16, 2022. doi: 10.1590/0103-8478cr20180788.
- RISSI, D. R.; ELSMO, E. J.; SANCHEZ, S. **Cystitis and peritonitis caused by *Staphylococcus xylosus* infection in a calf**. v.8, p.99–101, 2015. Available from: <<https://pesquisa.bvsalud.org/portal/resource/pt/vti-22003>>. Accessed: Jun. 16, 2022.
- ROCHA-E-SILVA, R. C. et al. O pombo (*Columba livia*) como agente carreador de *Salmonella* spp. e as implicações em saúde pública. **Arquivos do Instituto Biológico**, v.81, p.189–194, 2014. Available from: <<https://doi.org/10.1590/1808-1657000702012>>. Accessed: Jun. 16, 2022. doi: 10.1590/1808-1657000702012.
- ROSARIO MEDINA, I. et al. Pigeons and their droppings as reservoirs of *Candida* and other zoonotic yeasts. **Revista Iberoamericana de Micología**, v.34, p.211–214, 2017. Available from: <<https://linkinghub.elsevier.com/retrieve/pii/S1130140617300529>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.riam.2017.03.001.
- SACRISTÁN, C. et al. Virulence genes, antibiotic resistance and integrons in *Escherichia coli* strains isolated from synanthropic birds from Spain. **Avian Pathology: Journal of the W.V.P.A.**, v.43, p.172–175, 2014. Available from: <<https://pubmed.ncbi.nlm.nih.gov/24689431/>>. Accessed: Jun. 16, 2022. doi:10.1080/03079457.2014.897683.
- SANTANA, J. A. et al. *Clostridioides difficile* and multi-drug-resistant staphylococci in free-living rodents and marsupials in parks of Belo Horizonte, Brazil. **Brazilian Journal of Microbiology**, 2021. Available from: <<https://pubmed.ncbi.nlm.nih.gov/34761356/>>. Accessed: Jun. 16, 2022. doi: 10.1007/s42770-021-00640-x.
- SANTANA, J. A. et al. Isolation and antimicrobial resistance of coagulase-negative staphylococci recovered from healthy tortoises in Minas Gerais, Brazil. **Ciência Rural**, v.52, p.e20210354, 2022. Available from: <<https://doi.org/10.1590/0103-8478cr20210354>>. Accessed: Jun. 16, 2022. doi: 10.1590/0103-8478cr20210354.
- SASAKI, T. et al. Multiplex-PCR method for species identification of coagulase-positive staphylococci. **Journal of Clinical Microbiology**, v.48, p.765–769, 2010. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2832457/>>. Accessed: Jun. 16, 2022. doi: 10.1128/JCM.01232-09.
- SAWA, H.; HIRAI, K. An outbreak of *Salmonella* Typhimurium subserovar copenhagen infection in pigeons (*Chalcophaps indica*) imported from Hong Kong. **Nihon Juigaku Zasshi. The Japanese Journal of Veterinary Science**, v.43, p.277–279, 1981. Available from: <<https://agris.fao.org/agris-search/search.do?recordID=JP19820771480>>. Accessed: Jun. 16, 2022. doi: 10.1292/jvms.1939.43.277.
- SCHWARZ, S.; WERCKENTHIN, C. Antibiotic Resistance in Staphylococci Isolated from Pigeons. **Veterinary Dermatology**, v.5, p.9–12, 1994. Available from: <<https://doi.org/10.1111/j.1365-3164.1994.tb00003.x>>. Accessed: Jun. 16, 2022. doi: 10.1111/j.1365-3164.1994.tb00003.x.
- SIDHU, M. S. et al. Persistence of multidrug-resistant Staphylococcus haemolyticus in an animal veterinary teaching hospital clinic. **Microbial Drug Resistance (Larchmont, N.Y.)**, v.13, p.271–280, 2007. Available from: <<https://pubmed.ncbi.nlm.nih.gov/18184053/>>. Accessed: Jun. 16, 2022. doi: 10.1089/mdr.2007.756.
- SILVA, R. O. S. et al. Detection of A/B toxin and isolation of *Clostridium difficile* and *Clostridium perfringens* from foals. **Equine Veterinary Journal**, v.45, p.671–675, 2013. Available from: <<https://pubmed.ncbi.nlm.nih.gov/23452044/>>. Accessed: Jun. 16, 2022. doi: 10.1111/evj.12046.
- SILVA, R. O. S. et al. Detection of enterotoxin A and cytotoxin B, and isolation of *Clostridium difficile* in piglets in Minas Gerais, Brazil. **Ciência Rural**, v.41, p.1430–1435, 2011. Available from: <<https://doi.org/10.1590/S0103-84782011005000100>>. Accessed: Jun. 16, 2022. doi: 10.1590/S0103-84782011005000100.
- SILVA, V. L. et al. Diarrheagenic *Escherichia coli* strains recovered from urban pigeons (*Columba livia*) in Brazil and their antimicrobial susceptibility patterns. **Current Microbiology**, v.59, p.302–308, 2009. Available from: <<https://pubmed.ncbi.nlm.nih.gov/19504156/>>. Accessed: Jun. 16, 2022. doi: 10.1007/s00284-009-9434-7.
- SPENNEMANN, D. H.; WATSON, M. J. Dietary habits of urban pigeons (*Columba livia*) and implications of excreta pH – A review. **European Journal of Ecology**, v.3, p.27–41, 2017. Available from: <<https://journals.ku.edu/EuroJecol/article/view/11598>>. Accessed: Jun. 16, 2022. doi: 10.1515/eje-2017-0004.
- SUKKUA, K.; MANOTHONG, S.; SUKHUMUNGOON, P. Seroprevalence and molecular epidemiology of EAST1 gene-carrying *Escherichia coli* from diarrheal patients and raw meats. **Journal of Infection in Developing Countries**, v.11, p.220–227, 2017. Available from: <<https://pubmed.ncbi.nlm.nih.gov/28368855/>>. Accessed: Jun. 16, 2022. doi: 10.3855/jidc.6865.
- SWEENEY, M. T. et al. Applying definitions for multidrug resistance, extensive drug resistance and pandrug resistance to clinically significant livestock and companion animal bacterial pathogens. **Journal of Antimicrobial Chemotherapy**, v.73, p.1460–1463, 2018. Available from: <<https://pubmed.ncbi.nlm.nih.gov/29481657/>>. Accessed: Jun. 16, 2022. doi: 10.1093/jac/dky043.
- TANAKA, C. et al. Bacteriological survey of feces from feral pigeons in Japan. **The Journal of Veterinary Medical Science**, v.67, p.951–953, 2005. Available from: <https://www.jstage.jst.go.jp/article/jvms/67/9/67_9_951/article>. Accessed: Jun. 16, 2022. doi: 10.1292/jvms.67.951.
- TIZARD, I. Salmonellosis in wild birds. **Seminars in Avian and Exotic Pet Medicine**, Emerging Diseases. v.13, p.50–66, 2004. Available from: <<https://www.sciencedirect.com/science/article/abs/pii/S1055937X04000039>>. Accessed: Jun. 16, 2022. doi: 10.1053/j.saep.2004.01.008.

- TORRES-MEJÍA, A. M. et al. Zoonotic Agents in Feral Pigeons (*Columba livia*) from Costa Rica: Possible Improvements to Diminish Contagion Risks. **Vector-Borne and Zoonotic Diseases**, v.18, p.49–54, 2018. Available from: <<https://pubmed.ncbi.nlm.nih.gov/29243991/>>. Accessed: Jun. 16, 2022. doi: 10.1089/vbz.2017.2131.
- TURNER, N. A. et al. Methicillin-resistant *Staphylococcus aureus*: an overview of basic and clinical research. **Nature Reviews. Microbiology**, v.17, p.203–218, 2019. Available from: <<https://pubmed.ncbi.nlm.nih.gov/30737488/>>. Accessed: Jun. 16, 2022. doi: 10.1038/s41579-018-0147-4.
- VELA, J. et al. Characterization of *Staphylococcus xylosum* isolated from broiler chicken barn bioaerosol. **Poultry Science**, v.91, p.3003–3012, 2012. Available from: <<https://pubmed.ncbi.nlm.nih.gov/23155006/>>. Accessed: Jun. 16, 2022. doi: 10.3382/ps.2012-02302.
- VERSALOVIC, J.; KOEUTH, T.; LUPSKI, J.R. Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. **Nucleic Acids Research**, v.19, p.6823–6831, 1991. Available from: <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC329316/>>. Accessed: Jun. 16, 2022. doi: 10.1093/nar/19.24.6823.
- WALTHER B, TEDIN K, LÜBKE-BECKER A. Multidrug-resistant opportunistic pathogens challenging veterinary infection control. **Vet Microbiol.** 2017 Feb; 200:71-78. Available from: <<https://pubmed.ncbi.nlm.nih.gov/27291944/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.vetmic.2016.05.017.
- WEESE, J. S.; DUIJKEREN, E. VAN. Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine. **Veterinary Microbiology**, v.140, p.418–429, 2010. Available from: <<https://pubmed.ncbi.nlm.nih.gov/19246166/>>. Accessed: Jun. 16, 2022. doi: 10.1016/j.vetmic.2009.01.039.
- WEESE JS, ARMSTRONG J. Outbreak of *Clostridium difficile*-associated disease in a small animal veterinary teaching hospital. **J Vet Intern Med.** 2003 Nov-Dec; 17(6):813-6. Available from: <<https://pubmed.ncbi.nlm.nih.gov/14658717/>>. Accessed: Jun. 16, 2022. doi: 10.1111/j.1939-1676.2003.tb02519.x.
- WON, Y. S. et al. Identification of *Staphylococcus xylosum* isolated from C57BL/6J-Nos2(tm1Lau) mice with dermatitis. **Microbiology and Immunology**, v.46, p.629–632, 2002. Available from: <<https://pubmed.ncbi.nlm.nih.gov/12437030/>>. Accessed: Jun. 16, 2022. doi: 10.1111/j.1348-0421.2002.tb02744.x.
- WORLD HEALTH ORGANIZATION (WHO). Critically important antimicrobials for human medicine, 2019. Available from: <<https://www.who.int/publications/i/item/9789241515528>>. Accessed: Jun. 16, 2022.
- YAMAMOTO, T.; NAKAZAWA, M. Detection and sequences of the enteroaggregative *Escherichia coli* heat-stable enterotoxin 1 gene in enterotoxigenic *E. coli* strains isolated from piglets and calves with diarrhea. **Journal of Clinical Microbiology**, v.35, p.223–227, 1997. Available from: <<https://pubmed.ncbi.nlm.nih.gov/8968912/>>. Accessed: Jun. 16, 2022. doi: 10.1128/jcm.35.1.223-227.1997.
- ZHOU, Z. et al. An outbreak of gastroenteritis in Osaka, Japan due to *Escherichia coli* serogroup O166:H15 that had a coding gene for enteroaggregative *E. coli* heat-stable enterotoxin 1 (EAST1). **Epidemiology and Infection**, v.128, p.363–371, 2002. Available from: <<https://pubmed.ncbi.nlm.nih.gov/12113479/>>. Accessed: Jun. 16, 2022. doi: 10.1017/s0950268802006994.
- ZIGO, F. Changes in Bacterial Microflora in Young Carrier Pigeons during the Race Season. **International International Journal of Avian & Wildlife Biology**, v.2, 2017. Available from: <<https://medcraveonline.com/IJAWB/changes-in-bacterial-microflora-in-young-carrier-pigeons-during-the-race-season.html>>. Accessed: Jun. 16, 2022. doi: 10.15406/ijawb.2017.02.00013.