



OPEN ACCESS

EDITED BY

William Dowling,
Coalition for Epidemic Preparedness
Innovations (CEPI), Norway

REVIEWED BY

Henning Jacobsen,
Helmholtz Centre for Infection Research,
Germany
Diego Cantoni,
MRC-University of Glasgow Centre For
Virus Research (MRC), United Kingdom

*CORRESPONDENCE

Cristina Toscano Fonseca
✉ cristina.toscano@fiocruz.br

RECEIVED 10 May 2023

ACCEPTED 26 June 2023

PUBLISHED 14 July 2023

CITATION

Franco-Luiz APM, Fernandes NMGS,
Silva TBdS, Bernardes WPOS, Westin MR,
Santos TG, Fernandes GdR, Simões TC,
Silva EFE, Gava SG, Alves BM,
de Carvalho Melo M, da Silva-Pereira RA,
Alves PA and Fonseca CT (2023)
Longitudinal study of humoral immunity
against SARS-CoV-2 of health
professionals in Brazil: the impact of
booster dose and reinfection on
antibody dynamics.
Front. Immunol. 14:1220600.
doi: 10.3389/fimmu.2023.1220600

COPYRIGHT

© 2023 Franco-Luiz, Fernandes, Silva,
Bernardes, Westin, Santos, Fernandes,
Simões, Silva, Gava, Alves, de Carvalho Melo,
da Silva-Pereira, Alves and Fonseca. This is
an open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Longitudinal study of humoral immunity against SARS-CoV-2 of health professionals in Brazil: the impact of booster dose and reinfection on antibody dynamics

Ana Paula Moreira Franco-Luiz¹,
Nubia Monteiro Gonçalves Soares Fernandes¹,
Thais Bárbara de Souza Silva²,
Wilma Patrícia de Oliveira Santos Bernardes¹,
Mateus Rodrigues Westin³, Thais Garcia Santos¹,
Gabriel da Rocha Fernandes⁴, Taynãna César Simões⁵,
Eduardo Fernandes E. Silva⁶, Sandra Grossi Gava⁷,
Breno Magalhães Alves⁸, Mariana de Carvalho Melo⁹,
Rosiane A. da Silva-Pereira¹, Pedro Augusto Alves²
and Cristina Toscano Fonseca^{1*}

¹Grupo de Pesquisa em Biologia e Imunologia de Doenças Infecciosas e Parasitárias, Instituto René Rachou, Fundação Oswaldo Cruz, Belo Horizonte, Minas Gerais, Brazil, ²Grupo de Imunologia de Doenças Virais, Instituto René Rachou, Fundação Oswaldo Cruz, Belo Horizonte, Minas Gerais, Brazil, ³Faculdade de Medicina, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, ⁴Grupo de Pesquisa em Informática de Biosistemas, Instituto René Rachou, Fundação Oswaldo Cruz, Belo Horizonte, Minas Gerais, Brazil, ⁵Núcleo de Estudos em Saúde Pública e Envelhecimento, Instituto René Rachou, Fundação Oswaldo Cruz, Belo Horizonte, Minas Gerais, Brazil, ⁶Serviço de capacitação em métodos quantitativos -SAmEq, Instituto René Rachou, Fundação Oswaldo Cruz, Belo Horizonte, Minas Gerais, Brazil, ⁷Grupo de Pesquisa em Helminologia e Malacologia Médica, Instituto René Rachou, Fundação Oswaldo Cruz, Belo Horizonte, Minas Gerais, Brazil, ⁸Centro de Vigilância em Saúde e Segurança do Paciente, Hospital Metropolitano Doutor Célio de Castro, Belo Horizonte, Minas Gerais, Brazil, ⁹Serviço Especializado em Segurança e Medicina do Trabalho, Hospital Metropolitano Doutor Célio de Castro, Belo Horizonte, Minas Gerais, Brazil

Introduction: The pandemic caused by SARS-CoV-2 has had a major impact on health systems. Vaccines have been shown to be effective in improving the clinical outcome of COVID-19, but they are not able to fully prevent infection and reinfection, especially that caused by new variants.

Methods: Here, we tracked for 450 days the humoral immune response and reinfection in 52 healthcare workers from Brazil. Infection and reinfection were confirmed by RT-qPCR, while IgM and IgG antibody levels were monitored by rapid test.

Results: Of the 52 participants, 19 (36%) got reinfected during the follow-up period, all presenting mild symptoms. For all participants, IgM levels dropped sharply, with over 47% of them becoming seronegative by the 60th day. For IgG, 90% of the participants became seropositive within the first 30 days of follow-up.

IgG antibodies also dropped after this period reaching the lowest level on day 270 (68.5 ± 72.3 , $p < 0.0001$). Booster dose and reinfection increased the levels of both antibodies, with the interaction between them resulting in an increase in IgG levels of 130.3 arbitrary units.

Conclusions: Overall, our data indicate that acquired humoral immunity declines over time and suggests that IgM and IgG antibody levels are not associated with the prevention of reinfection.

KEYWORDS

antibody, COVID-19, healthcare workers, humoral immunity, reinfection, SARS-COV-2

1 Introduction

Coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), emerged in China in 2019, and has affected more than 200 countries. By the end of 2022, there had been 647,972,911 confirmed cases of COVID-19, including 6,642,832 deaths, in the world. In Brazil, the first case was diagnosed in February 2020, and by the end of December 2022, the country had had more than 35 million confirmed cases (1). Healthcare workers (HCW) were a group greatly impacted by the pandemic, leading COVID-19 to be recognized as an occupational disease (2).

Due to the significant impact of the pandemic on health systems, a global effort has sought various alternatives to reduce the harm caused by the disease (3). Thus, vaccines have been developed and approved (4). Due to their higher risk of exposure, HCW were the first group to be vaccinated (5). In January 2021, the vaccination schedule began in Brazil. Until December 2022, more than 80% of the population have had a complete vaccination schedule, and almost 50% have had a booster dose (6). Although these vaccines reduce virus levels in the body, and consequently reduce viral transmission, they do not fully prevent new SARS-CoV-2 infections (7–9). Vaccines have proven to be safe, effective, and timely tools to prevent severe outcomes of COVID-19, including hospitalization and death. However, the efficiency of vaccination can change depending on the type of vaccine and other factors, such as the emergence and/or introduction of new viral variants (9–11).

Since the first recorded cases of COVID-19, new variants of SARS-CoV-2 have been identified. Therefore, in order to establish control and monitoring goals for the new variants, the World Health Organization (WHO) has established three classification categories: variants of concern (VOCs), variants of interest (VOIs), and variants under monitoring (VUMs). The four previously circulating VOCs are Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), and Delta (B.1.617.2), while Omicron (B.1.1.529, including BA.1, BA.2, BA.3, BA.4, BA.5 and their descendent lineages) are currently circulating (1). Each SARS-CoV-2 VOC is associated with a new wave of infection, such as the Gamma variant in December 2020 and the Omicron variant in December 2021, which affect

human health worldwide (1, 12). Studies involving these new variants have elucidated fundamental aspects of SARS-CoV-2 biology, including viral transmissibility, disease severity, immune system escape, vaccine efficiency, clinical treatment, and management strategies (3, 13, 14).

It is already known that it is possible to become reinfected with SARS-CoV-2 (12, 15). However, since some people have a recurrence of positive test results for SARS-CoV-2 RNA detection during apparently the same infection (16), in order to be considered a reinfection, the CDC has established that the nucleotide sequences of positive samples must be from different lineages or there must be a difference of 90 or more days between positive results (17). Previously published data on the prevalence of SARS-CoV-2 reinfection have highlighted its low level, ranging from 0.10 to 0.65% in the range of 6–9 months (18–20). However, the reinfection rate in Brazil is still unknown, especially in risk groups such as health professionals. The emergence of new variants could increase the reinfection rate, as they can escape the immune response triggered by existing vaccines, as with VOCs (e.g., Omicron) (14, 21, 22). Identifying the potential for SARS-CoV-2 reinfection is crucial to understanding the long-term dynamics of the pandemic. Previous studies suggest that the presence of IgG antibodies reduce the risk of reinfection (23). According to a study carried out in England, a primary infection reduces the reinfection rate by 84% over the following seven months (24). Therefore, understanding the dynamic behavior of SARS-CoV-2 infection, and assessing reinfection rates, the impact of genetic variants and vaccines on immune memory kinetics, and their application in the global vaccination campaign are some key points that still need to be elucidated (25).

In this study, we investigated, in HCW in the city of Belo Horizonte in Brazil, the dynamics and longevity of the humoral immune response up to 450 days after the initial onset of COVID-19 disease symptoms and laboratory confirmation of SARS-CoV-2 infection. During the study, the kinetics of the humoral response were monitored both before and after the initial vaccination scheme and subsequent booster dose. Our study demonstrated the occurrence of reinfection in 19 study participants, and showed that (i) the humoral immunity of HCW declined over time, and (ii) the booster dose was essential to increase antibody levels, mainly IgG, but not enough to protect against

reinfection with new variants. Robust and constant surveillance is, therefore, essential for responding to future epidemic waves, and provides a basis for recommendations for immunization programs and vaccine updates.

2 Materials and methods

2.1 Ethical approval

This study was conducted in accordance with current legislation including the Declaration of Helsinki and Resolution No. 466/2012 of the Conselho Nacional de Saúde do Brasil. Ethical approval was obtained from the institutional review board of the Instituto René Rachou, Fundação Oswaldo Cruz, CAAE: 31.919520.8.0000.5091, approval numbers: 4177931; 4291836; 4343318; 4624187; 5294423. Written informed consent was obtained from all participants before any study procedure was undertaken.

2.2 Study population and enrollment

The population selected for this study was composed of HCW who worked in at least one of the public hospitals: Hospital das Clínicas (HC) at the Universidade Federal de Minas Gerais, the Unidade de Pronto Atendimento (UPA) Centro-Sul, and the Hospital Metropolitano Dr. Célio de Castro (HMDCC). All three health centers are in the city of Belo Horizonte, the state of Minas Gerais, Brazil. As inclusion criteria, in addition to what has already been mentioned above, all participants had to (i) present with at least one of the following symptoms within the previous seven days: fever (equal to or greater than 37.5°C), cough (dry or productive), fatigue, dyspnea, sore throat, anosmia/hyposmia, and/or ageusia; and (ii) have a positive result for the detection of SARS-CoV-2 RNA by RT-qPCR. Participants were invited by telephone, and those who met the inclusion criteria were included in the study. Individuals who reported volunteering in COVID-19 vaccine clinical trials, prior diagnosis of COVID-19, or reported pregnancy, were

excluded. Hospitalization also resulted in loss of follow-up due to inability to perform the tests. The enrollment of participants took place between October 2020 to April 2021. Individuals were followed up for 450 days, with capillary blood samples collected on days 7, 10, 15, 20, 25, 30, 60, 90, 180, 270, and 450 after being enrolled in the study (referred to as D7, D10, D15 and so on). In addition to capillary blood collections, participants were contacted every 30 days after the 60th day (from December 2020 to June 2022) to assess possible SARS-CoV-2 reinfection. Besides, all participants were instructed to contact the research team when they present any symptoms that constitute suspicion of COVID-19. The predetermined symptoms which were monitored as evidence for reinfection were: fever (equal to or greater than 37.5°C), cough (dry or productive), sore throat, fatigue, dyspnea, and diarrhea. Also, any suspicious symptom established by medical criteria was monitored. All suspected cases were tested for the presence of viral RNA by RT-qPCR and serological status. In January 2021, during the study period, public roll-out of the vaccination scheme using the CoronaVac vaccine was started in Brazil. Fifteen days after the second dose, the initial two-dose vaccination scheme is considered “complete”. However, in September 2021, a booster dose was rolled-out (6). The workflow for our study is shown in Figure 1.

2.3 Study design

All symptomatic healthcare professionals were tested for SARS-CoV-2 by RT-qPCR using the Charite Institute protocol (26). During the isolation period, kits were sent to participants to perform a self-collection of saliva or nasopharyngeal and oropharyngeal swabs on D1, D3, and D5 after enrollment in the study. Swab or saliva was placed in a tube containing a lysis buffer with guanidine (4 M Guanidine Isothiocyanate, 25 mM Sodium Citrate, pH 7.4) that were stored at 4°C. Samples were collected in participants’ houses within a 5-day interval and were transported under refrigeration to the central laboratory for RNA extraction. As demonstrated by Carvalho et al., 2021 (27) this buffer, besides decreasing the virus infectivity, maintains RNA integrity for up to 16 days at room




FIGURE 1
Flowchart illustrating our study workflow. The numbers inside the squares represent the days of visits on which patients were followed-up. The icons shown in the legend represent the study design. After 60 days, questionnaires were answered to assess symptoms consistent with suspected SARS-CoV-2 reinfection, if reported by participants, and clinical samples were collected. Fifty-four participants were confirmed positive for COVID-19, of which one participant became pregnant and another was hospitalized, both were subsequently excluded from the study.

temperature. In addition to the material for saliva/swab sample collection, the kits included materials, together with an instruction manual, for performing a rapid test to detect IgM and IgG antibodies against SARS-CoV-2 antigens (TR DPP[®] COVID-19 IgM/IgG - Bio-Manguinhos). Participants characterized as infected with SARS-CoV-2 must have had at least one positive result for detection of SARS-CoV-2 RNA on D1, D3 and/or D5. Participants characterized as non-infected presented negative results in all three samples analyzed and were not included in the follow-up period of the study.

2.4 RNA extraction

RNA was extracted from the nasopharyngeal and oropharyngeal swabs or saliva samples using the QIAamp[®] Viral RNA kit (QIAGEN[®], USA) according to the manufacturer's instructions. Briefly, 140 μ L of sample was added to 560 μ L of AVL buffer containing carrier RNA, and after 10 minutes at room temperature, 560 μ L ethanol were added. This solution was applied to an RNA affinity column and this column was centrifuged at 6,000 x g for 1 minute. Then, the column was washed with AW1 and AW2 buffer solutions in that order. After the washing process, RNA was eluted using AVE solution and used in the RT-qPCR test.

2.5 RT-qPCR (Quantitative PCR)

The RT-qPCR reactions were performed using the ViiA[™] 7 Real-Time PCR System of the communal Real-Time PCR platform at the Instituto René Rachou. The RT-qPCR assays were performed using 5 μ L of sample RNA, and the 200 nM GoTaq[®] Probe 1-Step RT-qPCR System Kit (Promega). This kit uses GoTaq Probe qPCR Master Mix with dUTP (10 μ L), GoScript RT Mix for one-step RT-qPCR (0.4 μ L), sense and antisense primers (400 nM) and nuclease-free water to a 20.0 μ L final volume. The conditions for the amplification were: 45°C for 15 minutes and 95°C for 2 minutes, followed by 40 cycles of denaturation at 95°C for 15 seconds and hybridization at 60°C for 1 minute. The results were analyzed using the Thermo Cloud platform, according to the following criteria: samples with amplification of the E gene (Ct<37) and the RNase P gene (RP) (Ct<35) were considered positive; samples without E gene amplification or with detection above Ct 37, with RP amplification (Ct<35), were considered negative. Samples with RP amplification above Ct 35 were considered invalid, and the test was performed again using RNA obtained from another extraction of the samples collected.

2.6 TR DPP[®] COVID-19 IgM/IgG test (Bio-Manguinhos)

In order to detect IgM and IgG antibodies against SARS-CoV-2, the DPP[®] COVID-19 IgM/IgG kit supplied by Bio-Manguinhos (FIOCRUZ, Brazil) was used according to the manufacturer's instructions. Briefly, a digital puncture was performed, and a blood sample was diluted in the buffer provided in the kit. The sample was then applied to the cassette. After 5 minutes, 9 drops of the buffer were added to the cassette, and the results were read after an additional ten-minute period. The interpretation of the test was performed with the aid of the DPP[®] Micro Reader, which

provides the intensity of the reactive line. The DPP[®] COVID-19 IgM/IgG system uses an algorithm that includes assay-specific cutoff values to determine test result. Values equal to or greater than 30 for the IgM and IgG antibodies were considered reactive. In order to assess antibody levels, the test was performed on scheduled days and/or when the participant was suspected of reinfection (Figure 1).

2.7 Next-generation sequencing

Positive samples, with cycle threshold (Ct) lower than 36, were sequenced by Next-Generation Sequencing (NGS) on the Illumina MiSeq Platform using the Illumina COVIDSeq Kit for library construction (Illumina, San Diego, USA) generating paired-end reads 150 bp long. The raw reads were trimmed using Trimmomatic version 0.39 (28) with a sliding window of 4 nucleotides with a minimum average Phred score of 20. Trimmed reads smaller than 50 bp were removed. The filtered reads were mapped to the SARS-CoV-2 reference genome (NC_045512) using BWA version 0.7.17 (Li 2009) with the default parameters. The nucleotide variants were identified using iVar version 1.3 (29), with a minimum frequency of 40% and depth of 30 reads. The consensus sequences generated by iVar were submitted to Pangolin version 4.2 (30) to identify the coronavirus lineage. Sequences that met the GISAID criteria were submitted to the EpiCoV database and are deposited at The European Nucleotide Archive under project number PRJEB49204 (<https://www.ebi.ac.uk/ena/browser/view/PRJEB49204>).

2.8 Statistical analysis

GraphPad Prism v.8.0.1., Jamovi 2.3.18.0 (<https://www.jamovi.org/>), and the statistical software R (<https://www.r-project.org/>) version 4.1.2 (31) were used for data analysis and generation of figures. Data organization and pre-processing of some graphs and figures were done using Excel and PowerPoint (Microsoft 365). The chi-square test or the Fisher's exact test were used to assess the association of categorical demographic variables and the infection status of individuals, as well as the correlation of symptoms reported during the first infection, reinfection and in suspected cases of reinfection. Paired and unpaired t-tests were used to analyze the difference in viral load between infection vs reinfection, and females vs males, respectively.

Generalized Estimating Equations (GEE) (version 1.3.9) were used to evaluate the longitudinal data. Proposed by Lian and Zeger (1986) (32), the proposed model jointly estimates an average effect and intra-individual variations, considering the structure of correlation or dependence between the repeated measures. The outcome of interest was the numeric scale of IgM and IgG levels and the covariates were vaccine status, follow-up time, booster dose and reinfection. In the GEE model, the peak of IgM and IgG levels in the interval from D7 to D30 was considered. A model segmented in time, 180 and 270 for IgM and IgG, respectively, was adjusted to make the average structure more accurate.

In order to assess whether there is an association between antibody levels and reinfection, differences in the levels of IgM and IgG antibodies against SARS-CoV-2 among individuals with a

confirmed case of reinfection and those not reinfected were evaluated. This analysis only considered samples taken during the period of the study when reinfection occurred (i.e., between July 2021 and July 2022), which was also the period when the delta and omicron variants circulated in Belo Horizonte. For this evaluation, the Mann-Whitney U-test was used. For individuals who did not become reinfected, the arithmetic means of all of the values of the antibody levels determined during the period stated above were considered. For individuals who became reinfected, we considered only the values of the antibody levels determined when reinfection was confirmed.

3 Results

3.1 Description of the study cohort

A total of 163 health professionals aged between 19 to 68 years were included in the initial study cohort between October 2020 to April 2021 (Figure 1). Fifty-four (33%) of these individuals subsequently had positive test results by RT-qPCR. Two individuals were excluded from further involvement in the study due to pregnancy or hospitalization, while two other participants withdrew from the study during the follow-up period, but allowed the use of their data and the samples already collected. Therefore, our final study cohort comprised 52 individuals, with 50 of these participants remaining until the end of the 450-day follow-up period. Among the starting 52 participants, four (7.7%) worked at UPA, 16 (30.8%) at HMDCC, and 32 (61.5%) at HC. Overall, the average age of the subjects was 37.38 ± 6.99 years, and 55.8% were female. The final study cohort consisted of 22 (42.3%) physicians, 14 (26.9%) nursing technicians, 10 (19.2%) nurses, and 4 (7.7%) physiotherapists (Table 1).

In January 2021, the vaccination schedule started in Brazil. Thus, 30 (58%) participants were infected with SARS-CoV-2 before starting the vaccination schedule, 11 (21%) during the schedule interval, and 11 (21%) after the complete vaccination, that is 15 days after the second dose of the initial vaccination scheme. During the 450 days of follow-up, 46 (88%) participants had suspected reinfections; of these, 19 (37%) cases were confirmed by RT-qPCR (Figure 2). The proportion of women who became reinfected was higher than that of men ($p=0.011$). Among the participants who became reinfected 58% had already received the booster vaccine dose when the reinfection was confirmed.

3.2 IgM and IgG antibodies against SARS-CoV-2

The antibody peak observed between the seventh and thirtieth day (D7-D30) of each participant was chosen to be included in the analysis and represented a reaction average of $72.6 (\pm 68.0)$ for IgM (Figure 3A) and $193.7 (\pm 121.2)$ for IgG (Figure 3B). Nine (17.3%) of the 52 participants did not IgG seroconvert within 30 days, and two (3.8%) individuals did not seroconvert at any time during the study to either of the two monitored immunoglobulins. In addition to

TABLE 1 Demographic characteristics of the final study cohort (n = 52).

Characteristics	Infection	Reinfection	Total	p-value
Health lefts				
HC	20 (38.5%)	12 (23.1%)	32 (61.5%)	1.000
HMDCC	10 (19.2%)	6 (11.5%)	16 (30.8%)	
UPA	3 (5.8%)	1 (1.9%)	4 (7.7%)	
Total	33 (63.5%)	19 (36.5%)	52 (100%)	
Age				
Range 21-30	7 (13.5%)	5 (9.6%)	12 (23.1%)	0.084
Range 31-40	7 (13.5%)	9 (17.3%)	16(30.8%)	
Range 41-60	19 (36.5%)	5 (9.6%)	24(46.2%)	
Total	33 (63.5%)	19 (36.5%)	52(100%)	
Sex				
Female	14 (26.9%)	15 (28.8%)	29 (55.8%)	0.011
Male	19 (36.5%)	4 (7.7%)	23 (44.2%)	
Total	33 (63.5%)	19 (36.5%)	52 (100%)	
Professional Category				
Physicians	15 (28.8%)	7 (13.5%)	22 (42.3%)	0.691
Nurse Technician	7 (13.5%)	7 (13.5%)	14 (26.9%)	
Nurse	6 (11.5%)	4 (7.7%)	10 (19.2%)	
Physiotherapist	3 (5.8%)	1 (1.9%)	4(7.7%)	
¹ Others	2 (3.8%)	0 (0.0%)	2 (3.8%)	
Total	33(63.5%)	19 (36.5%)	52 (100%)	
Complete vaccination scheme (i.e., initial two doses only)				
Yes	5 (9.6%)	19 (36.5%)	11 (21.2%)	<0.001
No	28 (53.8%)	0 (0.0%)	41 (78.8%)	
Total	33(63.5%)	19 (36.5%)	52 (100%)	

¹Professionals included in the study: social workers, nutritionists, and psychologists.

these two latter individuals, 12 (23%) did not seroconvert to IgM throughout the study. The reactivity rate for IgM varied from 65%, 36%, 24%, 10%, 15%, and 24%, (Figure 3A) whereas for IgG it was 83%, 85%, 90%, 76% 58%, and 72% (Figure 3B) on days D7-30, D60, D90, D180, D270, and D450, respectively. Immunoglobulin levels decreased over time, reaching their lowest level on day 270 after infection when the mean level for IgM was $13.8 (\pm 17, p < 0.0001)$ (Figure 3A) and for IgG was $68.5 (\pm 72.3, p < 0.0001)$ demonstrating a significant reduction in comparison to the values observed at D7-D30. Figures 3C, D show the individual profile for anti-SARS-CoV-




FIGURE 2
 Overview of participant cohorts. Longitudinal follow-up of 52 participants who had COVID-19 confirmed by RT-qPCR. The day of symptom onset was called 0 (Black stroke - Symptom onset). The longitudinal timeline shows the days of the start of the vaccination schedule (pink dots), of the complete vaccination schedule, considering 15 days after the second dose (blue dots), and of the booster dose (green dots). Nineteen participants had confirmed SARS-CoV-2 reinfection (red dots).

2 IgM and IgG in the study participants, respectively. Although the mean levels of anti-SARS-CoV-2 IgG antibody did not differ significantly between days 270 and 450, representing the period when the majority of reinfection and vaccination booster doses occurred, a tendency for IgG levels to increase was observed (Figures 3A-F). There was no difference between biological sex in the dynamics of antibodies (Figures 4A, B). However, significant differences in the dynamics of antibody levels were observed, with older individuals presenting the highest levels of IgM and IgG (Figures 4C, D) and a slower decrease in anti-SARS-CoV-2 IgG levels (Figure 4D) compared to IgM (Figure 4C). No significant differences in the Ct value obtained in the RT-qPCR at baseline were observed between the different sex and the age groups (Figures 4E, F).

The dynamics of IgM and IgG levels over time are shown in Figures 3E, F. For IgM and IgG immunoglobulins, there were two behaviors: the drop in the mean antibody levels up to 180 and 270 days, respectively, followed by a slight increase after this time point. Tables 2 and 3 demonstrate the behavior of IgM and IgG estimated by the segmented model. The GEE results indicate that the average IgM and IgG values decreased by 0.606 and 0.645 units each day until days 180 and 270, respectively. After this time, the slope is positive for both IgM (0.363) and IgG (0.224). As shown in Figure 3E, the observed mean IgM levels were below the cutoff despite this upward trend. We can also observe that 60 days after symptom onset, the mean anti-SARS-CoV-2 IgM levels are below the cutoff, while anti-SARS-CoV-2 IgG levels drop, but not below the cutoff, around 270 days after symptom onset (Figures 3E, F).

No significant difference in IgM and IgG antibody levels with regard to the vaccination status of the participants was observed (Tables 2, 3). However, after the booster dose, the anti-SARS-CoV-2 IgM and IgG levels both significantly increased. The IgM levels after the booster dose were 38.388 units higher. In addition, IgM levels

were significantly higher after reinfection by 20.378 units. The interaction between these two covariables was not significant. In contrast, when we analyzed the IgG levels, the interaction between the booster dose and reinfection was significant, with the mean IgG levels 130.3 [95%CI: 82.19-178.30] units greater in those that received the booster dose and got reinfected. Before the booster dose, there was no difference in the mean IgG level when comparing reinfected and non-reinfected participants ($p = 0.348$). After the booster dose, the mean level of IgG for reinfected individuals was 87.8 [27.3; 148.0] units, which was higher than for non-reinfected individuals. For non-reinfected individuals, after the booster dose, the mean IgG was 52.22 units higher than before.

Of the 19 participants were reinfected (Figure 5), 13 (68.4%) were after and 6 (31.6%) before the booster dose. In four (21.0%), increased levels of IgM and IgG antibodies were not observed, even after booster dose and re-infection, these individuals being non-reactive to SAR-CoV-2 antigens at day 450 for both immunoglobulin classes (Figures 5F, G, J, O). One individual (Figure 5F) did not seroconverted during the whole period of follow-up. Of the remaining 15 individuals were reinfected, considering IgM dynamics, four of them never seroconverted during the follow-up period (Figures 5B, H, K, S). Four individuals (Figures 5C-E, R) who were negative during visits D7-D30 only had IgM seroconversion after reinfection or the booster dose. For four other individuals (Figures 5L, M, P, Q) that became IgM reactive during period D7-D30, but turned non-reactive during the longitudinal assessment, the booster dose and reinfection were not able to stimulate production of this class of antibody. For the remaining three participants (Figures 5A, I, N), IgM levels declined to non-reactive, but the booster dose or reinfection seroconverted them again. For IgG, the dynamics are different, since all 15 reinfected participants seroconverted at some point during the study. Of these, one only seroconverted after the booster dose (Figure 5K). The other 14




FIGURE 3
 Longitudinal humoral immune response in participants infected with COVID-19. Kinetics of the levels of anti-SARS-CoV-2 IgM and IgG antibodies over time. A serum sample was considered positive when the reaction intensity had a value equal to or above 30. Colored dots indicate individuals who became reinfected over the course of the study. (A, B) Show the reactivity of IgM and IgG specific for SARS-CoV-2 antigens over time. The frequency of seroconversion is shown at the top of the figure. The number of individuals tested (*n*) varies according to the time point evaluated and is indicated on the graph. **p*<0.05; ***p*<0.01; ****p*<0.001 *****p*<0.0001. Statistical significance was measured using a Kruskal-Wallis test at a significance level of 5%. (C, D) Each line represents one participant (*n* = 52). Dashed lines represent participants infected once with COVID-19. Reinfected participants are represented by solid lines. The red lines indicate the interval in which reinfection occurred. (A–D) Values below 30 are shown in the gray zone of the graphs. (E, F) Mean IgM and IgG immunoglobulin levels over time. For the analysis, the peak antibody level for each participant, observed from the seventh to the thirtieth day, was considered (30). Follow-up evaluation of anti-SARS-CoV-2 IgM and IgG levels was performed at days 60, 90, 180, 270, and 450 after enrollment in the study. A test was considered positive when the detected value was equal to or greater than 30.

participants had reactive IgG antibodies at baseline. For all of them, anti-SARS-CoV-2 IgG antibody levels declined, with seven participants becoming non-reactive to SARS-CoV-2 antigens for this class of antibody. For the remaining seven seroreactive participants, six showed increased IgG levels after reinfection or booster dose (Figures 5H, I, L, M, P, R). For the individuals who became IgG seronegative over the course of the study, the booster dose or reinfection was able to cause secondary IgG seroconversion of all such participants (Figures 5A–E, N, S).

3.3 Reinfection

Every 30 days or when participants presented symptoms that indicated a suspicion of COVID-19 reinfection, saliva or nasopharyngeal swab collection was performed to detect viral RNA. At the same time, a capillary blood sample was collected to evaluate immunoglobulins. The interval between symptoms onset and swab/saliva collection or antibody assessment varies from 1 to 15 days,

with 75% of the participants being tested until 6 days of symptoms onset (Supplementary Figure S1). Of the 52 participants, 46 (88%) described at least some symptoms compatible with suspected COVID-19, with 9 (17%) reporting it once, 19 (36%) twice, 9 (17%) three times, five (10%) four times, and four (8%) participants reporting five times (Figure 6A). In total, 114 suspected episodes of COVID-19 were recorded, all suspected cases were tested by RT-qPCR and of these only 19 were confirmed positive by RT-qPCR. Of all the symptom episodes/types (*n* = 185) reported by participants during the study, cough (61.6%) followed by congestion or runny nose (49.2%) and sore throat (41.1%) were the most frequent. The most common symptoms reported during the suspected and confirmed reinfections were congestion or runny nose (57.0% and 74%), cough (40.4% and 63%), and sore throat (45.6% and 47%), respectively. The profile of symptoms presented by reinfected individuals differs from those reported during the first SARS-CoV-2 infection, with statistical significance (*p*<0.001) for headache, fever and chills, myalgia, anosmia, and ageusia between groups were observed (Table 4). Ageusia and anosmia were not




FIGURE 4 Dynamics of IgM and IgG levels against SARS-CoV-2 and Ct values over time stratified by sex and age. **(A-D)** The left and right panels represent the distribution of IgM and IgG levels, respectively. A serum sample was considered positive when the reaction intensity had a value equal to or above 30. Values below 30 are shown in the gray zone of the graphs. P values were determined using a Sidak’s multiple comparisons test, ANOVA, or unpaired t-test. **(A, B)** Stratification of the humoral response over time by sex. The triangles represent the antibody level of women and the squares that of men. There was no statistical difference. **(C, D)** Stratification of humoral response over time by age, 21-30 years (triangles), 31-40 years (squares), and 41-60 years (dots). “a” significant difference in comparison with D7-D30 ($p < 0.005$); “b” significant differences in comparison with D180 ($p < 0.005$); “c” significant differences in comparison with D270 ($p < 0.005$); * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. **(E)** Ct values of males (squares) and females (dots). Statistical significance was measured using an unpaired t-test at a significance level of 5%. **(F)** Ct values in participants stratified by age: 21-30 years (dots), 31-40 years (squares), and 41-60 years (triangles). Statistical significance ANOVA at a significance level of 5%.

TABLE 2 Parameter estimates according to Generalized Estimating Equations (GEE) analysis of the relative numeric scale of IgM levels.

Covariable	Estimated IgM level in arbitrary units (95%CI)	p-value
Time		
≤180	-0.606 (-0.884; -0.328)	< 0.001
>180	0.363 (0.172; 0.554)	< 0.001
Vaccination status		
Unvaccinated	Reference	
Partially vaccinated	22.151 (-10.597; 54.90)	0.185
Fully vaccinated	9.934 (-8.511; 28.38)	0.291
Booster dose		
Before the booster dose	Reference	
After the booster dose	38.388 (19.091; 57.685)	< 0.001
Reinfection status		
No reinfection	Reference	
Reinfected	20.378 (2.869; 37.888)	0.023

Vaccine status, follow-up time, booster dose, and reinfection were used as covariates.

TABLE 3 Parameter estimates according to Generalized Estimating Equations (GEE) analysis of the relative numeric scale of IgG levels.

Covariable	Estimated IgG level in arbitrary units (95%CI)	p-value
Time		
<=270	-0.645 (-0.811; -0.479)	< 0.001
>270	0.224 (0.127; 0.321)	< 0.001
Vaccination status		
Unvaccinated	Reference	
Partially vaccinated	15.977 (-15.343; 47.298)	0.317
Fully vaccinated	26.848 (-7.077; 60.773)	0.121
Booster dose		
Before the booster dose	Reference	
After the booster dose	52.221 (15.406; 89.035)	0.005
Reinfection status		
No reinfection	Reference	
Reinfected	9.715 (-10.566; 29.996)	0.348
Interaction		
Booster dose - Reinfection	87.8 (27.3; 148.0)	0.001
Reinfection - Booster dose	130.3 (82.19;178.30)	0.001

Vaccine status, follow-up time, booster dose, and reinfection were used as covariates.

reported in confirmed cases of reinfection (p<0.001). Other symptoms reported during the first infection in the suspected and confirmed cases of reinfection can be seen in Table 4.

The first five (26%) cases of reinfection occurred between August and September 2021 (Table 5). At this time, the Delta and Gamma variants were circulating in Belo Horizonte (Figure 7).

Between December 2021 and January 2022, 12 cases (63%) of reinfection were confirmed, a period that overlaps with the new wave of transmission caused by the Omicron variant (Figure 7). Of the reinfection samples collected during this period that were sequenced, nine were identified as the Omicron variant. Sequencing samples with low viral load (Ct value > 36) was not




FIGURE 5 IgM and IgG antibody profiles of participants who were reinfected. (A-S) The triangles represent the IgM and the dots the IgG values. The gray, blue and green dashed lines represent the beginning date of the vaccination, the complete vaccination scheme, and the booster dose, respectively. The red line represents the reinfection period. A serum sample was considered reactive when the reaction intensity had a value equal to or above 30. Values below 30 are shown in the gray zone of graphs.




FIGURE 6 Main self-reported symptoms, dynamics of IgM and IgG levels, and Ct-values stratified according to SARS-CoV-2 infection/reinfection status. **(A)** Main self-reported symptoms described by participants. Blue bars indicate the percentage of symptoms reported by the 52 COVID-19 positive participants. Red bars show the symptoms reported in cases of suspected reinfection, but without confirmation by detection of viral RNA. Green bars show the main symptoms of participants with confirmed reinfection by SARS-CoV-2. Statistical significance was measured using either the Chi-square test or Fisher’s exact test $**p < 0.001$, $*p = 0.002$. **(B)** SARS-CoV-2 viral load. Statistical significance was measured using a paired t-test at a significance level of 5%. **(C, D)** Antibody levels of reinfected and non-reinfected participants. For individuals who were not reinfected, the values shown are the arithmetic means of multiple measurements of their circulating antibody levels taken throughout only the period of the study when the waves of the Delta and Omicron variants occurred. For individuals who were reinfected, the antibody levels shown are single point estimates determined only at the time of reinfection confirmation. The left and right panels represent the distribution of IgM **(C)** and IgG **(D)** levels, respectively. A serum sample was considered positive when the reaction intensity had a value equal to or above 30. Values below 30 are shown in the gray zone of graphs. P values were determined using the Mann-Whitney U test at a significance level of 5%.

possible. The mean CT of infected participants was $30.17 (\pm 6.5)$ (Table 5). There was no significant difference between Ct values detected during infection and reinfection (Figure 6B). When assessing the humoral response of reinfected and non-reinfected

individuals, we did not observe differences in IgM and IgG levels between these groups (Figures 6C, D). The intervals between infection and vaccination and seroconversion status are given in Tables 5, 6.

TABLE 4 Main symptoms reported by the final study cohort.

Symptoms	First infection	Suspected reinfection	Confirmed reinfection	Total	p-value
Congestion or runny nose	35 (67.3%)	65 (57.0%)	14 (73.7%)	114 (61.6%)	0.234
Cough	33 (63.5%)	46 (40.4%)	12 (63.2%)	91 (49.2%)	0.010
Sore throat	15 (28.8%)	52 (45.6%)	9 (47.4%)	76 (41.1%)	0.106
Headache	27 (51.9%)	22 (19.3%)	7 (36.8%)	56 (30.3%)	<0.001
Fever or chill	26 (50.0%)	11 (9.6%)	3 (15.8%)	40 (21.6%)	<0.001
Fatigue/Tiredness	20 (38.5%)	19 (16.7%)	1 (5.3%)	40 (21.6%)	0.002
Myalgias	28 (53.8%)	3 (2.6%)	1 (5.3%)	32 (17.3%)	<0.001
Anosmia (loss smell)	20 (38.5%)	5 (4.4%)	0 (0.0%)	25 (13.5%)	<0.001
Ageusia (loss of taste)	12 (23.1%)	3 (2.6%)	0 (0.0%)	15 (8.1%)	<0.001
Diarrhea	4 (7.7%)	18 (15.8%)	1 (5.3%)	23 (12.4%)	0.287
Other ¹	8 (15.4%)	15 (13.2%)	3 (15.8%)	26 (14.1%)	0.855
Total	52 (28.1%)	114 (61.6%)	19 (10.31%)	185 (100%)	

¹Other symptoms reported to healthcare professionals included: malaise, throat irritation, pain in the face, abdominal pain, inappetence, asthenia, arthralgia, retro orbital, pain and sweating, nausea or vomiting, and red or irritated eyes.

TABLE 5 Genomic strain, Ct value, and classification of symptoms of confirmed cases of SARS-CoV-2 infection and reinfection.

Date of onset of symptoms	Sample RT-PCR	CT (value)	Symptoms	Genomic Strain	Period between infections (days)	Date of onset of symptoms	Reinfection			
							Sample RT-PCR	CT (value)	Symptoms	Genomic Strain
Nov-20	Saliva	28.17	Mild	Zeta (P.2-like) ^{&&}	214	Sept-21	Saliva	31.92	Mild	ND
Nov-20	Saliva	33.94	Mild	ND	433	Jan-22	Saliva	36.94	Mild	ND
Nov-20	NS	24.25	Mild	Zeta (P.2-like) ^{&1}	408	Jan-22	NS	27.37	Mild	Omicron (BA.1.15) ^{&&}
Dec-20	NS	19.22	Mild	B.1.1.33 ^{&2}	299	Dec-21	NS	36.89	Mild	ND
Jan-21	NS	27.95	Mild	Zeta (P.2-like) ^{&3}	388	Jan-22	Saliva	29.41	Mild	ND
Jan-21	NS	25.61	Mild	Zeta (P.2-like) ^{&&}	386	Jan-22	Saliva	16.8	Mild	Omicron (BA.1.1) ^{&10}
Jan-21	NS	14.85	Mild	B.1.1.28 ^{&4}	359	Sept-21	Saliva	36.45	Mild	ND
Jan-21	Saliva	25.52	Mild	ND	258	Jan-22	Saliva	18.58	Mild	Omicron (BA.1.1) ^{&11}
Jan-21	Saliva	30.44	Mild	B.1.1.28 ^{&5}	360	Jan-22	NS	28.95	Mild	Omicron (BA.1.1.1) ^{&&}
Jan-21	NS	23.07	Mild	Zeta (P.2-like) ^{&6}	370	Jan-22	Saliva	36.73	Mild	ND
Jan-21	Saliva	34.9	Mild	ND	354	Aug-21	Saliva	34.24	Mild	ND
Jan-21	Saliva	35.75	Moderate	B.1.1 ^{&&}	339	Jan-22	NS	31.91	Mild	Omicron (BA.1) ^{&&}
Mar-21	Saliva	30.38	Mild	ND	181	May-22	NS	27.17	Mild	Omicron (BA.2) ^{&12}
Mar-21	Saliva	28.98	Mild	B.1.1 ^{&&}	427	Aug-21	NS	34.78	Mild	B.1 ^{&&}
Mar-21	Saliva	35.19	Mild	ND	446	Jun-22	NS	24.16	Mild	Omicron (BA.2.23) ^{&13}
Mar-21	Saliva	32.38	Moderate	ND	314	Jan-22	Saliva	36.31	Mild	ND
Mar-21	NS	21.88	Mild	Gamma (P.1-like) ^{&7}	222	Aug-21	Saliva	34.78	Mild	ND
Mar-21	NS	22.79	Mild	Gamma (P.1-like) ^{&8}	287	Jan-22	Saliva	30.54	Mild	Omicron (BA.1.1.1) ^{&&}

(Continued)

TABLE 5 Continued

Date of onset of symptoms	Sample RT-PCR	CT (value)	Symptoms	Genomic Strain	Period between infections (days)	Date of onset of symptoms	Reinfection			
							Sample RT-PCR	CT (value)	Symptoms	Genomic Strain
Apr-21	NS	19.17	Mild	Gamma (P.1-like) ⁸⁹	267	Jan-22	Saliva	19.39	Mild	Omicron (BA.1) ^{81,4}

Mild symptoms: no oxygen support but other COVID-19 symptoms (which may include congestion or runny nose, cough, sore throat, fever, chills, shortness of breath, change in taste or smell, headache, nausea, vomiting, diarrhea, fatigue, myalgias, and other symptoms). Moderate symptoms: supplemental oxygen. Severe symptoms: intensive care unit admission. NS: nasopharyngeal swabs. ND: Not determined. & Submitted in GISAID. && No coverage for submission in GISAID. Submission number GISAID &1: EPI_ISL_3031326; &2: EPI_ISL_3031330; &3: EPI_ISL_3031331; &4: EPI_ISL_3031335; &5: EPI_ISL_3031336; &6: EPI_ISL_3031346; &7: EPI_ISL_12042548; &8: EPI_ISL_12042549; &9: EPI_ISL_10706100; &11: EPI_ISL_10706101; &12: EPI_ISL_14675090; &13: EPI_ISL_14675091; &14: EPI_ISL_10706102.

4 Discussion

Elucidating the kinetics of the humoral response to SARS-CoV-2 is crucial for controlling the pandemic, and for designing, planning and implementing the most appropriate vaccine schemes (1). Here, we investigated for 450 days the dynamics and longevity of IgM and IgG antibodies from healthcare professionals. The long follow-up period allowed us to monitor dynamics of the humoral response after vaccination booster doses and reinfection.

Many previous studies have evaluated this profile. However, such studies have a variety of follow-up durations ranging from 50 days (33), 100 days (34–37), 210 days (38), 360 days (39), and 480 days (40). Some of these studies followed specific antibodies against the nucleocapsid (N), the receptor-binding domain (RBD), or the spike (S) protein, using either ELISA assays, immunochromatographic tests, or looking for neutralizing antibodies. In our study, a commercial immunochromatographic test was used to detect both IgM and IgG against viral proteins. It has been shown by some authors that there is a correlation between the detection of total and neutralizing antibodies (36, 40, 41).

Our study reinforces the view, and provides evidence, that antibodies are initially produced, but decline over time. The drop of IgM starts on the 30th day, reaching non-reactive levels by the 60th day after symptom onset, the same profile as previously described (36, 37). For IgG, a less pronounced decline is observed, and around the ninth month, we observe the lowest mean reactivity. The stability of the IgG reactivity for three months has been previously demonstrated (36, 37). Gil-Manso et al. (2020) (42) and Gaebler (2021) (43) observed that the IgG response lasted longer, about six months, and the levels of neutralizing activity were proportional to anti-RBD IgG antibody titers. Some studies demonstrate that the duration of the response depends on the studied target: anti-RBD antibodies remained stable for between six to 12 months, while anti-N antibodies decreased over the same period (39). In contrast, Yang (2022) (40) described a peak of anti-RBD antibodies around 120 days after the onset of symptoms with a subsequent decline, maintaining positivity until day 400 after symptom onset. A recent study describes that for non-vaccinated individuals, IgG antibodies, evaluated by ELISA, persist for one year (44).

The results of longitudinal studies, however, may critically vary according to different conditions and variables. Mioch et al. (2023) (44) report that loosening epidemiological control measures increases the chance of re-exposure to the virus. The sensitivity of the tests and different methods used may vary. In addition, patients with comorbidities may have different antibody kinetics, as demonstrated by the rapid decline of antibodies in diabetic patients (45). Yang et al. (2022) (40) conducted a long follow-up of the humoral response in individuals confirmed to be free of re-exposure and vaccination against SARS-CoV-2. Although their study is significant, it does not portray the global reality since we have more than 60% of the population vaccinated worldwide, while in Brazil more than 80% of people completed the vaccination schedule and almost 50% took a booster dose (1). In our cohort, all healthcare professionals who worked on the front line completed the vaccination schedule with the Coronavac vaccine and took the Pfizer vaccine as a booster dose (6)




FIGURE 7
 Overlap of genomic strains sequenced in this study in relation to strains circulating in Belo Horizonte during the same period. Absolute frequency of SARS-CoV-2 genomic samples sequenced in Belo Horizonte from October 2020 to August 2022 (Stacked area graph). The colors inside the circles indicate the sequence of SARS-CoV-2 strains. White dots represent samples for which the SARS-CoV-2 strain was not determined. The dashed lines connect results obtained in samples from the same participant during their first infection and subsequent reinfection.

For our analyses, we considered the peak of reactivity to be between days 7-30 (D7-D30). Thus, there was no difference in the increase of antibody levels for fully and partially vaccinated and unvaccinated individuals. Based on the dynamics presented, the slight increase observed in the mean antibody levels (D60-D90) may have been induced by the Coronavac vaccination (46). After using the Generalized Estimating Equations (GEE), it is possible to establish that the Pfizer booster dose increased antibody levels for both IgM and IgG. The decline of antibodies after the Coronavac vaccination, even in individuals who became infected, has also been already described. The booster dose is essential to restimulate the humoral response (47–49) and the same profile is also observed for other vaccines (50–53).

Vaccines are essential to reduce the morbidity and mortality from COVID-19. However, they cannot completely prevent new infections and reinfections (50). In our cohort, 36% of participants were reinfected either before (31.6%) or after (68.4%) the booster dose. The reinfections coincided with the spread of new variants in Brazil, such as Delta and Gama in August-September 2021, and Omicron in December 2021. The emergence of new variants has been a matter of great concern, as they can reduce neutralization and even escape vaccination, as demonstrated for Delta (46, 54), Beta and Gamma (47, 55), and Omicron (56–58) variants.

Cases of reinfection have been reported since the beginning of the pandemic (12, 20, 59–61), including in Brazil (62–64). Studies have shown that an acquired immune response can reduce the risk of transmission by up to 90%, with an interval of 6-10 months (65–67). The reinfection rate is relatively low, ranging from 0.1-0.65 (18–20, 68–70), with the highest rates reported in the UK study at 1.9% and 4.5% in India (24). In our study, we reported a high rate of reinfection that might be due to some important factors. Most of the reinfections occurred nine months after the first infection, when

antibody levels were already low. Additionally, reinfections coincided with the entry of new variants into Brazil, which, as already mentioned, have a high rate of transmission, and escape from immune responses (14, 21, 25, 67, 71).

Some studies show that men are more likely to test positive and develop severe COVID-19 (72). Petersen et al. (2022) (73) showed a faster decrease in IgG in males. Frauke et al. (2022) (41) observed that the decline of neutralizing antibodies in men was faster than in women but that afterward, there was no difference in response. The same was observed by (74), in which no correlation exists between neutralizing antibodies and biological sex. We also did not notice any difference in the behavior of the humoral response between men and women. Evaluating reinfection cases, our data corroborate the data of Alexander Lawandi et al. (2022) (75), in which we observed a higher rate of reinfection in women, which contrasts with the review made by Sahar (2022) (24), while other studies did not find a relationship between sex and reinfection (76). This heterogeneity of results demonstrates that other factors must be evaluated, such as comorbidity, lifestyle, workplace, biological and immune differences. Understanding sex differences is fundamental to improving disease management, predicting outcomes, and planning specific interventions for men and women.

There is no relationship between age and cases of reinfection in health workers in our study, as demonstrated by Alejandra Svartz (2023) (77). There was also no difference between the general population and healthcare workers who were reinfected. Ren et al. (2022) (22) and Sahar Ghorbani (2022) (19) review that there is a wide age distribution among reinfected patients, ranging from 15 to 99 years. Hansen (2021) (78) described that an age greater than 65 years might influence the increase in the relative risk of reinfection. Healthcare professionals in our cohort who became reinfected were between 24 and 55 years old, and all of them had mild symptoms.

TABLE 6 Seroconversion of IgM and IgG after infection, vaccine doses, and reinfection.

Infection				Vaccination					Reinfection					
Date of onset of symptoms	Symptoms	Seroreactivity after 30 days (IgM)	Seroreactivity after 30 days (IgG)	Date of full vaccination	Vaccination time relative to infection time (days)	Seroreactivity after full vaccination (IgM)	Seroreactivity after full vaccination (IgG)	Booster dose date	Days between infection and booster dose	Period between infections (days)	Date of onset of symptoms	Symptoms	Seroreactivity after reinfection (IgM)	Seroreactivity after reinfection (IgG)
Nov-20	Mild	No	No	Feb-21	109	No	N/A	Nov-21	361	214	Sept-21	Mild	Yes	Yes
Nov-20	Mild	No	No	Mar-21	98	No	N/A	Nov-21	350	433	Jan-22	Mild	No	Yes
Nov-20	Mild	No	No	Feb-21	93	No	Yes	Oct-21	330	408	Jan-22	Mild	Yes	Yes
Dez-20	Mild	Yes	Yes	May-21	155	No	Yes	Feb-22	421	299	Dec-21	Mild	Yes	Yes
Jan-21	Mild	No	Yes	Feb-21	52	No	Yes	Dec-21	359	388	Jan-22	Mild	Yes	Yes
Jan-21	Mild	No	No	Mar-21	53	No	No	Oct-21	268	386	Jan-22	Mild	No	No
Jan-21	Mild	No	Yes	Feb-21	46	No	Yes	Oct-21	280	359	Sept-21	Mild	No	No
Jan-21	Mild	No	Yes	Apr-21	85	No	Yes	Oct-21	281	258	Jan-22	Mild	No	Yes
Jan-21	Mild	Yes	Yes	Apr-21	85	Yes	Yes	Jun-22	504	360	Jan-22	Mild	Yes	Yes
Jan-21	Mild	Yes	Yes	Apr-21	73	No	Yes	Oct-21	276	370	Jan-22	Mild	No	No
Jan-21	Mild	No	No	Feb-21	24	No	No	Oct-21	265	354	Aug-21	Mild	No	No
Jan-21	Moderate	Yes	Yes	Mar-21	48	No	Yes	Nov-21	276	339	Jan-22	Mild	No	Yes
Mar-21	Mild	Yes	Yes	Feb-21	-33	Yes	Yes	Oct-21	212	181	May-22	Mild	No	Yes
Mar-21	Mild	Yes	Yes	Feb-21	-25	Yes	Yes	Nov-21	245	427	Aug-21	Mild	No	No
Mar-21	Mild	Yes	Yes	Mar-21	-9	Yes	Yes	Oct-21	222	446	Jun-22	Mild	No	Yes
Mar-21	Moderate	Yes	Yes	Feb-21	-26	Yes	Yes	Oct-21	220	314	Jan-22	Mild	No	Yes
Mar-21	Mild	Yes	Yes	Mar-21	-21	Yes	Yes	Oct-21	205	222	Aug-21	Mild	Yes	Yes
Mar-21	Mild	No	Yes	May-21	-52	No	Yes	Nov-21	229	287	Jan-22	Mild	No	Yes
Apr-21	Mild	No	Yes	Mar/21	-50	No	Yes	Out-21	169	267	Jan-22	Mild	No	Yes

Even though age was not associated with reinfection, we observed that the dynamics of antibody levels varied over time and behaved differently across age groups. The level of antibodies produced by the younger group was lower than the older groups, as observed by others works (73, 79, 80).

Many studies have sought to understand and relate antibody profiles and Ct values with disease severity and protection against reinfection (34, 40, 81, 82). Omid Dadras et al., 2022 (83), concluded that the relationship between viral load and disease severity is inconclusive. We observed no significant difference in viral load at first infection and reinfection. Except for two participants who required oxygen support for their first infection, all other participants had mild disease. As expected, the main symptoms reported were nasal congestion, coughing, sore throat, and headache. Other symptoms were also reported as described in other studies (22). The reinfection period of most participants in our study overlaps with the spread of the Omicron variant, as described by Menni et al. (2022), Karina Vihta (2022) (84, 85) and Machado Curbelo (2022) (86), who demonstrated that anosmia and ageusia were less frequently associated with the Omicron variant. No reinfected participants reported anosmia or ageusia in our study. Symptoms such as cough, fever, shortness of breath, myalgia, fatigue, and headache were less frequently reported by participants who had suspicion of COVID-19 either confirmed or not by RT-qPCR. In contrast, sore throat was more frequently reported by those participants, although no significant difference in frequency was observed among infection/reinfection status. These clinical conditions corroborate those described by Karina Vihta (2022) (85).

Our study has some limitations. We evaluated total antibodies and did not determine if they were neutralizing, nor the quality of memory B cells necessary to produce antibodies against reinfection. Also, we monitored reinfection in symptomatic participants, but there is a possibility that cases of asymptomatic reinfection also occurred (24).

The humoral response declines after the first infection and vaccination but increases substantially after reinfection and booster doses, especially for the IgG antibody class. There is no association between circulating antibody levels and cases of reinfection. Overall, we demonstrated that even after the booster dose, health professionals can become reinfected with new variants of SARS-CoV-2. Therefore, our study demonstrates that prolonged protection after COVID-19 infection, even after the booster dose, does not prevent reinfection by new variants, which contrasts with the prolonged immune response cited by other studies (73, 87). Studies show that previous infection and booster dose reduce the risk of reinfection, as seen in Switzerland (88), Qatar (68), and United States of America (89). However, studies have shown a significant decrease in the effectiveness of the vaccine against the Omicron variant within a few months after administration (90). Although our study was not designed to assess whether the booster dose would prevent reinfection, the evidence suggests that the booster dose was not effective in preventing reinfection. Those differences might have been impacted by the vaccine scheme and/or type used during the vaccination campaign.

In this context, our data reinforce the importance of robust surveillance in viral genomics and in the immune response of

individuals, especially in high-risk individuals, such as the immunocompromised and health care professionals. Investing in these tools is essential for preparing and responding to new variants and future pandemics.

Data availability statement

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: <https://www.ebi.ac.uk/ena/browser/view/PRJEB49204>, PRJEB49204 (The European Nucleotide Archive).

Ethics statement

The studies involving human participants were reviewed and approved by Institutional review board of the Instituto René Rachou, Fundação Oswaldo Cruz, CAAE: 31.919520.8.0000.5091, approval numbers: 4177931; 4291836; 4343318; 4624187; 5294423. Written informed consent was obtained from all participants before any study procedure was undertaken. The patients/participants provided their written informed consent to participate in this study.

Author contributions

Conceptualization, CF, MW, PA and RS-P; methodology, CF, GF, PA and SG; validation, NF, TS, TGS and WB; formal analysis, AF-L, GF, ES and TS; investigation, NF, TS, TGS and WB; resources, BA, CF, MM, MW and PA; data curation, CF, NF, TGS, RS-P and WB; writing—original draft preparation, AF-L; writing review and editing, CF, GF, MW, NF, PA, RS-P, SG, TS, TCS, TGS and WB; visualization, AF-L, CF and PA; supervision, CF; project administration, CF and WB; funding acquisition, CF. All authors contributed to the article and approved the submitted version.

Funding

This research was funded by the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) (Grant number: APQ-00635-20) and the Fundação Oswaldo Cruz – Inova Program (Grant number: VPPCB-005-FIO-20-2-24) and CNPq (Fellowship Grant number: CTF- 306188/2022-8).

Acknowledgments

The authors thank all participants for their involvement in the study, and the healthcare and administrative personnel at the hospitals participating in this study. We acknowledge the Program for Technological Development in Tools for Health—PDTIS-FIOCRUZ for the use of its Real-Time PCR and Next Generation Sequencing facilities; the Clinical Research Platform of

Fiocruz for monitoring the study documents and for generating and managing the study database; the project support service of the IRR for project management and Bio-Manguinhos for providing the DPP[®] COVID-19 IgM/IgG kits. We also thank Edward José Oliveira, Marina Moraes Mourão and Roberta Lima Caldeira for their contributions towards the conceptualization of the study and Luke Baton for revising the language of the manuscript.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. WHO Coronavirus (COVID-19) Dashboard. *WHO coronavirus (COVID-19) dashboard with vaccination data* (2023). Available at: <https://covid19.who.int/>.
2. Smallwood N, Harrex W, Rees M, Willis K, Bennett CM. COVID-19 infection and the broader impacts of the pandemic on healthcare workers. *Respirology* (2022) 27:411–26. doi: 10.1111/resp.14208
3. Zheng C, Shao W, Chen X, Zhang B, Wang G, Zhang W. Real-world effectiveness of COVID-19 vaccines: a literature review and meta-analysis. *Int J Infect Diseases* (2022) 114:252–60. doi: 10.1016/j.ijid.2021.11.009
4. Uysal EB, Gümüş S, Bektöre B, Bozkurt H, Gözalan A. Evaluation of antibody response after COVID-19 vaccination of healthcare workers. *J Med Virol* (2022) 94(3):1060–6. doi: 10.1002/jmv.27420
5. Alejandro Gómez-Ochoa S, Franco OH, Rojas LZ, Raguindin PF, Roa-Díaz M, Wyssmann BM, et al. COVID-19 in healthcare workers: a living systematic review and meta-analysis of prevalence, risk factors, clinical characteristics, and outcomes. *Am J Epidemiol* (2021) 190(1):161–75. doi: 10.1093/aje/kwaa191
6. Conselho Nacional de Saúde do Brasil. *Vacinômetro da COVID-19*. (2022). Available at: <https://www.gov.br/saude/pt-br/composicao/seidigi/demas/covid19>.
7. Sciscent BY, Eisele CD, Ho L, King SD, Jain R, Golamari RR. COVID-19 reinfection: the role of natural immunity, vaccines, and variants. *J Community Hosp Intern Med Perspect* (2021) 11(6):733–9. doi: 10.1080/20009666.2021.1974665
8. Hadj Hassine I. Covid-19 vaccines and variants of concern: a review. *Rev Med Virol* (2022) 32:e2313. doi: 10.1002/rmv.2313
9. Fiolet T, Kherabi Y, MacDonald CJ, Ghosn J, Peiffer-Smadja N. Comparing COVID-19 vaccines for their characteristics, efficacy and effectiveness against SARS-CoV-2 and variants of concern: a narrative review. *Clin Microbiol Infection* (2022) 28:202–21. doi: 10.1016/j.cmi.2021.10.005
10. Havers FP, Patel K, Whitaker M, Milucky J, Reingold A, Armistead I, et al. Morbidity and mortality weekly report laboratory-confirmed COVID-19-Associated hospitalizations among adults during SARS-CoV-2 omicron BA.2 variant predominance-COVID-19-Associated hospitalization surveillance network, 14 states. *Morb Mortal Wkly Rep* (2022) 71(34):1085–91. doi: 10.15585/mmwr.mm7134a3
11. Havers FP, Pham H, Taylor CA, Whitaker M, Patel K, Anglin O, et al. COVID-19-Associated hospitalizations among vaccinated and unvaccinated adults 18 years or older in 13 US states, January 2021 to April 2022. *JAMA Intern Med* (2022) 182(10):1071–81. doi: 10.1001/jamainternmed.2022.4299
12. Nordström P, Ballin M, Nordström A. Risk of SARS-CoV-2 reinfection and COVID-19 hospitalisation in individuals with natural and hybrid immunity: a retrospective, total population cohort study in Sweden. *Lancet Infect Dis* (2022) 22(6):781–90. doi: 10.1016/S1473-3099(22)00143-8
13. He X, Hong W, Pan X, Lu G, Wei X. SARS-CoV-2 omicron variant: characteristics and prevention. *MedComm* (2021) 2(4):838–45. doi: 10.1002/mco.2.110
14. Araf Y, Akter F, dong TY, Fatemi R, Parvez MSA, Zheng C, et al. Omicron variant of SARS-CoV-2: genomics, transmissibility, and responses to current COVID-19 vaccines. *J Med Virology* (2022) 94:1825–32. doi: 10.1002/jmv.27588
15. Bowe B, Xie Y, Al-Aly Z. Acute and postacute sequelae associated with SARS-CoV-2 reinfection. *Nat Med* (2022) 28(11):2398–405. doi: 10.1038/s41591-022-02051-3
16. Duggan NM, Ludy SM, Shannon BC, Reisner AT, Wilcox SR. Is novel coronavirus 2019 reinfection possible? interpreting dynamic SARS-CoV-2 test results. *Am J Emergency Med* (2021) 39:256.e1–3. doi: 10.1016/j.ajem.2020.06.079

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2023.1220600/full#supplementary-material>

17. CDC. *Coronavirus disease 2019 (COVID-19) 2021 case definition* | CDC. (2022). Available at: <https://ndc.services.cdc.gov/case-definitions/coronavirus-disease-2019-2021/>.
18. Flacco ME, Acuti Martellucci C, Soldato G, Carota R, Fazio P, Caponetti A, et al. Rate of reinfections after SARS-CoV-2 primary infection in the population of an Italian province: a cohort study. *J Public Health (Oxf)* (2022) 44(4):e475–e478. doi: 10.1093/pubmed/fdab346
19. Sotoodeh Ghorbani S, Taherpour N, Bayat S, Ghajari H, Mohseni P, Hashemi Nazari SS. Epidemiologic characteristics of cases with reinfection, recurrence, and hospital readmission due to COVID-19: a systematic review and meta-analysis. *J Med Virol* (2022) 94(1):44–53. doi: 10.1002/jmv.27281
20. Nguyen NN, Houhamdi L, Hoang VT, Delerce J, Delorme L, Colson P, et al. SARS-CoV-2 reinfection and COVID-19 severity. *Emerg Microbes Infect* (2022) 11(1):894–901. doi: 10.1080/22221751.2022.2052358
21. Pulliam JRC, van Schalkwyk C, Govender N, von Gottberg A, Cohen C, Groome MJ, et al. Increased risk of SARS-CoV-2 reinfection associated with emergence of omicron in south Africa. *Science* (2022) 376(6593):eabn4947. doi: 10.1126/science.abn4947
22. Ren X, Zhou J, Guo J, Hao C, Zheng M, Zhang R, et al. Reinfection in patients with COVID-19: a systematic review. *Glob Health Res Policy* (2022) 7(1):1–20. doi: 10.1186/s41256-022-00245-3
23. Havervall S, Ng H, Jernbom Falk A, Greilert-Norin N, Månberg A, Marking U, et al. Robust humoral and cellular immune responses and low risk for reinfection at least 8 months following asymptomatic to mild COVID-19. *J Intern Med* (2022) 291(1):72–80. doi: 10.1111/joim.13387
24. Hall VJ, Foulkes S, Charlett A, Atti A, Monk EJM, Simmons R, et al. SARS-CoV-2 infection rates of antibody-positive compared with antibody-negative health-care workers in England: a large, multicentre, prospective cohort study (SIREN). *Lancet* (2021) 397(10283):1459–69. doi: 10.1016/S0140-6736(21)00675-9
25. Negi N, Maurya SP, Singh R, Das BK. An update on host immunity correlates and prospects of re-infection in COVID-19. *Int Rev Immunol* (2022) 41(4):367–92. doi: 10.1080/08830185.2021.2019727
26. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DKW, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* (2020) 25(3):2000045. doi: 10.2807/1560-7917.ES.2020.25.3.2000045
27. Carvalho AF, Rocha RP, Gonçalves AP, Silva TBS, Sato HI, Vuitika L, et al. The use of denaturing solution as collection and transport media to improve SARS-CoV-2 RNA detection and reduce infection of laboratory personnel. *Braz J Microbiol* (2021) 52(2):531–9. doi: 10.1007/s42770-021-00469-4
28. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics* (2014) 30(15):2114–20. doi: 10.1093/bioinformatics/btu170
29. Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, et al. An amplicon-based sequencing framework for accurately measuring intrahost virus diversity using PrimalSeq and iVar. *Genome Biol* (2019) 20(1):8. doi: 10.1186/s13059-018-1618-7
30. O'Toole Á, Scher E, Underwood A, Jackson B, Hill V, McCrone JT, et al. Assignment of epidemiological lineages in an emerging pandemic using the pangolin tool. *Virus Evol* (2021) 7(2):veab064. doi: 10.1093/ve/veab064
31. R Core Team. (2022).

32. Liang KY, Zeger SL. Longitudinal data analysis using generalized linear models. *Biometrika* (1986) 73(1):13–22. doi: 10.1093/biomet/73.1.13
33. Prévost J, Gasser R, Beaudoin-Bussièrès G, Richard J, Duerr R, Laumaea A, et al. Cross-sectional evaluation of humoral responses against SARS-CoV-2 spike. *Cell Rep Med* (2020) 1(7):100126. doi: 10.1016/j.xcrm.2020.100126
34. Minervina AA, Komech EA, Titov A, Koraichi MB, Rosati E, Mamedov IZ, et al. Longitudinal high-throughput tcr repertoire profiling reveals the dynamics of t-cell memory formation after mild covid-19 infection. *Elife* (2021) 10:1–17. doi: 10.7554/eLife.63502
35. Talla A, Vasaikar SV, Lemos MP, Moodie Z, Pebworth MPL, Henderson KE, et al. Longitudinal immune dynamics of mild COVID-19 define signatures of recovery and persistence. *bioRxiv [Preprint]* (2021) 2021.05.26.442666. doi: 10.1101/2021.05.26.442666
36. Seow J, Graham C, Merrick B, Acors S, Pickering S, Steel KJA, et al. Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. *Nat Microbiol* (2020) 5(12):1598–607. doi: 10.1038/s41564-020-00813-8
37. Isho B, Abe KT, Zuo M, Jamal AJ, Rathod B, Wang JH, et al. Persistence of serum and saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. *Sci Immunol* (2020) 5(52):eabe5511. doi: 10.1126/sciimmunol.abe5511
38. Figueiredo-Campos P, Blankenhaus B, Mota C, Gomes A, Serrano M, Ariotti S, et al. Seroprevalence of anti-SARS-CoV-2 antibodies in COVID-19 patients and healthy volunteers up to 6 months post disease onset. *Eur J Immunol* (2020) 50(12):2025–40. doi: 10.1002/eji.202048970
39. Teyssou E, Zafilaza K, Sayon S, Marot S, Dropy M, Soulie C, et al. Long-term evolution of humoral immune response after SARS-CoV-2 infection. *Clin Microbiol Infection* (2022) 28(7):1027.e1–1027.e4. doi: 10.1016/j.cmi.2022.03.012
40. Yang Y, Yang M, Peng Y, Liang Y, Wei J, Xing L, et al. Longitudinal analysis of antibody dynamics in COVID-19 convalescents reveals neutralizing responses up to 16 months after infection. *Nat Microbiol* (2022) 7(3):423–33. doi: 10.1038/s41564-021-01051-2
41. Muecksch F, Wise H, Templeton K, Batchelor B, Squires M, McCance K, et al. Longitudinal variation in SARS-CoV-2 antibody levels and emergence of viral variants: implications for the ability of serological assays to predict immunity. *medRxiv* (2021) 2021.07.02.21259939. doi: 10.1101/2021.07.02.21259939
42. Gil-Manso S, Alonso R, Catalán P, Sánchez-Arcilla I, Marzola M, Correa-Rocha R, et al. IgG anti-RBD levels during 8-month follow-up post-vaccination with BNT162b2 and mRNA-1273 vaccines in healthcare workers: a one-center study. *Front Cell Infect Microbiol* (2022) 12. doi: 10.3389/fcimb.2022.1035155
43. Gaebler C, Wang Z, Lorenzi JCC, Muecksch F, Finkin S, Tokuyama M, et al. Evolution of antibody immunity to SARS-CoV-2. *Nature* (2021) 591(7851):639–44. doi: 10.1038/s41586-021-03207-w
44. Mioch D, Vanbrabant L, Reimerink J, Kuiper S, Lodder E, van den Bijllaardt W, et al. SARS-CoV-2 antibodies persist up to 12 months after natural infection in healthy employees working in non-medical contact-intensive professions. *Int J Infect Dis* (2023) 126:155–63. doi: 10.1016/j.ijid.2022.11.025
45. Sourij C, Aziz F, Kojzar H, Obermayer AM, Sternad C, Müller A, et al. Severe acute respiratory syndrome coronavirus 2 spike antibody level decline is more pronounced after the second vaccination, but response to the third vaccination is similar in people with type 1 and type 2 diabetes compared with healthy controls: the prospective COVAC-DM cohort study. *Diabetes Obes Metab* (2023) 25(1):314–8. doi: 10.1111/dom.14855
46. Cheng ZJ, Huang H, Zheng P, Xue M, Ma J, Zhan Z, et al. Humoral immune response of BBIBP COVID-19 vaccination before and after the booster immunization. *Allergy* (2022) 77(8):2404–14. doi: 10.1111/all.15271
47. Mok CKP, Cohen CA, Cheng SMS, Chen C, Kwok KO, Yiu K, et al. Comparison of the immunogenicity of BNT162b2 and CoronaVac COVID-19 vaccines in Hong Kong. *Respirology* (2022) 27(4):301–10. doi: 10.1111/resp.14191
48. Cao Y, Hao X, Wang X, Wu Q, Song R, Zhao D, et al. Humoral immunogenicity and reactivity of CoronaVac or ZF2001 booster after two doses of inactivated vaccine. *Cell Res* (2022) 32(1):107–9. doi: 10.1038/s41422-021-00596-5
49. Costa Clemens SA, Weckx L, Clemens R, Almeida Mendes AV, Ramos Souza A, Silveira MBV, et al. Heterologous versus homologous COVID-19 booster vaccination in previous recipients of two doses of CoronaVac COVID-19 vaccine in Brazil (RHH-001): a phase 4, non-inferiority, single blind, randomised study. *Lancet* (2022) 399(10324):521–9. doi: 10.1016/S0140-6736(22)00094-0
50. Bates TA, McBride SK, Leier HC, Guzman G, Lyski ZL, Schoen D, et al. Vaccination before or after SARS-CoV-2 infection leads to robust humoral response and antibodies that effectively neutralize variants. *Sci Immunol* (2022) 7(68):eabn8014. doi: 10.1126/sciimmunol.abn8014
51. Falsey AR, Frenck RW, Walsh EE, Kitchin N, Absalon J, Gurtman A, et al. SARS-CoV-2 neutralization with BNT162b2 vaccine dose 3. *N Engl J Med* (2021) 385(17):1627–9. doi: 10.1056/NEJMc2113468
52. Eliakim-Raz N, Leibovici-Weisman Y, Stemmer A, Ness A, Awwad M, Ghantous N, et al. Antibody titers before and after a third dose of the SARS-CoV-2 BNT162b2 vaccine in adults aged ≥ 60 years. *JAMA* (2021) 326(21):2203–4. doi: 10.1001/jama.2021.19885
53. Choi A, Koch M, Wu K, Chu L, Ma LZ, Hill A, et al. Safety and immunogenicity of SARS-CoV-2 variant mRNA vaccine boosters in healthy adults: an interim analysis. *Nat Med* (2021) 27(11):2025–31. doi: 10.1038/s41591-021-01527-y
54. Hu J, Peng P, Wang K, Fang L, Luo Fy, Jin As, et al. Emerging SARS-CoV-2 variants reduce neutralization sensitivity to convalescent sera and monoclonal antibodies. *Cell Mol Immunol* (2021) 18(4):1061–3. doi: 10.1038/s41423-021-00648-1
55. Chen M, Qin R, Jiang M, Yang Z, Wen W, Li J. Clinical applications of detecting IgG, IgM or IgA antibody for the diagnosis of COVID-19: a meta-analysis and systematic review. *Int J Infect Diseases* (2021) 104:415–22. doi: 10.1016/j.ijid.2021.01.016
56. Willett BJ, Grove J, MacLean OA, Wilkie C, De Lorenzo G, Furnon W, et al. SARS-CoV-2 omicron is an immune escape variant with an altered cell entry pathway. *Nat Microbiol* (2022) 7(8):1161–79. doi: 10.1038/s41564-022-01143-7
57. Pérez-Then E, Lucas C, Monteiro VS, Miric M, Brache V, Cochon L, et al. Neutralizing antibodies against the SARS-CoV-2 delta and omicron variants following heterologous CoronaVac plus BNT162b2 booster vaccination. *Nat Med* (2022) 28(3):481–5. doi: 10.1038/s41591-022-01705-6
58. Wang Q, Bowen A, Valdez R, Gherasim C, Gordon A, Liu L, et al. Antibody response to omicron BA.4–BA.5 bivalent booster. *New Engl J Med* (2023) 388(6):567–9. doi: 10.1056/NEJMc2213907
59. Keeling MJ. Patterns of reported infection and reinfection of SARS-CoV-2 in England. *J Theor Biol* (2023) 556:111299. doi: 10.1016/j.jtbi.2022.111299
60. Carazo S, Skowronski DM, Brisson M, Barkati S, Sauvageau C, Brousseau N, et al. Protection against omicron (B.1.1.529) BA.2 reinfection conferred by primary omicron BA.1 or pre-omicron SARS-CoV-2 infection among health-care workers with and without mRNA vaccination: a test-negative case-control study. *Lancet Infect Dis* (2023) 23(1):45–55. doi: 10.1016/S1473-3099(22)00578-3
61. Tillet RL, Sevinsky JR, Hartley PD, Kerwin H, Crawford N, Gorzalski A, et al. Genomic evidence for reinfection with SARS-CoV-2: a case study. *Lancet Infect Dis* (2021) 21(1):52–8. doi: 10.1016/S1473-3099(20)30764-7
62. Nonaka CKV, Franco MM, Gráf T, De Lorenzo Barcia CA, De Ávila Mendonça RN, De Sousa KAF, et al. Genomic evidence of SARS-CoV-2 reinfection involving E484K spike mutation, Brazil. *Emerg Infect Dis* (2021) 27(5):1522–4. doi: 10.3201/eid2705.210191
63. Yu ALF, Liphau BL, Ferreira PM, Tanamachi AT, Masuda ET, Trevisan CM, et al. SARS-CoV-2 reinfection: report of two cases in southeast Brazil. *Rev Inst Med Trop Sao Paulo* (2021) 63. doi: 10.1590/s1678-9946202163050
64. Camargo CH, Gonçalves CR, Pagnoca EVRG, Campos KR, Montanha JOM, Flores MNP, et al. SARS-CoV-2 reinfection in a healthcare professional in inner sao paulo during the first wave of COVID-19 in Brazil. *Diagn Microbiol Infect Dis* (2021) 101(4):115516. doi: 10.1016/j.diagmicrobio.2021.115516
65. Lumley SF, Rodger G, Constantinides B, Sanderson N, Chau KK, Street TL, et al. An observational cohort study on the incidence of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and B.1.1.7 variant infection in healthcare workers by antibody and vaccination status. *Clin Infect Dis* (2022) 74(7):1208–19. doi: 10.1093/cid/ciab608
66. Helfand M, Fiordalisi C, Wiedrick J, Ramsey KL, Armstrong C, Gean E, et al. Risk for reinfection after SARS-CoV-2: a living, rapid review for American college of physicians practice points on the role of the antibody response in conferring immunity following SARS-CoV-2 infection. *Ann Intern Med* (2022) 175(4):547–55. doi: 10.7326/M21-4245
67. Murchu E O, Byrne P, Carty PG, De Gascun C, Keogan M, O'Neill M, et al. Quantifying the risk of SARS-CoV-2 reinfection over time. *Rev Med Virol* (2022) 32(1):e2260. doi: 10.1002/rmv.2260
68. Abu-Raddad LJ, Chemaitelly H, Ayoub HH, Yassine HM, Benslimane FM, Al Khatib HA, et al. Association of prior SARS-CoV-2 infection with risk of breakthrough infection following mRNA vaccination in Qatar. *JAMA* (2021) 326(19):1930–9. doi: 10.1001/jama.2021.19623
69. Bean DJ, Monroe J, Turcinovic J, Moreau Y, Connor JH, Sagar M. Severe acute respiratory syndrome coronavirus 2 reinfection associates with unstable housing and occurs in the presence of antibodies. *Clin Infect Dis* (2022) 75(1):e208–15. doi: 10.1093/cid/ciab940
70. Monroe JM, Haralambieva IH, Warner ND, Grill DE, Quach HQ, Kennedy RB. Longitudinal antibody titer, avidity, and neutralizing responses after SARS-CoV-2 infection. *Heliyon* (2022) 8(11):e11676. doi: 10.1016/j.heliyon.2022.e11676
71. Karim SSA, Karim QA. Omicron SARS-CoV-2 variant: a new chapter in the COVID-19 pandemic. *Lancet* (2021) 398(10317):2126–8. doi: 10.1016/S0140-6736(21)02758-6
72. Vahidy FS, Pan AP, Ahnstedt H, Munshi Y, Choi HA, Tiruneh Y, et al. Sex differences in susceptibility, severity, and outcomes of coronavirus disease 2019: cross-sectional analysis from a diverse US metropolitan area. *PLoS One* (2021) 16(1):e0245556. doi: 10.1371/journal.pone.0245556
73. Petersen MS, Pérez-Alós L, Armenteros JJA, Hansen CB, Fjallsbak JP, Larsen S, et al. Factors influencing the immune response over 15 months after SARS-CoV-2 infection: a longitudinal population-wide study in the faroe islands. *J Intern Med* (2023) 293(1):63–81. doi: 10.1111/joim.13560
74. Bates TA, Leier HC, Lyski ZL, Goodman JR, Curlin ME, Messer WB, et al. Age-dependent neutralization of SARS-CoV-2 and P.1 variant by vaccine immune serum samples. *JAMA* (2021) 326(9):868–9. doi: 10.1001/jama.2021.11656
75. Lawandi A, Warner S, Sun J, Demirkale CY, Danner RL, Klompas M, et al. Suspected severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) reinfections: incidence, predictors, and healthcare use among patients at 238 US

healthcare facilities, 1 June 2020 to 28 February 2021. *Clin Infect Dis* (2022) 74(8):1489–92. doi: 10.1093/cid/ciab671

76. Shui TJ, Li C, Liu Hb, Chen X, Zhang B. Characteristics of recovered COVID-19 patients with recurrent positive RT-PCR findings in wuhan, China: a retrospective study. *BMC Infect Dis* (2020) 20(1):749. doi: 10.1186/s12879-020-05463-z

77. Svartz A, Rial MJ, Insúa C, Stedile G, Haleblan E, Peverini A, et al. Brief report 1. *Arch Argent Pediatr* (2023) 121(1):202202595. doi: 10.5546/aap.2022-02595.eng

78. Hansen CH, Michlmayr D, Gubbels SM, Mølbak K, Ethelberg S. Assessment of protection against reinfection with SARS-CoV-2 among 4 million PCR-tested individuals in Denmark in 2020: a population-level observational study. *Lancet* (2021) 397(10280):1204–12. doi: 10.1016/S0140-6736(21)00575-4

79. Xu X, Nie S, Wang Y, Long Q, Zhu H, Zhang X, et al. Dynamics of neutralizing antibody responses to SARS-CoV-2 in patients with COVID-19: an observational study. *Signal Transduct Target Ther* (2021) 6(1):197. doi: 10.1038/s41392-021-00611-6

80. Yang HS, Costa V, Racine-Brzostek SE, Acker KP, Yee J, Chen Z, et al. Association of age with SARS-CoV-2 antibody response. *JAMA Netw Open* (2021) 4(3):e214302. doi: 10.1001/jamanetworkopen.2021.4302

81. Liu Y, Yan LM, Wan L, Xiang TX, Le A, Liu JM, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* (2020) 20(6):656–7. doi: 10.1016/S1473-3099(20)30232-2

82. Wang Y, Zhang L, Sang L, Ye F, Ruan S, Zhong B, et al. Kinetics of viral load and antibody response in relation to COVID-19 severity. *J Clin Invest* (2020) 130(10):5235–44. doi: 10.1172/JCI138759

83. Dadras O, Afsahi AM, Pashaei Z, Mojdeganlou H, Karimi A, Habibi P, et al. The relationship between COVID-19 viral load and disease severity: a systematic review. *Immun Inflammation Dis* (2022) 10(3):e580. doi: 10.1002/iid3.580

84. Menni C, Valdes AM, Polidori L, Antonelli M, Penamakuri S, Noyal A, et al. Symptom prevalence, duration, and risk of hospital admission in individuals infected with SARS-CoV-2 during periods of omicron and delta variant dominance: a prospective observational study from the ZOE COVID study. *Lancet* (2022) 399(10335):1618–24. doi: 10.1016/S0140-6736(22)00327-0

85. Vihta KD, Pouwels KB, Peto TE, Pritchard E, House T, Studley R, et al. Omicron-associated changes in SARS-CoV-2 symptoms in the united kingdom. *Clin Infect Dis* (2022) 76(3):e133–e141. doi: 10.1101/2022.01.18.22269082

86. Machado-Curbelo C, Gutiérrez-Gil J, González-Quevedo A. A shift in SARS-CoV-2 omicron variant's entry pathway might explain different clinical outcomes. *MEDICC Rev* (2022) 24(3–4):68–71. doi: 10.37757/MR2022.V24.N3-4.10

87. Kojima N, Klausner JD. Protective immunity after recovery from SARS-CoV-2 infection. *Lancet Infect Dis* (2022) 22(1):12–4. doi: 10.1016/S1473-3099(21)00676-9

88. Babouee Flury B, Güsewell S, Egger T, Leal O, Brucher A, Lemmenmeier E, et al. Risk and symptoms of COVID-19 in health professionals according to baseline immune status and booster vaccination during the delta and omicron waves in Switzerland—a multicentre cohort study. *PLoS Med* (2022) 19(11):e1004125. doi: 10.1371/journal.pmed.1004125

89. Accorsi EK, Britton A, Fleming-Dutra KE, Smith ZR, Shang N, Derado G, et al. Association between 3 doses of mRNA COVID-19 vaccine and symptomatic infection caused by the SARS-CoV-2 omicron and delta variants. *JAMA* (2022) 327(7):639–51. doi: 10.1001/jama.2022.0470

90. Patalon T, Saciuk Y, Peretz A, Perez G, Lurie Y, Maor Y, et al. Waning effectiveness of the third dose of the BNT162b2 mRNA COVID-19 vaccine. *Nat Commun* (2022) 13(1):3203. doi: 10.1038/s41467-022-30884-6