

## IX – REFERÊNCIAS BIBLIOGRÁFICAS

- ABREU, S.F. (1965). *Recursos minerais do Brasil*, vol. I (minerais não-metálicos). Rio de Janeiro, 489p.
- ADDAD, J. (2001). *Minerais pesados: uma ferramenta para prospecção, proveniência, paleogeografia e análise ambiental*. São Paulo: edição independente, 80p.
- ADDAD, J., KARFUNKEL, J., RENGER, F., QUÉMÉNEUR, J., MARTINS, M. & SCHOLZ, R. (1999). Os extremos nordeste e sudeste da protobacia sanfranciscana em Minas Gerais: A serra do Pai Albano e o alto do Cascabulho. *Bol. do 5º Simp. sobre o cretáceo do Brasil*. 189-193.
- ALKMIM, F.F. (2004). O que faz de cráton um cráton? O Cráton do São Francisco e as revelações Almeidianas ao delimitá-lo. In: V. Mantesso Neto, A. Bartorelli, C. Dal Ré Carneiro e B. Brito Neves (orgs.) *Geologia do Continente Sul-Americano: evolução da obra de Fernando Flávio Marques Almeida*. São Paulo; Editora Beca, 17-35.
- ALKMIM, F.F., CHEMALE Jr., F., ENDO, I. (1996). A deformação das coberturas proterozóicas do Cráton do São Francisco e o seu significado tectônico. *Rev. Escola de Minas*, **49**(1):22-38.
- ALKMIM, F.F., BRITO NEVES, B.B. & CASTRO, J.A. (1993). Arcabouço tectônico do Cráton do São Francisco – Uma revisão. In: O Cráton do São Francisco. Reunião Preparatória do II Simp. Sobre o Cráton do São Francisco. J.M.L. DOMINGUES e A. MISI (eds.). *Soc. Bras. Geol. Minas e Energia*. CNPq. Salvador, 45-62.
- ALKMIM, F.F., MARSHAK, S., FONSECA, M.A. (2001). Assembling West Gondwana in the Neoproterozoic: clues from São Francisco craton region, Brazil. *Geology*, **29**:319-322.
- ALLEN, P.A., ALLEN, J.R. (1990). *Basin analysis: Principles and applications*. Blackwell, Oxford, 415p.
- ALMEIDA, F.F. (1968). Origem e evolução da Plataforma Brasileira. DNPM/DGM, Bol. 241, 36p, Rio de Janeiro.
- ALMEIDA, F.F., HASUI, Y., BRITO NEVES, B.B. (1976). The Upper Precambrian of South América. *Bol. Inst. Geoc.*, USP/IG, **7**:45-80.
- ALMEIDA, F.F.M. (1977). O Cráton do São Francisco – *Rev. Bras. Geol.* **7**(4): 349-364.
- ALMEIDA, F.F.M. & HASUI, Y. (1984). *O Pré-cambriano do Brasil*. Editora Blücher, São Paulo, 604p.
- ALMEIDA-ABREU, P.A. (1993). *A evolução geodinâmica da Serra do Espinhaço Meridional, Minas Gerais, Brasil*. Tese de doutoramento, Univ. Freiburg, 150p.
- ALMEIDA-ABREU, P.A. (1995). O Supergrupo Espinhaço da Serra do Espinhaço Meridional. Minas Gerais: o rifte, a bacia e o orógeno. *Geonomos*, **3**(1): 1-18.
- ALMEIDA-ABREU, P.A. & PFLUG, R. (1994). The geodynamic evolution of the southern Serra do Espinhaço, Minas Gerais, Brazil. *Zbl., Geol., Palaönt., Teil I*, (1/2): 21-44.
- ALMEIDA-ABREU, P.A. & RENGER, F. (2000). Paleogeography and significance of the Mesoproterozoic mountain glaciation within the Serra do Espinhaço, Minas Gerais, Brasil. *31<sup>TH</sup> Intern. Cong. Geol. Rio de Janeiro. Abstracts*.
- ALMEIDA-ABREU, P.A. & RENGER, F. (2002). Serra do Espinhaço meridional: um orógeno de colisão Mesoproterozóico. *Rev. Bras. Geoc.*, **32**/1:1-14.
- ANDERSON, D.L. (1980). Chemical stratification of the mantle. *Jour. of Geophys. Res. (suplly)*, **84**:6297-6298.
- ARAÚJO, D.P. (2002). *Mineralogia dos diamantes da Província kimberlítica de Juína, MT*. Tese de doutoramento, IG/UnB, 202p.
- ARIMA, M., KOZAI, Y., AKAISHI, M. (2002). Diamond nucleation and growth by reduction of carbonate melts under high-pressure and high-temperature conditions. *Geology*, **30**:691-694.
- ASHLEY, G.M. (1990). Classification of a large-scale subaqueous bedforms: a new at an old problem. *Jour. Sedim. Petrol.* **60**:160-172.
- ASSUNÇÃO, C.T. (1964). Temas clássicos em cristalografia. Classes de simetria e formas simples. *Soc. Portuguesa de Ciências Naturais, Coleção "Natura"*, 56p.
- BABINSKI, M. & KAUFMAN, A.J. (2003). First direct dating of a Neoproterozoic post-glacial cap carbonate. In: IV South American Symposium on Isotope Geology, Cordoba, Short Papers, 321-323.
- BANKO, A.G. (1997). *FTIR-Spektroskopische Untersuchungen na Beryllen unterschiedlicher Herkunft, sowie die Geologische, Mineralogische und Gemmologische Charakterisierung der Diamanten des Espinhaço-Gebirges (Minas Gerais, Brasilien)*. Dissertation am Institut für Mineralogie und Kristallographie der Universität Wien, 186p.
- BARBOSA, O. (1965). Geologia econômica e aplicada a uma parte do planalto brasileiro. In: Cong. Bras. Geol., 19, Rio de Janeiro, 1965. Anais..., RJ, SBG.

- BARCELOS, J.H. & SUGUIO, K. (1980). Ambiente de sedimentação da Formação Areado, Cretáceo Inferior da bacia Sanfranciscana, MG. *Rev. Bras., Geoc.*, **10**(4):237-242.
- BARTOSHINSKY, Z.V., BEKESHA, S.N., VINNICHENKO, T.G. (1987)a. Volatiles in diamonds from Northern part of Russian Platform. *Mineralogical collected book*, **44**(2):14-18 (em russo).
- BARTOSHINSKY, Z.V., BEKESHA, S.N., VINNICHENKO, T.G. (1987)b. Gas impurities of Yakutian diamonds. *Mineralogical collection*, **41**(1):25-32.
- BERGER, S.D. & PENNYCOOK, S.J. (1982). Detection of nitrogen at {100} platelets in diamond. *Nature*, **298**:635-637.
- BERMAN, R. (1965). Physical properties of diamonds. *Oxford*, Clarendon Press. 443p.
- BOUMA, A.M. (1962). *Sedimentology of some flysch deposits*. Amsterdam, Elsevier, 168p.
- BOYD, F.R., & FINNERTY, A.A. (1980). Conditions of origin of natural diamonds of peridotitic affinity. *Jour. Geophys. Res.*, **85**:6911-6918.
- BOYD, S.R., & PILLINGER, C.T. (1994). A preliminary study of  $^{15}\text{N}/^{14}\text{N}$  in octahedral growth from diamonds. *Chem. Geol.*, **116**:43-59.
- BRANCO, J.J.R. & COSTA, M.T. (1961). Roteiro para a excursão Belo Horizonte-Brasília. UFMG/IPR, Publ. 15, 127p.
- BRANNER, J.C. (1919). Outline of the geology of Brazil to accompany the geological map of Brazil. *Bulletin of the Geological Society of America*, **30**:189-338.
- BRITO NEVES, B.B., CORDANI, U.G., KAWASHITA, K., DELHAL, J. (1979). A evolução geocronológica da Cordilheira do Espinhaço – dados novos e integração. *Rev Bras. Geoc.*, **9**(1): 71-85.
- BRITO NEVES, B.B., WINGE, M., CARNEIRO, M.A. ((1996). Orogênese precedendo e tafrogêneses sucedendo Rodínia na América do Sul. *Boletim IG-USP*, **27**:1-40.
- BRODZIKOWSKI, K. & van LOON, A.J. (1987). *Glacigenic sediments*. Amsterdam, Elsevier. 674p.
- BRULEY, J. & BROWN, L.M. (1989). Quantitative electron energy-loss spectroscopy microanalysis of platelets and voidite defects in natural diamonds. *Phil. Magazine*, **59**:247-261.
- BUCHWALDT, R., TOULKERIDIS, T., BABINSKI, M., SANTOS, R., NOCE, C.M., MARTINS-NETO, M. & HERCOS (1999). Age determination and age related provenance analysis of the Proterozoic glaciation event in central eastern Brazil. *In: South America Symposium on Isotope Geology*, 2. Cordoba, 1999. *Anais 4*, p. 387-390.
- BULANOVA, G.P. (1995). The formation of diamond. *Journal of Geochemical Exploration*, **53**:1-23.
- BULANOVA, G.P., PEARSON, D.G., HAURI, E.H., GRIFFIN, B.J. (2002). Carbon and nitrogen isotope systematics within a sector-growth diamond from the Mir kimberlite, Yakutia. *Chem. Geol.*, **188**:105-123.
- BURNS, R.C. & DAVIES, G.J. (1997). Growth of synthetic diamond. *In: The properties of natural and synthetic diamond*. J.E. Field (ed.). Academic Press, 398-421.
- BURSILL, L.A. & GLAISHER, R.W. (1985). Aggregation and dissolution of small and extended defect structures in type Ia diamond. *Am. Mineral.*, **70**:608-618.
- CAMPOS, J.E.G., DARDENNE, M.A. & GONZAGA, G.M. (1993). O potencial diamantífero do conglomerado Abaeté no NW de Minas Gerais. *In: Simp. Geol. Diamante. I. Anais. Cuiabá. UFMT, Publ. Esp. 2/93*:101-113.
- CAMPOS, J.E.G. & GONZAGA, G.M. (1999). O transporte de diamantes por sistemas fluviais a longas distâncias: uma visão crítica. *Revista Brasileira de Geociências*. **29**(2):113-118.
- CAMPOS NETO, M.C. (2000). Orogenic systems from southwestern Gondwana. An approach to Brasiliano-Pan Africano cycle and orogenic collage in southeastern Brazil. *In: U.G. Cordani, E.J. Milani, A. Thomaz Filho, D.A. Campos (eds.). Tectonic evolution of South America*. 31<sup>TH</sup> Int. Geol. Cong., Rio de Janeiro, 335-365.
- CAPDEVILA, R., ARNDT, N., LETENDRE, J., SAUVAGE, J-F. (1999). Diamonds in volcanoclastic komatiite from French Guiana. *Nature*, **399**:456-458.
- CARLISLE, D.B. & BRAMAN, D. (1991). Nanometre-size diamonds in the cretaceous/tertiary boundary clay of Alberta. *Nature*. **352**:708-709.
- CARTIGNY, P., HARRIS, J.W. & JAVOY, M. (1998)a. Eclogitic diamond formation at Jwaneng.. *Science*. **280**:1421-1424.
- CARTIGNY, P., HARRIS, J.W., PHILLIPS, D., GIRARD, M., JAVOY, M. (1998)b. Subduction-related diamonds? The evidence for a mantle-derived origin from coupled  $\delta^{13}\text{C}$ -  $\delta^{15}\text{N}$  determinations. *Chem Geol.*, **147**:147-159.
- CARTIGNY, P., HARRIS, J.W. & JAVOY, M. (1999). Eclogitic, peridotitic and metamorphic diamonds and problems of carbon recycling – The case

- of Orapa (Botswana). In: *Proc. VII Conf. Intern. Kimberlitos* (ed. Gurney, J.J., Gurney, J.L., Pascoe, M.D. e Richardson, S.H.). África do Sul, Cidade do Cabo, **1**:117-124.
- CARTIGNY, P., HARRIS, J.W. & JAVOY, M. (2001). Diamond genesis, mantle fractionations and mantle nitrogen content: a study of  $\delta^{13}\text{C-N}$  concentrations in diamonds. *Earth and Plan. Science Letters*, **185**:85-98.
- CENSIER, C. & TOURENQ, J. (1995). Crystal forms and surface textures of alluvial diamonds from the western region of the Central African Republic. *Mineralium Deposita*. **30**:314-322.
- CHAVES, A.O., RIGHI, A., NEVES, J.M.C., HARTMAN, M., MARTINS, M. (2005)a. Nitrogen centres and mineral inclusions of diamonds from Minas Gerais (Brazil) as revealed by vibrational spectroscopy. In: *II South American Symp. On Diamond Geology*, Diamantina, 43-48.
- CHAVES, M.L.S.C. (1987). Conglomerados diamantíferos da região do rio Salobro, Canavieiras, BA. *SBG/MG, Boletim* **6**:155-162. Sistemas deposicionais no Precambriano.
- CHAVES, M.L.S.C. (1997). *Geologia, mineralogia e gemologia do diamante da Serra do Espinhaço em Minas Gerais*. Tese de doutoramento, IG-USP, São Paulo. 289p.
- CHAVES, M.L.S.C. & KARFUNKEL, J. (1997). A protobacia do rio Jequitinhonha (Minas Gerais) durante o cretáceo inferior e seu potencial diamantífero. *Geociências, São Paulo*. **16**(1):191-203.
- CHAVES, M.L.S.C. & SVISERO, D.P. (1993). Características geológicas e origem dos conglomerados diamantíferos das regiões de Diamantina (Mesoproterozóico) e de Romaria (Cretáceo Superior), MG. *Boletim IG/USP, Série Científica*, **24**:49-57.
- CHAVES, M.L.S.C. & UHLEIN, A. (1991). Depósitos diamantíferos da região do Alto/Médio Rio Jequitinhonha, Minas Gerais. In: C. Schobbenhaus, E.T. Queiroz, & C.E.S. Coelho (eds.), *Principais depósitos minerais do Brasil*, v.IV-A. DNPM/CPRM, Brasília, 117-138.
- CHAVES, M.L.S.C., DUPONT, H., KARFUNKEL, J., SVISERO, D.P. (1993). Depósitos diamantíferos de Minas Gerais: uma revisão. *Anais do 1º. Simpósio Brasileiro de Geologia do Diamante*, Cuiabá, **1**:79-100.
- CHAVES, M.L.S.C., GERALDES, M.C., SANO, Y., CHAMBEL, L. (2005)b. Primeiros resultados da datação Pb-Pb em diamantes carbonados da Chapada Diamantina (BA) e Rio Jequitinhonha (MG), Serra do Espinhaço. *Rev Bras. Geociências*, **35**(3), 419-422.
- CHAVES, M.L.S.C., KARFUNKEL, J. & ADDAD, J. (1999). Geologia da região diamantífera de Grão Mogol, Minas Gerais. *Geociências, São Paulo*. **18**(1): 129-155.
- CHAVES, M.L.S.C., KARFUNKEL, J., HOPPE, A., HOOVER, D.B. (2001). Diamonds from the Espinhaço Range (Brazil) and their redistribution through the geologic record. *J. South Am. Earth Sci.*, **14**:277-289.
- CHAVES, M.L.S.C., KARFUNKEL, J., RIBEIRO, S.H. (1996). A protobacia do Rio Jequitinhonha durante o Eocretáceo e seu potencial diamantífero. In: *simp. Sobre o cretáceo do Brasil*, 4, Águas de São Pedro. Bol. do...Rio Claro, UNESP, p.195-198.
- CHAVES, M.L.S.C., KARFUNKEL, J., MARTINS, M.S., OLIVEIRA, L.H.B. (1998). O conglomerado diamantífero da Formação Grão Mogol na área do Papo D'Emá (Grão Mogol). In: *Cong. Bras. Geol.*, 1, Belo Horizonte. *Anais*. Belo Horizonte, p.290.
- CHAVES, M.L.S.C., KARFUNKEL, J., STASIULEVICIUS, R., BRANDÃO, P.R.G., CHAMBEL, L. (2001). Color nature of green and brown (coated and body color) diamonds in SE-Brazil (Minas Gerais) and Angola (Lunda): a discussion. In: *3 Simp. Bras. Geol. do Diamante / 1 Simp. Sul-Am. Geol. do Diamante. Abstracts*. p.14, Brasília-DF.
- CHAVES, M.L.S.C., KARFUNKEL, J., WEGNER, R. (1997). Diamantvorkomen von Grão Mogol: Geologische end gemologische untersuchungen in Minas Gerais, Brasilien. *Geowissenschaften*, **15**:73-78.
- CHEPUKROV, A.I., FEDOROV, I.I., SONIN, V.M. (1997). Experimental modelling of the diamond formation process. *Novosibirsk, Siberian Branch*, 197p. (em russo).
- CHOPIN, C. (1984). Coesite and pure pyrope in high-grade blueschists of the western Alps: a first Record and some consequences. *Contributions to Mineralogy and Petrology*, **86**:107-118.
- CHULA, A.M.D. (1996). *Caracterização e geoquímica dos metamagmatitos e metassedimentos da região de Planalto de Minas, Diamantina, Minas Gerais*. IGC/UFMG, Dis. mestrado, 162p.
- CLIFFORD, T.N. (1966). Tectono-metalogenetic units and metallogenetic provinces of Africa. *Earth and Planetary Science Letters*, **1**: 421-434.
- CLOIZEAUX, A.L.D. (1855). Note sur le diamante noir. *Annales des Mines (Paris)*, 5 ed., **8**:304/6.
- COLLINSON, D.W. (1998). Magnetic properties of polycrystalline diamonds. *Earth and Planetary Science Letters*. **161**:179-188.

- COLLINSON, J.D. & THOMPSON, D.B. (1982). *Sedimentary structures*. George Allen & Unwin (publ.), 194p.
- COMPSTON, W., WILLIAMS, I.S., KIRSCHVINK, J.L. (1992). Zircon U–Pb ages of early Cambrian time-scale. *J. Geol. Soc.* **149**:171–184.
- CORDANI, U.G.; SATO, K.; TEIXEIRA, W.; TASSINARI, C.C.G & BASEI, M.A.S. (2000). Crustal evolution of the South American Platform. In: CORDANI, U.G.; MILANI, E.J.; THOMAZ FILHO, A.; CAMPOS, D.A. (eds) *Tectonic evolution of South America*, Rio de Janeiro, 19–40.
- COX, K.C., BELL, J.D., PANKHURST, R.J. (1979). *The interpretation of igneous rock*. London, Allen and Unwin, 450p.
- CRAIG, H. (1953). The geochemistry of the stable carbon isotopes. *Geochim. Cosmochim Acta*, **3**:53–92.
- CUKROV, N. (1999). *A glaciação neoproterozóica na porção sul do Cráton do São Francisco e suas litofácies nas regiões de Jequiá-MG e Cristalina-GO*. Dis. Mestrado, IG/UnB, 104p.
- D'AGRELLA-FILHO, M.S., PACCA, I.G., TEIXEIRA, W., ONSTOTT, T.C. & RENNE, P.R. (1990). Paleomagnetic evidence for the evolution of meso to neoproterozoic glaciogenic rocks in central-eastern, Brazil. In: *Palaeogeogr., Palaeoclim., Palaeocol.*, **80**:255–265.
- DANDERFER FILHO, A. (2000). *Geologia sedimentar e evolução tectônica do Espinhaço Setentrional, Estado da Bahia*. Tese de doutorado, UnB, Brasília, 498p.
- DARDENNE, M.A. (1978). Síntese sobre a estratigrafia do Grupo Bambuí no Brasil Central. In: XXX Cong. Bras. Geol., *Anais...*, **2**:597–610, Recife.
- DARDENNE, M.A. (1979). *Lês mineralizations de plomb, Zinc, Flúor du Proterozoïque Superior dans lê Brasil Central. Avec une mise an point sur la cronologie du Précambrien Brésilien*. Tese de doutorado, Univ. Paris, 251p.
- DARDENNE, M.A., & SCOBENHAUS, C. (2000). Crustal evolution of the South American Platform. In: CORDANI, U.G.; MILANI, E.J.; THOMAZ FILHO, A.; CAMPOS, D.A. (eds) *Tectonic evolution of South America*, Rio de Janeiro, p.19–40.
- DARDENNE, M.A., & SCOBENHAUS, C. (Eds.) (2001). *Metalogênese do Brasil*. Brasília - Editora Universidade de Brasília, 392p.
- DARDENNE, M.A., & WALDE, D.G. (1979). Estratigrafia dos grupos Bambuí e Macaúbas no Brasil Central. In: *I Simp. Geol. Minas Gerais*, *Anais*, 43–53.
- DAULTON, T.L. & OZIMA, M. (1996). Radiation-induced diamond formation in uranium-rich carbonaceous materials. *Science*. **271**:1260–1262.
- DAVIES, G. (1984). *Diamond*. Adam Hilder Ltd., Bristol, UK, 255p.
- DAWSON, J.B. (1980). *Kimberlites and their xenoliths*. Springer-Verlag, Berlin, 324p.
- DAWSON, J.B. (1985)a. Ascent and emplacement of kimberlite magma. In: *Kimberlite occurrence and origin: a basics for conceptual models in exploration*. J.E. Glover & P.G. Harris. **8**:113–124.
- DAWSON, J.B. (1985)b. Petrogenesis of kimberlite. In: *Kimberlite occurrence and origin: a basics for conceptual models in exploration*. J.E. Glover & P.G. Harris. **8**:103–111.
- DAWSON, J.B. & STEPHENS, W.E. (1976). Statistical classification of garnets from kimberlites and associated xenoliths. *Journal of Geology*, **83**:589–607.
- DE, S., HEANEY, P.J., HARGRAVES, R.B., VICENZI, E.P. & TAYLOR, P.T. (1998). Microstructural observations of polycrystalline diamond: a contribution to carbonado conundrum. *Earth and Planetary Science Letters*. **164**:421–433.
- DE, S., HEANEY, P.J., VICENZI, E.P. & WANG, J. (2001). Chemical heterogeneity in carbonado, an enigmatic polycrystalline diamond. *Earth and Plan. Science Letters*. **185**:315–330.
- DEINES, P. (1980). The carbon isotopic composition of diamonds: relationship to diamond shape, color, occurrences and vapor composition. *Geochim. Cosmochim Acta*, **44**:943–961.
- DEINES, P. (1992). Mantle carbon: concentration, mode of occurrence and isotopic composition. In: *Schidrowski, M., Golubic, S., Kimberley, M.M., McKirdy, D.M., Trudinger, E.A. (eds.)* Early organic evolution. Springer, Berlin, 133–146p.
- DEINES, P. (2002). The carbon isotope geochemistry of mantle xenoliths. *Earth Sci. Rev.*, **58**:247–278.
- DEINES, P., HARRIS, J.W., GURNEY, J.J. (1987). Carbon isotopic composition, nitrogen content and inclusion composition of diamonds from the Roberts Victor kimberlite, South Africa: evidence for <sup>13</sup>C deposition in the mantle. *Geochim. Cosmochim Acta*, **51**:1227–1243.
- DEINES, P., VILJOEN, F., HARRIS, J.W. (2001). Implications of the carbon isotope and mineral inclusion record for the formation of diamonds in the mantle underlying a mobile belt: Venetia, South Africa. *Geochim. Cosmochim Acta*, **65**:813–838.

- DEINES, P. (2002). The carbon isotope geochemistry of mantle xenoliths. *Earth Science Review*, **58**:247-278.
- DEINES, P. & HARRIS, J.W. (1994). On the importance of fluids for diamond growth. In: *Goldschmit Conf. (ext. abstract)*, Edimburgh, 219-220.
- DERBY, O.A. (1879). Observações sobre algumas rochas diamantíferas da Província de Minas Gerais – *Arch. Mus. Nac.*, IV, 121-132, Rio de Janeiro.
- DERBY, O.A. (1881). Observações sobre algumas rochas diamantíferas da província de Minas Gerais. *Arch., Mus. Nac.*, 4: 121-132; Rio de Janeiro.
- DERBY, O.A. (1882). Modes of occurrence of the diamond in Brazil – *Amer. Jour. Sci., 3rd. Ser.*, Vol. XXIV, p. 34-42, New Haven.
- DERBY, O.A. (1905a). The geology of the diamond and carbonado washings of Bahia, Brazil – *Econ. Geol.*, **1**:134-142, Urbana.
- DERBY, O. A. (1905b). Lavras diamantinas. *Revista Instituto Geográfico e Histórico da Bahia*, **11**(30):143-153.
- DERBY, O.A. (1906). The Serra do Espinhaço, Brazil – *Jour. Geol.*, **14**:374-401, Chicago.
- DOSSIN, I.A., UHLEIN, A., DOSSIN, T.M. (1984). Geologia da Faixa Móvel Espinhaço em sua porção meridional, MG. In: XXXIII Cong. Bras. Geol., Anais **7**: 3118-3134. Rio de Janeiro.
- DOSSIN, I.A., DOSSIN, T.M., CHAVES, M.L.S.C. (1990). Compartimentação estratigráfica do Supergrupo Espinhaço em Minas Gerais – Os grupos Diamantina e Conselheiro Mata. *Revista Bras. Geociências*, **20**(1-4): 178-186.
- DUPONT, H. (1995). O Grupo Conselheiro Mata no seu quadro paleogeográfico e estratigráfico. In: Simp. Geol. Minas Gerais, 8. Diamantina, 1995. Anais. Belo Horizonte, SBG-Núcleo MG, 9-10pg.
- DUPONT, H., ESPINOSA, J., KARFUNKEL, J., & NOCE, C.M. (2000). Glacial deposits of the neoproterozoic glaciation in the Serra do Cabral na vicinities (Minas Gerais, Brasil). Facies distribution and genesis. *31<sup>th</sup> Intern. Geol. Cong. Rio de Janeiro. Abstracts*.
- DUPONT, H., KARFUNKEL, J., & SOARES, H.M. (2001). Paleogeografia das coberturas sedimentares neoproterozóicas do Cráton do São Francisco. In: *XI Simp. Geol. Minas Gerais*, em CD-Rom.
- DUSSIN I.A. (1994a). Evolution structurale de l'Espinhaço meridional, bordure sud-est du craton São Francisco, Brésil. Orléans, Tese de doutorado, 200p.
- DUSSIN, T.M. (1994b). Association volcano-plutoniques de l'Espinhaço meridional (SE-Brésil). Orléans, Tese de doutorado, 177p.
- DUSSIN I.A. & DUSSIN, T.M. (1995). Supergrupo Espinhaço: Modelo de Evolução Geodinâmica. *Geonomos*, **1**: 19-26.
- DYER, H.B., RAAL, F.A., PREEZ, L.DU., LOBUSER, H.N. (1965). Optical absorption features associated with paramagnetic nitrogen in diamond. *Mat. Sci. Earth Interior*, I. Sunagawa (ed.):763-774.
- EINSELE, G. (1992). *Sedimentary basins*. Berlin Springer-Verlag, 628p.
- ELDRIDGE, C.S., COMPSTON, W., WILLIAMS, I.S., HARRIS, J.W., BRISTOW, J.W. (1991). Isotope evidence for the involvement of recycled sediments in diamond formation. *Nature*, **353**:243-245.
- ESCHWEGE, W.L. von (1833). Pluto brasiliense – Tradução em português, 2 vol., (Cia. Ed. Nac.), São Paulo, 1944.
- EVANS, T. (1992). Aggregation of nitrogen in diamond. In: *The properties of natural and synthetic diamond* (ed. J.E. Field), Academic Press, 710p.
- EVANS, T. & HARRIS, J.W. (1989). Nitrogen aggregation, inclusion equilibration temperatures and the age of diamonds. *Geology Society Australian, especial publication*. **14**(2):1001.
- EVANS, T. & QI, Z. (1982). The kinetics of the aggregation of nitrogen atoms in diamond. *Proc. R. Soc. London*. **A270**:538-552.
- EVDOKIMOV, M.D., LADYGINA, M.Y., NESTEROV, A.R. (2001). Morphology of diamonds as a possible indicator of their genesis. *N. Jb. Miner. Abh.*, **176**(2):153-177.
- EYLES, N. & EYLES, C.H. (1992). Glacial depositional systems: In R.G. Walker & N.P. James (eds.): *Facies models*. Geotext. 1, *Geol. Assoc. of Canada*, 73-100.
- EYLES, N. (1995). Characteristics and origin of coarse gold in Late Pleistocene sediments of the Cariboo placer mining district, British Columbia, Canada. *Sedim. Geol.*, **95**:69-95.
- FAIRCHILD, I.J. & SPIRO, B. (1990). Carbonate minerals in glacial sediments: geochemical clues to palaeoenvironment. In: *Glacimarine environments: processes and sediments*. J.A. Dowdeswell & J.D. Scourse (eds.). *Geol. Soc. Spec. Publ.* **53**:201-206.
- FIELD, J.E. (1997). *The properties of natural and synthetic diamonds*. Academic Press, Second edition, New York, 710p.

- FIPKE, C., GURNEY, J.J., MOORE, R.O. (1989). The development of advanced technology to distinguish between diamondiferous and barren diatremes. *Geol. Surv. of Canada, Open File Report*, 2124.
- FLEISCHER, R. (1998). A rift model for the sedimentary diamond deposits of Brazil. *Mineralium Deposita*, **33**: 238-254.
- FLINT, R.F. (1962). Geological evidence of cold climates. In: NAIRN, A.E.M. (ed.): *Problems in Paleoclimatology*: 141-155; New York (interscience).
- FOGAÇA, A.C.C., ALMEIDA-ABREU, P.A., SCHORSCHER, H.D. (1984). Estratigrafia da sequência supracrustal arqueana da porção meridiana-central da Serra do Espinhaço Meridional, Minas Gerais. In: XXXIII Cong. Bras. Geol., 6: 2654-2667, Rio de Janeiro.
- FRANK, F.C. & PUTTICK, K.E. (1958). Etch pits and trigons in diamond (1). *Phil. Magazine*. **3**:1262-1272.
- FRANTSESSON, E.V. & KAMINSKIY, F.V. (1974). Carbonado, a diamond variety of nonkimberlitic origin. *Doklady Akademii Nauk SSSR*, **219**:187-189.
- FREYBERG, B. von (1932). Ergebnisse geologischer forschungen in Minas Gerais, Brasilien. *Neves Jahrb. Minel. Geol. Paleontol. Abh. Mineral. Petrogr.*, Stuttgart, (2):403p.
- FREYBERG, B. von (1934). Bodenschätze des staates Minas Gerais (Brasilien). *E. Schweizerbart'sche Verlagsbuchhandlung*. P.306-317.
- FRITSCH, E., & SCARRATT, K. (1992). Natural-color nonconductive gray-to-blue diamonds. *Gems & Gemology*, Spring Issue, 35-42.
- FRONDEL, C. & MARVIN, U.B. (1967). Lonsdaleite, a hexagonal polymorph of diamond. *Nature*, 214:587-589. Blücher, São Paulo, p. 159-208.
- FROSTICK, L.E. & REID, I. (1987). Tectonic control of desert sediments in rift basins ancient and modern. In: L.E. FROSTICK & I. REID (eds.), *Desert Sediments: Ancient and Modern*. **35**:53-68.
- GALIMOV, E.M. (1991). Isotope fractionation related to kimberlitic magmatism and diamond formation. *Geochim. Cosmochim Acta*, **55**:1697-1708.
- GALIMOV, E.M., KAMINSKY, F.V., KODINA, L.A. (1985). Recent data on isotopic composition of carbonado carbon. *Geochemistry*, **5**:723-726.
- GALLOWAY, W.D. & HOBDAV, D.K. (1996). *Terrigenous clastic depositional systems. Applications to fossil fuel and groundwater resources*. 2 ed., Berlim Springer Verlag, 489p.
- GOLUBEV, Yu. K. (1995). Diamond exploration in glaciated terrain: a Russian perspective. *Journal of Geochemical Exploration*, **53**:265-275.
- GONZAGA, G.M., TEIXEIRA, N.A. & GASPAR, J.C. (1994). The origin of diamonds in western Minas Gerais, Brazil. *Mineralium Deposita*. **29**:414-421.
- GONZAGA, G.M. & TOMPKINS L.A. (1991). Principais depósitos minerais do Brasil. Gemas e rochas ornamentais, *DNPM-CPRM*, vol. 4-A, p. 53-116.
- GORESY, A. & DONEY, G. (1968). A new allotropic form of carbon from the Ries crater. *Science*, **161**(3839):363-366.
- GORSHKOV, A.I., BAO YANAN, BERSHOV, L.V., RYABCHIKOV, I.D., SIVTSOV, A.V., LAPINA, M.I. (1997)a. Inclusions in Diamond from the Liaoning Deposit (China) and Their Genetic Meaning. *Geochemistry*. **1**:58-65.
- GORSHKOV, A.I., BERSHOV, L.V., VINOKUROV, S.F., OTON, H.L., SIVTSOV, A.V., MOKHOV, A.V., BOGACHEVA, E.O. (1997)b. Carbonado from the Lençois region, Bahia State, Brazil. Mineral inclusions, physical and geochemical characteristics and conditions for formation. *Geol.Rudn. Mestorozhd.*, 39(3), 269-277, 1997. (Russian).
- GORSHKOV, A.I., TITKOV, S.V., PLESHAKOV, A.M., SIVTSOV, A.V. & BERSHOV, L.V. (1996)a. Inclusions of Native Metals and Other Mineral Phase in Carbonado from Ubanga region (Central Africa)., *Geology of ore Deposits*. V.38. 02. P.131-136.
- GORSHKOV, A.I., TITKOV S.V., SIVTSOV A.V., BERSHOV L.V. (1996)b. Rare-earth Minerals in Carbonado from Yakutian Diamond Deposits. *Geochemistry*. 6. p.501-506.
- GORSHKOV, A.I., TITKOV, S.V., SIVTSOV, A.V., BERSHOV, L.V. & MARFUNIN, A.S.(1996)c. The first finds of native Cr, Ni, and  $\alpha$ -Fe in carbonado from the diamond deposits of Yakutia. *Geochimistry Internation*. Vol. 33, No. 1: p.59-63.
- GORSHKOV, A.I., VINOKUROV, S.F., RYABCHIKOV, I.D., BERSHOV, L.V., MAGAZINA, L.O., SIVTSOV, A.V., SOLODOV, D.I. (2000). Mineralogical and geochemical characteristics of gold-containing carbonado from the Poxoreu District, Mato Grosso State (Brazil). *Geokhimiya*, (1), 3-15 (Russian) 2000.
- GORSHKOV, A.I., VINOKUROV, S.F., SOLODOV, D.I., BERSHOV, L.V., MOKHOV, A.V., SOLODOVA, YU. P., SIVTSOV, A.V. (1998). Polycrystalline diamond from the Udachnaya pipe (Yakutia).Mineralogical, geochemical, and genetic characteristics. *Litol. Polezn.Iskop.*, (6):588-603 1998. (Em russo).

- GORSHKOV, A.I., YAN NAN BAO, BERSHOV, L.V., RYABCHICOV, I.D., SIVTISOV, A.V., LAPINA, M.I. (1997)c. Inclusions of Native Metals and Other Minerals in Diamond from Kimberlite Pipe 50, Liaoning, China. *Geochemistry*, **8**:794-804.
- GRADIM, R. (2003). *Geologia da região de Olhos d'Água (MG)*. Trabalho geológico de graduação, IGC/UFGM, 63p.
- GRAVENOR, C.P., BRUNN, V.V., DREIMANIS, A. (1984). Nature and classification of waterlain glaciogenic sediments, exemplified by Pleistocene, Late Paleozoic and late Precambrian deposits. *Earth science Reviews*, **20**: 105-166.
- GRAVENOR, C.P. & MONTEIRO, R.L.B.P. (1983). Ice-thrust features in the Proterozoic Macaúbas Group, Jequitaiá area, Minas Gerais, Brazil. *Jour. of Geology*, **91**:113-116.
- GREGORY, R.T. & TAYLOR, H.P. (1981). An oxygen isotope profile in a section of Cretaceous oceanic crust. Samail ophiolite, Oman: evidence for  $\delta^{18}\text{O}$  buffering of the oceans by deep (>5km) seafloor-hydrothermal circulation at mid-ocean ridges. *Jour. Geophys. Res.*, **86**:2737-2755.
- GRIFFIN, W.L. & RYAN, C.G. (1995). Trace elements in indicator minerals: area selection and target evaluation in diamond exploration. *Journal of Geochemical Exploration*, **53**:311-337.
- GRIFFIN, W.L., JACQUES, A.I., SIE, S.H., RYAN, C.G., COUSENS, D.R., SUTER, G.F. (1988). Conditions of diamond growth: a proton microprobe study of inclusions in west Australian diamonds. *Contr. Miner. Petrol.*, **99**:143-158.
- GROSSI-SAD, J.H., LOBATO, L.M., PEDROSA-SOARES, A.C., SOARES FILHO, B.S. (1997). Projeto Espinhaço em CD-Rom. Belo Horizonte, *COMIG*. 2693p.
- GUIMARÃES, J.T. (1996). *A Formação Bebedouro no Estado da Bahia: faciologia, estratigrafia e ambientes de sedimentação*. Dis. Mestrado, Univ. Federal Bahia, 155p.
- GURNEY, J.J. (1984). A correlation between garnets and diamonds in kimberlites. In: *Glover, J.E. & Harris, P.G. (Eds.): Kimberlite occurrence and origin: A basis of conceptual models in exploration*. Univ. of Western Australia, Publ. **8**:143-166.
- GURNEY, J.J. (1989). Diamonds. In: *Diamonds and related rocks*. J. Ross (ed.). Geological Soc. Of Australia, Special Publication **14**:935-965.
- GURNEY, J.J. & SWITZER, G.S. (1973). The discovery of garnets closely to diamonds in the Finsch Pipe, South Africa. *Contr. Mineral. Petrol.*, **39**:103-116.
- GURNEY, J.J. & ZWEISTRA, P. (1995). The interpretation of the major element compositions of mantle minerals in diamond exploration. *Jour. of Geochemical Exploration*, **53**:293-309.
- GUTHRIE, G.D., VEBLEN, D.R., NAVON, O., ROSSMAN, G.R. (1991). Submicrometer fluid inclusions in turbid-diamonds coats. *Earth Plan. Sci. Lett.*, **105**(1-3):1-12.
- HAGGERTY, E.S. (1986). Diamond genesis in a multiply-constrained model. *Nature*, **320**:34-38.
- HAGGERTY, E.S. (1999). A diamond trilogy: Superplumes, supercontinents, and supernovae. *Science*. **285**:851-860.
- HARALYI, N.L.E., HASUI, Y., MORALES, N. (1991). O diamante pré-cambriano da Serra do Espinhaço, MG. Gemas e rochas ornamentais, *DNPM-CPRM*, vol. 4-A, p. 117-138.
- HARRIS, J.W. (1987). Recent physical, chemical and isotopic research of diamond. In: *Mantle Xenoliths*, Edited by P. H. Nixon, John Wiley & Sons, p. 477-500.
- HARRIS, J.W. (1992). Diamond geology. In: *The properties of natural and synthetic diamond* (ed. J.E. Field), Academic Press, 710p.
- HARRIS, J.W., HAWTHORNE, J.B., OOSTERVELD, M.M., WEHMEYER, E. (1975). A classification scheme for diamond and a comparative study of South African diamond characteristics. *Phys. Chem. of the Earth*, **9**:765-783.
- HARRIS, J.W., HAWTHORNE, J.B., OOSTERVELD, M.M. (1979). Regional and local variations in the characteristics of diamonds from some southern african kimberlites. *Proceedings of the Second International Kimberlite Conference*, Edited by J. F. Boyd & H. O. A. Meyer, **1**:27-41.
- HARRIS, J.W., HAWTHORNE, J.B., OOSTERVELD, M.M. (1983). A comparison of diamond characteristics from the De Beers pool mines, Kimberley South Africa. *Proceedings 3rd International Kimberlite Conference*, Edited by J. Kornprobst, Elsevier Press, **1**:1-13.
- HARRIS, J.W., HAWTHORNE, J.B., OOSTERVELD, M.M., WEHMEYER, E. (1975). A classification scheme for diamond, and a comparative study of south african diamond characteristics. *Physics and Chemistry of the Earth*, **9**:477-506.
- HARTE, B., GURNEY, J.J., HARRIS, J.W. (1980). The formation of peridotite suite inclusions in diamonds. *Cont. Min. Petrol.*, **72**:181-190.

- HARTE, B. & HARRIS, J.W. (1994). Lower mantle associations preserved in diamonds. *Miner. Mag.*, **58A**:384-385.
- HARTE, B., FITZSIMONS, C.W., HARRIS, J.W., OTTER, M.L. (1999). Carbon isotopes ratios and nitrogen abundances in relation to cathodoluminescence characteristics for some diamonds from Kaapvaal provinces. *South Africa Min. Mag.*, **63**(6):829-856.
- HASUI, Y., SADOWSKI, G.R., SUGUIO, K., FUCK, G.F. (1975). The Phanerozoic tectonic evolution of the western Minas Gerais State. *An. Acad. Bras. Ciências*, **47**:431-438.
- HAURI, E.H., WANG, J., PEARSON, D.G., BULANOVA, G.P. (2002). Microanalysis of  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , and N abundances in diamonds by secondary ion mass spectrometry. *Chem. Geol.*, **185**:149-163.
- HELMSTAEDT, H.H. (2002). Background for short course on diamond exploration. ADIMB, Brasília, 1-88.
- HELMSTAEDT, H.H. & GURNEY, J.J. (1994). Geotectonic controls on the formation of diamonds and their kimberlitic and lamproitic host rocks: applications to diamond exploration. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference, Araxá, Brazil*. **2**:236-250.
- HELMSTAEDT, H.H. & GURNEY, J.J. (1995). Geotectonic controls of primary diamond deposits: implications for area selection. *Journal of Geochemical Exploration*, **53**:125-144.
- HELMREICHEN, V. von (1846). Über das geognostische Vorkommen der Diamanten und ihre Gewinnungs – Methoden auf der Serra do Grão Mogol – 74 p., (*Braunmüller & Seidel*), Wien.
- HERVIG, R.L., SMITH, J.V., STEELE, I.M., GURNEY, J.J., MEYER, H.O.A., HARRIS, J.W. (1980). Diamonds: minor elements in silicate inclusions. Pressure-temperature implications. *Jour. Geophys. Res.*, **85**: 6919-6929.
- HETTICH, M. (1973). *Zur stratigraphie und gênese dès Macaúbas nordlich der Serra Negra, Espinhaço-Zone, Minas Gerais, Brasilien*. Tese doutorado, Univ. Freiburg, 42p.
- HETTICH, M. (1977). A glaciação proterozóica do centro-norte de Minas Gerais. *Rev. Bras. Geoc.*, **7**:87-101.
- HETTICH, M. & KARFUNKEL, J. (1978). Um esker, um varvito e seixos estriados no Grupo Macaúbas, norte de Minas Gerais. *EMOP*, **34**(2):5-8, Ouro Preto.
- HIPPERT, J. (1999). Are *s-c* structures, duplexes and conjugate shear zones different manifestations of the same scale invariant phenomenon? *Journal of Structural Geology*, **21**:975-984.
- HIRSCH, P.B., HUTCHISON, J.L., TICHMARSH, J. (1986). Voidites in diamond. Evidence for a crystalline phase containing nitrogen. *Phil. Magazine*, **54**:49-54.
- HOLLAS, M. (2002). *Basic atomic and molecular spectroscopy*. The Royal Society of Chemistry (publ.), London, 184p.
- HOFFMAN, P.F., KAUFMAN, A.J., HALVERSON, G.P. (1998). A Neoproterozoic snow ball Earth. *Science*, **281**:1342-1346.
- HUSSON, L. & MUGNIER, J.L. (2003). Three dimensional horizon reconstruction from outcrop structural data, restoration, and strain field of the Baisahi anticline, Western Nepal. *Journal of Structural Geology*, **25**:79-90.
- IBGE (1977). Folha Bocaiúva, SE-23-X-C-III, Carta do Brasil, escala 1:100.000.
- IBGE (1977). Folha Botumirim, SE-23-X-B-IV, Carta do Brasil, escala 1:100.000.
- IBGE (1977). Folha Itacambira, SE-23-X-D-I, Carta do Brasil, escala 1:100.000.
- INDA, H.A.V., SCHORSCHER, H.D., DARDENNE, M.A., SCHOBENHAUS, C., HARALYI, N.L.E., AZEVEDO BRANCO, P.C., RAMALHO, R. (1984). O Cráton do São Francisco e a Faixa de Dobramentos Araçuaí. In: *SCHOBENHAUS, C., ALMEIDA CAMPOS, C., DERZE, G.R., ASMUS, H.E. (eds.). Geologia do Brasil. DNPM*, p.194-284.
- IRVINE, T.N., & BARAGAR, W.R.A. (1971). A guide to the chemical classification of the common volcanic rocks of orogenic areas. *Bull. Geol. Soc. Am.*, **83**:29-40.
- ISOTTA, C.A., ROCHA-CAMPOS, A.C., YOSHIDA, R., (1969). Striated pavement of the upper-Precambrian Glaciation in Brazil. *Nature*, **5192**: 466-468; London.
- IZRAELI, E.S., HARRIS, J.W., NAVON, O. (2001). Brine inclusions in diamonds: a new upper mantle fluids. *Earth and Plan. Sci Lett.*, **187**(3-4):323-332.
- IWANUCH, W., MAKEEV, A.B., KONDO, M.M. (2005)a. Mineralogia, inclusões, catodoluminescência e composição isotópica do carbono dos carbonados de Andaraí, Chapada Diamantina, Bahia. In: *II South American Symp. On Diamond Geology*, Diamantina, 76-79.
- IWANUCH, W., MAKEEV, A.B., KONDO, M.M. (2005)b. Películas metálicas sobre os diamantes de Juína, Mato Grosso. In: *II South American Symp. On Diamond Geology*, Diamantina, 73-75.



- JACOB, D., JAGOUTZ, E., LOWRY, D., MATTEY, D., KUDRJAVTSEVA, G. (1994). Trace elements in diamondiferous eclogites from Siberia: remnants of Archean oceanic crust. *Geochim. Cosmochim. Acta*, **58**:5191-5207.
- JACOBSEN, S.B. & KAUFMAN, A.L. (1999). The Sr, C and O isotopic evolution of Neoproterozoic seawater. *Chemical Geology*, **16**:37-57.
- JANSE, A.J.A. (1991). Review of supposedly non-kimberlitic and non-lamproitic diamond host rocks. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. **2**:144-159
- JANSE, A.J.A. (1994). Is Clifford's Rule still valid? Affirmative examples from around the world. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. **2**:215-235.
- JANSE, A.J.A. & SHEAHAN, P.A. (1995). Catalogue of world wide diamond and Kimberlite occurrence: a selective and annotative approach. *Journal of Geochemical Exploration*, **53**:73-111.
- JAQUES, A.L., HALL, A.E., SHERATON, J., SMITH, C.B. & ROKSANDIC, Z. (1994). Peridotitic planar octahedral diamonds from the Ellendale lamproite pipes, Western Australia. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. **2**:69-77.
- JAVOY, M., PINEAUS, F., DELORME, H. (1986). Carbon and nitrogen isotopes in the mantle. *Chem. Geol.*, **57**:41-62.
- JAYNES, C. (1978). Natural polycrystalline diamond. *Industrial Diamond Review*, January, p. 14-23.
- JENNINGS, C.M.H. (1995). The exploration context for diamonds. *Journal of Geochemical Exploration*, **53**:113-124.
- KAGI, H., TAKAHASHI, K., HIDAKA, H., MASUDA, A. (1994). Chemical properties of Central African carbonado and its genetic implications. *Geochimica et Cosmochimica Acta*, **58**(12):2629-2638.
- KAISER, W. & BOND, W.L. (1959). Nitrogen, a major impurity in common type I diamond. *Phys. Rev.*, **115**:857-863.
- KAMINSKIY, F.V. (1984). The diamond content of non-kimberlitic eruptive rocks. Izd-vo Nedra, Moscow.
- KAMINSKIY, F.V. (1987). Genesis of carbonado-polycrystalline aggregate of diamond. *Doklady Akademii Nauk SSSR*, **294**:439-440.
- KAMINSKIY, F.V. (1994). Carbonado and yakutite: properties and possible genesis. *Proceedings 5th International Kimberlite Conference*, CPRM Special Publication, **2**:136-143.
- KAMINSKIY, F.V., KULAKOVA I.I., OGLOBINA, A.I. (1985). Polycyclic aromatic hydrocarbons in carbonado and diamond. *Doklady Akademii Nauk SSSR*, **283**:985-988.
- KAMINSKY, F.V., ZAKHARCHENKO, O.D., DAVIES, R., GRIFFIN, W.L., KHACHATRYAN-BLIONVA, G.K., SHIRYAEV, A.A. (2001). Super diamonds from the Juina area, Mato Grosso State, Brazil. *Contr. Min. Petrol.*, **140**:734-753.
- KAMINSKY, F.V., FELDMAN, A.A., VARLAMOV, V.A., BOYCO, A.N., OLOFINSKY, L.N., SHOFMAN, I.L. & VAGANOV, V.I. (1995). Prognostication of primary diamond deposits. *Journal of Geochemical Exploration*, **53**:167-182.
- KANE, R.E. (1987). Three notable fancy-color diamond: purplish red, purple-pink, and reddish purple. *Gems & Gemology*. vol. XXIII, p. 90-95.
- KARFUNKEL, J. & CHAVES, M.L.S.C. (1994). Diamante na região do Médio São Francisco, Minas Gerais. In: *Cong. Bras. Geol.*, **38, Bol.**, **1**:191-193.
- KARFUNKEL, J. & CHAVES, M.L.S.C. (1995). Conglomerados cretácicos da Serra do Cabral, Minas Gerais: um modelo para a redistribuição colúvio-aluvionar dos diamantes do médio São Francisco. *Geociências*, UNESP, **14**: 59-72.
- KARFUNKEL, J. & DUPONT, H. (2001). Depósitos glaciogênicos neoproterozóicos em Minas Gerais: controvérsias e consenso. In: *XI Simp. Geol. Minas Gerais*. Em CD-ROM.
- KARFUNKEL, J. & HOPPE, A. (1988). Late Proterozoic Glaciation in Central-East Brazil: Synthesis and model. *Palaeogr., Palaeoclimatol., Palaeoecol.*, **65**:1-21.
- KARFUNKEL, B. & KARFUNKEL, J. (1975). Fazielle Entwicklung der mittleren Espinhaço-Zone mit besonderer Berücksichtigung des Tillitproblems. *Tese Doutorado*, Univ. Freiburg, 86 p., Freiburg.
- KARFUNKEL, B. & KARFUNKEL, J. (1977). Fazielle entwicklung der mittleren Espinhaço-zone mit besonderer Berücksichtigung des tillit-problems. *Geogisches Jahrbuch, Beihefte* **24**:3-91.
- KARFUNKEL, J., PEDROSA-SOARES, A.C., DOSSIN, I.A. (1985). O Grupo Macaúbas em Minas Gerais, revisão dos conhecimentos. In: *Simp. Geol. Minas Gerais 3, Anais*. Belo Horizonte, SBG, p.45-59.
- KARFUNKEL, J., CHAVES, M.L.S.C., SVISERO, D.P., MEYER, M.O.A. (1994). Diamonds from Minas Gerais, Brazil: an update and sources, origin and production. *Inter. Geology Review*, **36**:1019-1032.

- KARFUNKEL, J., CHAVES, M.L.S.C., BANKO, A.G., HOOVER, D.B. (1998). Colluvial diamond and quartz deposits from the Espinhaço Range (MG): genesis and economic importance. *In: XL Cong. Bras. Geol., Belo Horizonte, 1998. Anais...*, SBG-MG.
- KARFUNKEL, J., DUPONT, H., SCHOBENHAUS, C., NOCE, C.M. (2000). The Neoproterozoic São Francisco Glaciation in Central-Eastern Brazil. *Profil, Bd.*, 19 (in CD-ROM).
- KARFUNKEL, J., HORTA MOREIRA, P.C., RIBEIRO, M.C., & COSTA FRANCO, A.L. (1984). Aspectos genéticos e deposicionais do Grupo Macaúbas na região da Barragem do Paraúna e sua importância na contribuição para o modelo paleogeográfico e geotectônico. *In: Soc. Bras. Geol., Rio de Janeiro, Cong. Bras. Geol.*, 40, *Anais*, 5:489-493.
- KARFUNKEL, J., DANTAS, M.S., PIMENTA, M.A., PINHEIRO, M.V.B., PERSIANO, A.I.C., LAMEIRAS, F.S., ADDAD, J.E. (2000). Métodos espectroscópicos na análise gemológica. Espectroscopia Micro-Raman (EMR). *REM*, 53(4):239-245.
- KARFUNKEL, J., MARTINS, M.S., SCHOLZ, R., McCANDLESS, T.E. (2001)a. Diamonds from the Macaúbas River Basin (MG): Characteristics and Possible Source. *In: 3º Simp. Bras. Geol. do Diamante / 1º Simp. Sul-Amer. de Geol. Diamante, Abstracts, Brasília*, p. 23.
- KARFUNKEL, J., MARTINS, M., SCHOLZ, R., SAMPAIO, N.P., BRITO, W. (2001)b. Características e origem do diamante da bacia do rio Macaúbas, centro-norte de Minas Gerais. *In: XI Simp. Geol. Minas Gerais*. Em CD-ROM.
- KARFUNKEL, J., SCHOBENHAUS, C., DUPONT, J.H., NOCE, C.M. (2001)c. The Neoproterozoic São Francisco Glaciation (Brazil): facies distribution and model. LAK/Stuttgart. Em CD-ROM.
- KATTAH, C. (1991). *Análise faciológica e estratigráfica do Jurássico Superior/Cretáceo Inferior na porção meridional da Bacia do São Francisco, oeste do Estado de Minas Gerais*. Ouro Preto, EM-UFOP, Dis, Mestrado, 227p.
- KAUFMAN, A.J., KNOLL, A.H., NARBONNE, G.M. (1997). Isotopes, ice ages, and Proterozoic earth history. *Proc. Natl., Acad. Sci., USA*, 94, 6600-6605.
- KERR, P.F., GRAF, D.L., BALL, S.H. (1948). Carbonado from Venezuela. *Am. Mineral.*, 33: 251-254.
- KENNEDY, M.J., RUNNEGAR, B., PRAVE, A.R., HOFFMANN, K.H. & ARTHUR, M.A. (1998). Two or four Neoproterozoic glaciations? *Geology*, 26:1059-1063.
- KESSON, S.E. & RINGWOOD, A.E. (1989). Slab-mantle interactions: 2. The formation of diamonds. *Chem. Geol.*, 78:97-118.
- KING, L.C. (1956). A geomorfologia do Brasil Oriental. *Revista Brasileira de Geografia*, 18(2):147-265.
- KIRK, E.C., WILLIAMS, D.A., AHMED, H. (1989). Cross-sectional transmission electron microscopy of precisely selected regions from semiconductor devices. *Inst. Phys Conf. Series*, Vol. 100, 501p.
- KIRKLEY, M.B. (1998). The origin of diamonds: Earth processes. *In: The nature of diamonds*, G.E. Harlow (ed.). Univ. Press and American Museum of Natural History, Cambridge, 48-65.
- KIRKLEY, M.B., GURNEY, J.J., OTTER, M.L., HILL, S.J., DANIELS, L.R. (1991). The application of C isotope measurements to the identification of the sources of C in diamonds: a review. *Applied Geochemistry*, 6:477-494.
- KIRKLEY, M.B., GURNEY, J.J. & RICHARD, R.S. (1994). Jwaneng framesites: carbon isotope and inclusion compositions. *In: CPRM/ Núcleo Brasília, Fifth international kimberlite conference, Araxá, Brazil*. 2:127-135.
- KLEIN, G.V. (1982). Probable sequential arrangement of depositional systems on cratons. *Geology*, 10(1):17-22.
- KLEIN, C. & HURLBUT JR. C.S. (1999). Manual of mineralogy. 21 ed. New York, *John Wiley & Sons*. 681p.
- KLEIN-BENDAVID, O., WIRTH, R., NAVON, O. (2006). TEM imaging and analysis of microinclusions in diamonds: a close look at diamond-growing fluids. *American Mineralogist*, 21 (no prelo).
- KNAUER, L.G. (1990). *Evolução geológica do Precambriano da porção centro-leste da Serra do Espinhaço Meridional e metalogênese associada*. Dissertação de mestrado, IG/UNICAMP, 298p.
- KNAUER, L.G. (1999). *Serra do Espinhaço Meridional: considerações sobre a estratigrafia e análise da deformação das unidades proterozóicas*. Tese de Doutorado, IGCE/UNESP, Rio Claro, 324p.
- KNOLL, A.H. & WALKER, M.R. (1992). Latest Proterozoic stratigraphy and Earth history. *Nature*, 356:673-678.
- KNOLL, A.H., HAYES, J.M., KAUFMAN, A.J., SWETT, K., LAMBERT, I.B. (1986). Secular variation in carbon isotope ratios from Upper

- Proterozoic successions of Svalbard and East Greenland. *Nature*, **321**:832-838.
- KUKHARENKO, A.A. (1954). On rounded diamond crystals. *Uchem. Zap. Leningrad Gos. Univ.*, **178**(4), 55p. (em russo).
- LEONARDOS, O.H. (1937). Diamante e carbonado no estado da Bahia. *Mineração e Metalurgia*, **V**:183-192.
- LEONARDOS, O.H. (1940). Tilito metamórfico de Carandaí, Minas Gerais. *An. Acad. Bras. Ciências*, **12**:243-259.
- LEOST, I., STACHEL, T., BREY, G.P., HARRIS, J.W., RYABCHIKOV, I.D. (2003). Diamond formation and source carbonation: mineral association in diamonds from Namibia. *Contr. Miner. Petrol.*, **145**:15-24.
- LEUNG, I.S., TAYLOR, L.A., TSAO, C.S., HAN, Z. (1996). SiC in diamond and kimberlites: implications for nucleation and growth of diamond. *Int. Geol. Rev.*, **37**:483-496.
- LIMA, S.A.A.; MARTINS-NETO, M.A.; PEDROSA-SOARES, A.C.; CORDANI, U.G. & NUTMAN, A. (2002). A Formação Salinas na área-tipo, NE de Minas Gerais: uma proposta de revisão da estratigrafia da Faixa Araçuaí com base em evidências sedimentares, metamórficas e de idades U-Pb SHRIMP. *Revista Brasileira de Geociências*, **32**(4): 491-500.
- LISBOA, M.A.R. (1906). Ocorrência de seixos facetados no Planalto Central do Brasil. *An. Esc. Minas, Ouro Preto*, **8**:23-74.
- LOMONOSOFF, M. (1843). Note sur le gisement des diamants au Brésil. *Ann. de Chemie et de Physique* 3, série, vol. VII. Paris.
- LORENTZ, N.J., CORSETTI, F.A., LINK, P.K. (2004). Seafloor precipitates and C-isotope stratigraphy from the Neoproterozoic Scout Mountain Member of the Pocatello Formation, southeast Idaho: implications for Neoproterozoic earth system behavior. *Precambrian Research*, **130**:57-70.
- LUDWIG, K.R. (1999). Using Isoplot/EX, version 2, a geochronological Toolkit for Microsoft Excel. *Berkeley Geochronological Center Special Publication*, **1a**:47.
- LUDWIG, K.R., 2001. Squid 1.02: a user manual. In: *Berkeley Geochronological Center Special Publication*, **12**:19.
- LUTOEV, V.P., GLUKHOV, Y.V., ISAENKO, S.I. (2000). Epigene nitrogen defects and metallic films on the surface of diamonds from the Middle Timan. *Doklady Earth Sciences*, **375**(8):1251-1254.
- LUTOEV, V.P., GLUKHOV, Y.V., MARTINS, M., ISAENKO, S.I., PETROVSKY, V.A., KARFUNKEL, J. (2004). Spectroscopic characterization of diamonds and carbonados from Macaúbas river basin (MG). In: *42<sup>TH</sup> Cong. Bras. Geol. Araxá*. CD-Rom.
- MACAHADO, N., SCHRANK, A., ABREU, F.R., KNAUER, L.G. & ALMEIDA-ABREU, P.A. (1989). Resultados preliminares da geocronologia U/Pb na Serra do Espinhaço Meridional. *V Simp. Geol. Minas Gerais, Anais*, p.171-174, Belo Horizonte.
- MAGALHÃES, L. (1988). *Análise estrutural qualitativa nos sedimentos do Grupo Bambuí, região sudeste da bacia do São Francisco (Faixa Sete Lagoas-Serra do Cipó)*. Dis. Mestrado, Escola de Minas/DEGEO/UFOP, 110p.
- MAGEE, C.W. & TAYLOR, W.R. (1998). Constraints on the history and origin of carbonado from luminescence studies. In: *7<sup>TH</sup> Int. Kimb. Conf., Cape Town*, 527-528.
- MAGEE, C.W. & TAYLOR, W.R. (1999). Constraints on the history and origin of carbonado from luminescence studies. In: *7<sup>TH</sup> Int. Kimb. Conf., Cape Town*, 529-532. J.J. Gurney, M.D. Pascoe, R.S.H. (eds.).
- MAKEEV, A.B. & FILIPPOV, V.N. (1999). Películas metálicas sobre a superfície de diamantes naturais (Jazida de Ichetiyyu), Médio Timan. *Dok. Ros.Ak. Nauk*. **368**(6): 808-812.
- MAKEEV, A.B., FILIPPOV, V.N., OBIDEN, S.K., SAPARIN, G.V., IWANUCH, W., MARTINS, M. & KARFUNKEL, J. (2001). Características Mineralógicas dos carbonados brasileiros dos Estados da Bahia e Minas Gerais. p.162-165. In: *Borintseva N.A., Bryanchaninova N.I., Kulikova K.V., Makeev A.B., Samomolkova, M.F., Ponomareva G.V. e Pystin A.M. 2001. Diamantes e potencial diamantífero da região de Timan-Urais. Syktyvkar. Geoprint*. 240p.(em russo).
- MAKEEV, A.B., IWANUCH, W., FILIPPOV, V.N., OBIDEN, G.V., SAPARIN, W. (2001). Mineralogia, composição das inclusões e catodoluminescência de carbonados do Estado da Bahia, Brasil. *Geol. Rudn. Mestorozhd.*
- MALZAHN, H. (2000). *Diamante*. Extralapis, 18p.
- MARTINS, M.S. (2002). *Geologia da bacia do Rio Macaúbas (MG), com ênfase nas ocorrências diamantíferas*. Dis. Mestrado, IGC/UFMG, 86 p.
- MARTINS, M.S., KARFUNKEL, J., ADDAD, J., SCHOLZ, R., PEREGOVICH, B. (2000). Das Macaubas-Becken in zentral-nord Minas Gerais, Brasilien und Diamantvorkommen. *Profil, Bd.*, 19 (in CD-ROM).
- MARTINS, M., KARFUNKEL, J., SCHOLZ, R., HAYDEN, S. (2001). Geologia da bacia do rio

- Macaúbas (MG). In: *Simp. Geol. Minas Gerais*. Em CD-ROM.
- MARTINS, M., PETROVSKY, V.A., SAMOILOVICH, M.I., FILIPPOV, V.N., KARFUNKEL, J., SUKHAREV, A.Y. (2003). Experimental and natural polycrystalline diamond systems: a case study for carbonado genesis. In: *XIX Cong. Bras. Geoquímica, Belém*. CD-Rom.
- MARTINS, M. & KARFUNKEL, J. (2004). Magmatismo neoproterozóico ao longo do anticlinal de Itacambira. In: *42° Cong. Bras. Geologia, Araxá*. CD-Rom.
- MARTINS, M., PETROVSKY, V.A., SAMOILOVICH, M.I., FILIPPOV, V.N., KARFUNKEL, J., SUKHAREV, A.Y. (2004). The evolution of natural polycrystalline diamond system based on carbonado characteristics. In: *42° Cong. Bras. Geologia, Araxá*. CD-Rom.
- MARTINS-NETO, M.A. (1993). *The sedimentary evolution of a Proterozoic rift: the basal Espinhaço Supergroup, Minas Gerais, Brazil*. Univ. Freiburg, Tese de doutorado, 155p.
- MARTINS-NETO, M.A. (2000). Tectonics and sedimentation in a Paleo/Mesoproterozoic Rift-Sag Basin (Espinhaço Basin, southeastern Brazil). *Prec. Res.* **103**:147-173.
- MARTINS-NETO, M.A. & ALKMIM, F.F. (2001). Estratigrafia e evolução tectônica das bacias neoproterozóicas do Paleocôntinente São Francisco e suas margens: registro da quebra de Rodínia e colagem de Gondwana. In: C.P. Pinto, M.A. Martins Neto (eds.). *Bacia do São Francisco: geologia e recursos minerais*. SBG MG, Belo Horizonte, 31-54.
- MARSHAL, T.R. & BAXTER-BROWN, R. (1995). Basic principles of alluvial diamond exploration. *Journal of Geochemical Exploration*, v.53: p.277-292.
- MARSHALL, T.R. (1994). The diamondiferous gravels of the southwestern Transvaal, South Africa. In: CPRM/ Núcleo Brasília, *Fifth international kimberlite conference*, Araxá, Brasil. **2**:187-201.
- MASCARENHAS, J.F., PEDREIRA, A.J., MISI, A., MOTTA, A.C., SILVA-SÁ, J.H. (1984). Província São Francisco. In: O Pré-Cambriano do Brasil. F.F.M. Almeida & Y. Hasui (coord.), Edgar Blücher, São Paulo, 46-122.
- McCALLUM, M.E. (1976). An emplacement model to explain contrasting mineral assemblages in adjacent kimberlite pipes. *Journal of Geology*. **84**:673-684.
- McCALLUM, M.E., HUNTLEY, P.M., FALK, R.W. & OTTER, M.L. (1991). Morphological, resorption and etch feature trends of diamonds from kimberlite populations within the Colorado Wyoming State line district, USA. In: CPRM/ *Fifth international kimberlite conference*, Araxá, Minas Gerais, Brasil. Vol. 2: Diamonds: characterization, genesis and exploration, p.32-50.
- McCALLUM, M.E., MABARAK, C.D., COOPERSMITH, H.G. (1979). Diamonds from kimberlites in the Colorado-Wyoming State Line District. Proceedings of the Second International Kimberlite Conference, Edited by J. F. Boyd & H. O. A. Meyer, vol. 1, p. 42-58.
- McCALLUM, M.E., HUNTLEY, P.M., FAK, R.W., OTTER, M.L. (1994). Morphological, resorption and etch feature trends of diamonds from kimberlite populations within the Colorado-Wyoming State Line District, USA. In: *Proceedings da V Conf. Intern. de kimb.* (ed. H.O.A. Meyer e O.H. Leonardos). V.2 – Diamonds:characteruzation, gênesis, and exploration. Araxá, Brasil:32-50.
- McCANDLESS, T.E. (1990). Kimberlite xenocryst wear in high-energy fluvial systems: experimental studies. *Journ. Geochem. Expl.*, **37**: 323-331.
- McCANDLESS, T.E. (2001). Textural analysis of the diamonds from the Macaúbas river basin (MG). *Rel. Interno. Universidade do Arizona, USA*, 23p.
- McCANDLESS, T.E. & NASH, W.P. (1996). Detrital mantle indicator minerals in southwestern Wyoming, USA: evaluation of mantle environment, igneous host, and diamond exploration. *Expl. Mining Geology*, **5**(1):33-44.
- McCANDLESS, T.E. & GURNEY, J.J. (1989). Sodium in garnet and potassium in clinopiroxene: criteria for classifying mantle eclogites. In: *Kimberlites and related rocks*. J. Ross (ed.). **2**:827-832.
- McCANDLESS, T.E. & GURNEY, J.J. (1997). Diamond eclogites: comparison with carbonaceous chondrites, carbonaceous shales, and microbial carbon-enriched MORB. *Russian Geology and Geophysics*. **38**(2):394-404.
- McCANDLESS, T.E., LETENDRE, J. & EASTOE, C.J. (1996). Morphology and carbon isotope composition of microdiamonds from Dachine, French Guiana. *Unpublished report*. **2**:550-556.
- McCANDLESS, T.E., NASH, W.P. & HANSEL, W. (1995). Mantle indicator minerals in ant mounds and conglomerates of the southern Green River Basin, Wyoming. *Wyoming Geology Association, Field Guidebook*: p.153-164.
- McCANDLESS, T.E., WALDMAN, M.A. & GURNEY, J.J. (1994). Macrodiamonds and microdiamonds from Murfreesboro lamproites, Arkansas: morphology, inclusions and carbon isotope geochemistry. In: CPRM/ Núcleo Brasília, *Fifth*

- international kimberlite conference*, Araxá, Brazil. **2**: 78-97.
- McCNAUGHT, M.A. & MITRA, G. (1996). The use of finite strain data in constructing a retrodeformable cross-section of the Meade thrust sheet, southeastern Idaho, USA. *Jour. Estructural Geology*, **16**:585-602.
- MELTON, C.E., & GIARDINI, A.A. (1974). The composition and significance of gas released natural diamonds from Africa and Brazil. *American Mineralogist*, **59**(7-8): 775-782.
- MEMMI, J.M. & McCALLUM, M.E. (1994). Finite element modeling of resistivity data from kimberlite intrusions in Wyoming, USA. In: CPRM/ Núcleo Brasília, *Fifth international kimberlite conference*, Araxá, Brazil. **2**:382-388.
- MENDELSSOHN, M.J. & MILLEDGE, H.J. (1995). Geologically significant information from routine analysis of the mid-infrared spectra of diamonds. *International Geology Review*. **37**:95-110.
- MENDELSSOHN M.J. & MILLEDGE, H.J. (1995). Morphological characteristics of diamond populations in relation to temperature-dependent growth and dissolution rates. *International Geology Review*. Vol. **37**: p.285-312.
- MESCHEDE, M. (1986). A method of discriminanting between different types of mid-ocean ridge basalts and continental tholeiites with the Nb-Zr-Y diagram. *Chem. Geol.*, **56**:207-218.
- MEYER., H.O.A. (1985). Genesis of diamond. A mantle saga. *American Mineralogist*, **70**:344-355.
- MEYER., H.O.A. (1987). Inclusions in diamond. In: *Mantle Xenoliths*. P. H. Nixon (ed.), New York, Wiley, p. 501-522.
- MEYER, H.O.A. & McCALLUM (1997). Diamonds and their sources in the Venezuela portion of the Guiana Shield. *Economic Geology*, **88**:989-998.
- MEYER, H.O.A. & TSAI, H.M. (1976). Mineral inclusions in natural diamond: their nature and significance. *Min. Sci. Eng.*, **8**:242-261.
- MEYER., H.O.A., GARWOOD, B.L., SVISERO, D.P., SMITH, C.B. (1994). Alkaline ultrabasic intrusions of western Minas Gerais, Brazil. IN: *CPRM/DNPM Special Publication*, Edited by H. O. A. Meyer & O. H. Leonardos, Brasília, **1**:140-155.
- MEYER, H.O.A., MILLEDGE, H.J., SUTHERLAND, F.L. & KENNEWELL, P. (1997). Unusual diamonds and unique inclusions from New South Wales, Austrália. *Russian Geology and Geophysics* .**38**(2):305-331.
- MEYER, H.O.A., ZHANG ANDI, MILLEDGE, H.J. & MENDELSSOHN, M.J. (1994). Diamonds and mineral inclusions in diamonds from Shandong and Liaoning Provinces, China. In: CPRM/ Núcleo Brasília, *Fifth international kimberlite conference*, Araxá, Brazil. **2**:98-105.
- MIALL, A.D. (1990). *Principles of sedimentary basin analysis*. 2 ed. New York, Springer-Verlag. 668p.
- MIALL, A.D. (1996). *The geology of stratigraphic sequences*. Berlim, Springer-Verlag, 433p.
- MILLEDGE, H.J., SHELKOV, D., PILLINGER, C.T., VERSHOVSKY, A.B. (1996). Comparative studies of carbonado, impactite, and other polycrystalline diamond. *Eos*, **77**(17):S143.
- MILLEDGE, H.J., WOODS, P.A., BEARD, A.D., SHELKOV, D., WILLIS, B. (1998). Cathodoluminescence of polished carbonado. *7th International Kimberlite Conference, Extended Abstracts*, Cape Town, p. 589-590.
- MITCTHELL, R.H. (1991). Kimberlites and lamproites: primary sources of diamonds. *Geoscience Canada*, **18**:1-16.
- MITCTHELL, R.H. (1995). The role of petrography and lithochemisrty in exploration for dimondiferous rocks. *Journal of Geochemical Exploration*, **53**:339-350.
- MITCTHELL, R.H. & BERMAN, S.C. (1991). *Petrology of lamproites*. Plenum Press, New York.
- MITRA, S. (1986). Duplex structures and imbricate thrust systems: geometry, structural position and hydrocarbon potential. *AAPG, Bul.* **70**:1087-1112.
- MONTES, M.L. (1977)a. Os conglomerados diamantíferos da Chapa Diamantina – Bahia, Brasil. Dis. Mestrado, IG/UnB, 102p.
- MONTES, A.S.L. (1977)b. O contexto estratigráfico e sedimentológico da Formação Bebedouro na Bahia. Dis. Mestrado, IG/UnB, 100p.
- MOORE, J.M. & MOORE, A.E. (2004). The roles of primary kimberlitic and secondary Dwyka glacial sources in the development of alluvial and marine diamond deposits in Southern Africa. *Journal of African Earth Sciences*, **38**(6):115-134.
- MOISSAN, H. (1895). Sur la presance du graphite, du carbonado, et le diamants microscopiques dans la terre bleue du Cap. *Comptes Rendu, Academie Sciences de Paris*, **116**:292-295.
- MOORE, M. & LANG, A.R. (1974). On the origin of the rounded rhombic dodecahedral habit of natural diamond. *Journal of Crystal Growth*, **26**:133-139.
- MORAES, L.J. (1928)a. O carbonado em Minas Gerais. *Min. Comb. e Transp.*, Ano I, **10**, 239p.

- MORAES, L.J. (1928)b. Geologia da região diamantina de Minas Gerais - Relatório de atividades. DNPM, *Serv. Geol. Min. Rel. Ann. Dir.* 1928, 29-34, Rio de Janeiro.
- MORAES, L.J. (1932). Área ocupada pela Formação Macaúbas no norte de Minas Gerais – *Ann. Acad. Bras. Sci.*, **4**:111-114.
- MORAES, L. J. (1934). Depósitos diamantíferos do norte do Estado de Minas Gerais. *Boletim DNPM/SFPM*, Rio de Janeiro, **3**:1-61.
- MORAES, L.J., BARBOSA, O., LISBOA, J.M.A., LACOURT, F., GUIMARÃES, D. (1937). Geologia Econômica do Norte de Minas Gerais. *Boletim DNPM/SFPM*. 19, 191p. Rio de Janeiro.
- MORAES, L.J. & GUIMARÃES, D. (1929). Estudos sobre a rocha matriz do Diamante – *DNPM, Serv. Geol. Min. Rel. Ann. Dir.* 1928, 171-174, Rio de Janeiro.
- MORAES, L.J. & GUIMARÃES, D. (1930). Geologia da região diamantífera do Norte de Minas Gerais – *An. Acad. Bras. Cienc.*, **2**:153-186, Rio de Janeiro.
- MORGAN, P. (1995). Diamond exploration from the bottom up: regional geophysical signature lithosphere conditions favorable for diamond exploration. *Journal of Geochemical Exploration*, **53**:145-165.
- MORRIS, T.F. & KASZYCKI, C.A. (1997). Prospector's guide to drift prospecting for diamonds, northern Ontario, Ontario. *Geological Survey, Miscellaneous Paper* 167, 63p.
- MORTON, A.C. (1984). Stability of detrital heavy minerals in tertiary sandstones from the North Sea basin. *Clay Minerals*, **19**:287-308.
- MPSOKOS, E.D. & KOSTOPOULOS, D.K. (2001). Diamond, former coesite and supersilicic garnet in metasedimentary rocks from the Greek Rhodope: a new ultrahigh-pressure metamorphic province established. *Earth and Planetary Science Letters*, **192**:497-506.
- NAVON, O. (1999). Diamond formation in the Earth's mantle. *Proc. do 7<sup>th</sup> Con. Intern. Kimb., África do Sul*, **2**:584-605.
- NAVON, O., HUTCHEON, I.D., ROSSMAN, G.R., WASSERBURG, G.J. (1988). Mantle-derived fluids in diamonds micro-inclusions. *Nature*, **335**:784-789.
- NISBET, D.E.G., MATTEY, D.P., LOWRY, D. (1994). Can diamonds be dead bacteria? *Nature*, **367**, 694.
- NIXON, P.H. (1995). The morphology and nature of primary diamondiferous occurrences. *Journal of Geochemical Exploration*, **53**:41-71.
- NOCE, C.M., GUIMARÃES, M.L.V., ROQUE, N.C., SGARBI, G.N., KNAUER, L.G., GROSSI SAD, J.H. (1996). Mapa geológico (1:100.000) da Folha Itacambira, Projeto Espinhaço, COMIG-UFMG.
- NOCE, CM. (1997). Texto explicativo da Folha Itacambira. In: J.H. GROSSI-SAD, L.M. LOBATO, A.C. PEDROSA-SOARES, B.S. SOARES FILHO (1997) - Projeto Espinhaço em CD-Rom. Belo Horizonte, COMIG.
- NOCE, CM. (1997). Geologia da Folha Carbonita. In: J.H. GROSSI-SAD, L.M. LOBATO, A.C. PEDROSA-SOARES, B.S. SOARES FILHO (1997) - Projeto Espinhaço em CD-Rom. Belo Horizonte, COMIG.
- NOCE, C.M., PEDROSA-SOARES, A.C., GROSSI-SAD, J.H., BAARS, F.J., GUIMARÃES, M.L.V., MOURÃO, M.A.A., OLIVEIRA, M.J.R. & ROQUE, N.C. (1993). Nova divisão estratigráfica regional do Grupo Macaúbas na Faixa Araçuaí: o registro de uma bacia neoproterozóica. *Anais 9<sup>o</sup> Simp. Geol. Minas Gerais – SBG Núcleo MG*. Bol. **14**: 29-31.
- OLIVEIRA, F.P. (1881). A exploração das minas de galena do Ribeirão do Chumbo, afluente do Abaeté. Estudo da zona percorrida de Ouro Preto até o lugar. *An. Esc. Minas, Ouro Preto*, **1**:39-105.
- OLIVEIRA, J.R.P. (1989). *Comportamento estrutural dos grupos Macaúbas e Bambuí na porção centro norte de Minas Gerais*. Dis. de Mestrado, EM?DEGEO/UFOP, 102p.
- OLIVEIRA, A.I., & LEONARDOS, O.H. (1940). *Geologia do Brasil*. Com. Bras. Centenários. Portugal.
- OLIVEIRA, L.H.B. & MARTINS, M.S. (1999). *Mapeamento Litoestrutural da região diamantífera de Grão Mogol*. Trabalho de Graduação, IGC/UFMG, Belo Horizonte, 91p.
- ORLOV, Yu. L. (1966). Evidence that the rounding of diamond is caused by dissolution process. *Min. Sbor. Louvsk. Univ.*, **20**:129-130.
- ORLOV, Yu. L. (1977). The mineralogy of diamond. *Izdatel'stva Nanka*. Translated from *Russian Geology and Geophysics* by WILEY, J. & SONS. New York. 235 p.
- OTTER, M.L. & GURNEY, J.J. (1989). Mineral inclusions in diamonds from the Sloan diatremes, Colorado-Wyoming State Line district, North America. In: Proc. da V Conf. Intern. Kimb. Kimberlites and related rocks, **2**:1042-1053.
- OTTER, M.L., GURNEY, J.J., KIRKLEY, M.B., HARTE, B., McCANDLESS, T.E. (1991). The carbon isotope composition of Sloan diamonds: evidence of a subducted ophiolite sequence? *Geochim., Cosmochim., Acta*, **58**:247-256.

- OTTER, M.L., McCALLUM, M.E., GURNEY, J.J. (1994). A physical characterization of the Sloan (Colorado) diamonds using a revised diamond description scheme. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. **2**: 15-31
- OZIMA, M. & TATSUMOTO, M. (1997). Radiation-induced diamond crystallization: origin of carbonados and its implications on meteorite nano-diamonds. *Geochimica et Cosmochimica Acta*. **61**(2):369-376.
- OZIMA, M., ZACHU, S., MATSUHISA, Y. (1991). Constrain from noble-gas contents on the origin of carbonado diamonds. *Nature*, **351**:472-474.
- PALY'ANOV, Y.N., SOKOL, A.G., BORZDOV, Y.M., KHOKHRYAKOV, A.F. (2002). Fluid-bearing alkaline carbonate melts as the medium for the formation of diamonds in the earth's mantle: an experimental study. *Lithos*, **60**:145-159.
- PARFENOFF, A., POMEROL, C., TOURENQ, J. (1970). Les minéraux en grain - Méthodes d'étude et détermination. *Masson et Cie, Éditeurs, Paris*, p. 50-87.
- PASTEL, A.R. & RAMATHAN, S. (1962). Etch pits in diamond surfaces. *Phil. Magazine*. **7**:1305-1314.
- PATTISON, D.R.M. & LEVINSON, A.A. (1995). Are enedral microdiamonds formed during ascent and decompression of kimberlite magma? Implications for use of microdiamonds in diamond grade estimation. *Appl. Geochem*. **10**:725-738.
- PEARCE, T.H. & CANN, J.R. (1973). Tectonic setting of basic volcanic rocks determined using trace element analyses. *Earth Plant. Sci. Lett.*, **19**:290-300.
- PEARSON, D.G., SHIRLEY, S.B., BULANOVA, G.P., CARLSON, R.W., MILLEDGE, H.J. (1999). Dating and paragenetic distinction of diamonds using Re-Os isotope system: application to some Siberian diamonds. *Proc. of the 7<sup>th</sup> Int. Kimb. Conf., Cape Town*. **2**:637-643.
- PEDREIRA, A.J. (1996). *Geologia e recursos minerais da bacia Metassedimehtar do Rio Pardo*. CBPM, ser. Arquivos Abertos, **11**, 17p.
- PEDROSA-SOARES, A.C. (1981). *Geologia da folha de Virgem da Lapa, Minas Gerais*. CPMT/IGC/UFGM, Relat. Final de bolsa de Aperfeiçoamento CNPq, Belo Horizonte, 52p.
- PEDROSA-SOARES, A.C. (1995). *Potencial aurífero do Vale do Araçuaí, MG: história da exploração, geologia e controle tectono-metamórfico*. Tese de doutoramento, IG/UnB, Brasília, 177p.
- PEDROSA-SOARES, A.C. & WIEDEMANN-LEONARDOS, C.M. (2000). Evolution of the Araçuaí Belt and its connection to the Ribeira Belt, Eastern Brazil. In: U.G. CORDANI; E.J. MILANI; A. THOMAZ FILHO; D.A. CAMPOS (eds) *Tectonic evolution of South America*. Rio de Janeiro, p.265-285.
- PEDROSA-SOARES, A.C., VIDAL, P., LEONARDOS, O.H., BRITO NEVES, B.B. (1998). Neoproterozoic ocanic remnants in eastern Brazil: further evidence and refutation of an exclusively ensialic evolution for the Araçuaí-West Congo orogen. *Geology*, **26**:519-522.
- PEDROSA-SOARES, A.C., NOCE, C.M., VIDAL, Ph., MONTEIRO, R.L.B.P., LEONARDOS, O.H. (1992). Towards a new tectonic model for the Late Proterozoic Araçuaí (SE Brazil)-West Congolian (SW Africa) Belt. *Jour. South Amer. Earth Sci.* **6**(1/2):33-47.
- PELTZIER, G. & SAUCIER, F. (1996). Present-day kinematics of Asia derived from geologic faults rates. *Jour. Geophys. Res.*, **101**(27):943-956.
- PENHA, U.C. (2001). *Geologia dos conglomerados cretácicos da Serra da Água Fria e dos depósitos diamantíferos de Jequitaiá*. Tese de doutorado, UNESP, 148p.
- PENHA, U.C., KARFUNKEL, J., ANGELI, N. (2005). Diamondiferous deposits in the Jequitaiá area (Minas Gerais, Brazil): a consequence of neotectonic processes. *N. Jb. Geol. Paläont. Abh., Stuttgart*, **236**(1/2):207-224.
- PEREIRA, L.F. (1992). *Relações tectono-estratigráficas entre as unidades Canastra e Ibiá na região de Coromandel e Guarda-Mor, MG*. Brasília, IG-UnB, 75p.
- PETRI, S., COIMBRA, A.M., AMARAL, G., OJEDA, H.O., FÚLFARO, V.J., PONÇANO, W.P. (1986). Código Brasileiro de nomenclatura estratigráfica. *Rev. Bras. Geoc.*, **16**:372-376.
- PETROVSKY, V.A., MARTINS, M., LUTOEV, V.P., KARFUNKEL, J., GLUKHOV, Y.V., SUKHAREV, A.E., FILIPPOV, V.N. (2002). Minerological and genetic features of carbonados from the Minas Gerais State (Brazil). *Vestnik, Revista do Inst. Geológico da República de Komi, RAS*, **2**(1): 7-12. Em russo.
- PETROVSKY, V.A., MARTINS, M., KARFUNKEL, J., SAMOILOVICH, M.I., BELYANIN, A.F., LUTOEV, V.P., SUKHAREV, A.E., GLUKHOV, Y.V. (2003). Carbonados: physical properties, mineral inclusions and conditions of formation. In: *Proc. 9<sup>th</sup> Int. Conference – High Technology in Russian Industry, Moscow*, 206-220 (em russo).
- PETROVSKY, V.A., MARTINS, M., KARFUNKEL, J., YAKOVLEV, E.N., FILONENKO, V.P., SUKHAREV, A.E., LUTOEV, V.P. (2004). Carbonado genesis on the natural and experimental data. In: *32<sup>th</sup> Int. Cong. Geologia, Florence*. CD-Rom.

- PETROVSKY, V.A., SILAEV, V.I., MARTINS, M., SUKHAREV, A.E., KARFUNKEL, J. (2005). Carbonados from Brazil: mineralogical characteristics and structures. In: *II South American Symp. On Diamond Geology, Diamantina*, 113-116.
- PETTIJOHN, F.J., & POTTER, P.E. (1964). *Atlas and glossary of primary sedimentary structures*. Springer, New York, 370p.
- PETTIJOHN, F.J., POTTER, P.E., SIEVER, R. (1973). *Sand and sandstones*. Springer, New York, 618p.
- PFLUG, R. (1965). A geologia da parte meridional da Serra do Espinhaço e zonas adjacentes, Minas Gerais – *DNPM, Div. Geol. Min., Bol.* 226., 51p., Rio de Janeiro.
- PFLUG, R. (1968). Observações sobre a estratigrafia da Série Minas na região de Diamantina, Minas Gerais – *DNPM, Div. Geol. Min. Not. Prel. Estud.*, 142, 20 p., Rio de Janeiro.
- PFLUG, R. & RENGER, F. (1973). Estratigrafia e evolução geológica da Margem SE do Cráton Sanfranciscano – *An. XXVII, Congr. Bras. Geol.*, 2:5-19, Aracaju.
- PLUIJM, B.A. & MARSHAK, S. (1997). *Earth structure. An introduction to structural geology and tectonics*. WCB/McGraw-Hill, 495p.
- PUPIN, J.P. (1980). Zircon and granite petrology. *Contr. to Mineralogy and Petrology*, 73:207-220.
- PUSHAREV, Y.D. (1997). Two types of interaction crust-mantle substance and a new approach to the problems of deep ore-formation. *Proc. of RAS*, 355(4):524-526.
- PUSHAREV, Y.D., GOROKHOVSKY, B.M., LARIN, A.M. (2001). Interaction role of crust and mantle substance in the formation of endogenic giant deposits: isotopic-geochemical approach. *Regional geology and metallogeny*, 11:73-80.
- QUEIROZ, W.P. (2002). *O Grupo Macaúbas na região de Couto Magalhães de Minas: estratigrafia e evolução*. Dis. Mestrado, IGC/UFMG, 214p.
- RAAL, F.A., & ROBINSON, D.N. (1980). Green for rarity. *Nuclear Active*, 23:5-8.
- RAKIN, V., MARTINS, M., KARFUNKEL, J. (2004). The shape of curve-faced diamonds. In: *32<sup>TH</sup> Intern. Cong. of Geology, Florence*. CD-Rom.
- RAMSAY, J.G. (1980). Shear zones geometry: a review. *Journ. Struct. Geol.*, 2:83-99.
- READING, H.G. (1986). *Sedimentary environments and facies*. Blackwell Scientific Publications, London, 615p.
- RENGER, F. (1979). Evolução dos conceitos geológicos da Serra do Espinhaço. *I Simp. Geol. Minas Gerais, Anais*, p.9-27, Diamantina.
- RENGER, F.E. & ALMEIDA-ABREU (2000). *Fieldtrip Guide Serra do Espinhaço*. 31<sup>TH</sup> Int. Geol. Cong., Rio de Janeiro. 35p.
- RENGER, F.E., KNAUER, L.G. (1995). Espinhaço Quo vais? (onde está? onde vai?) – A evolução dos conhecimentos sobre a cordilheira do Espinhaço Meridional em Minas Gerais entre 1979 e 1995. *Geonomos*. 3:31-39.
- RICHARDSON, S.H (1986). Latter-day origin of diamonds of eclogitic paragenesis. *Nature*. 322:623-626.
- RICHARDSON, S.H., ERLANK, A.J., HARRIS, J.W. (1984). Origin of diamonds in old enriched mantle. *Nature*. 30:198-202.
- RICHARDSON, S.H., ERLANK, A.J., HARRIS, J.W. & HART, S.R. (1990). Eclogitic diamonds of proterozoic age cretaceous kimberlites. *Nature*. 346:54-56.
- RICHARDSON, S.H., HARRIS, J.W. & GURNEY, J.J. (1993). Three generations of diamonds from old continental mantle. *Nature*. 366:256-258.
- RIMANN, E. (1915). Über kimberlit und alnoit in Brasilien. *Tschermaks Miner. Petrog. Mitt*, 23:2244-2262.
- RIMANN, E. (1917). A kimberlita no Brasil. *An. Esc. Minas*, 15:27-32.
- RINGWOOD, A.E. (1991). Phase transformations and their bearing on the constitution and dynamics of the mantle. *Geochimica et Cosmochimica Acta*. Vol. 55 : p.2083-2110.
- ROBERTSON, R. FOX, J.J., MARTIN, A.E. (1934). Two types of diamond. *Phil. Trans. Royal Soc. London*, A232:463-538.
- ROBINSON, D.N. (1978). The characteristics of natural diamond and their interpretation. *Mineral Science Engineering*, 10:55-72.
- ROBINSON, D.N. (1980). *Surface textures and other features of diamonds*. Tese doutorado, Natal, África do Sul, 221 p.
- ROBINSON, D.E., SCOTT, J.A., VAN NIEKERK, A., ANDERSON, V.G. (1989). The sequence of events reflected in the diamonds of some southern African kimberlites. *Geol. Soc. Australia. Sp. Publ.*, 14(2): 990-1000.



- ROGOZINA, E.A. (1983). Gás formation in katagenesis of organic substance from sedimentary rocks. Leningrad: *Nauka*, 164p. (em russo).
- ROLIM, V.K. (1992). Uma interpretação das estruturas tectônicas do Supergrupo Espinhaço, baseada na geometria dos falhamentos de empurrão. *REM, EM-Ouro Preto*, **45**(1/2):75-77.
- ROMBOUTS, L. (1995). Statistical distributions for diamonds. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. **2**:202-214.
- ROUFFAER, E.J.H.F. (1988). *On the origin of alluvial diamonds along the Atlantic Coast of southwest Africa*. Ph.D. Thesis. Leuven. 155p.
- RUDNICK, R.I., McDONALD, W.F., CHAPPEL, B.W. (1993). Carbonatite metasomatism in the northern Tanzanian mantle: petrography and geochemical characteristics. *Earth Plan. Sci. Lett.*, **114**:463-475.
- RUSSELL, S.S., PILLINGER, C.T., ARDEN, J.W., LEE, M.R., OTT, U. (1992). A new type of meteoritic diamond in the enstatite chondrite Abee. *Nature*, **256**:206-209.
- SALVADOR, A. (1994). Lithostratigraphic units. In: A. Salvador (ed.). *International Stratigraphic Guide. A guide to stratigraphic classification, terminology and procedure*. 2<sup>o</sup> ed. Colorado, Geological Society of America (International Subcommission on Stratigraphic Classification of International Union of Geological Sciences – IUGS; International Commission on Stratigraphy). 214p.
- SAMOILOVICH, M.I., SANZHARLINSKY, N.G., LAPTEV, V.A. (1985). Research of processes of spontaneous origin and continuous growth of diamonds in system metal-graphite. In: *Synthesis of Minerals*. Moscow, Nedra: 368-390 (em russo).
- SAMOILOVICH, M.I., PETROVSKY, V.A., SHILOV, Y.A., KHITUNIN, V.P. (2000). Experimental simulation on natural diamond growth. In: *31<sup>TH</sup> Int. Geological Cong., Rio de Janeiro*. Cd-Rom.
- SANO, Y., YOKOSHI, R., CHAVES, M.L.S.C., OZIMA, M. (2002). Ion microprobe Pb-Pb dating of carbonado, polycrystalline diamond. *Precambrian Research*, **11**:155-168.
- SANTOS, R.V., ALVARENGA, C.J.S., DARDENNE, M.A., SIAL, A.N., FERREIRA, V.P. (2000). Carbon and oxygen isotope profiles across Meso-Neoproterozoic limestones from central Brazil, Bambuí and Paranoá groups. *Precambrian Res.*, **104**:107-122.
- SANTOS, R.V., ALVARENGA, C.J.S., RAMOS, M.L.S., FONSECA, M.A., CUKROV, N. (2002). Isótopos de carbono na transição meso-neoproterozóico: implicações paleoambientais. *XLI Cong. Bras. Geol.*, pg. 337, João Pessoa – PB.
- SCHOBENHAUS, C. (1993). *O Proterozóico Médio no Brasil com ênfase à região Centro-leste, uma revisão*. Freiburg, Univ. Freiburg, Tese Doutorado, 166p.
- SCHOBENHAUS, C., CAMPOS, D.A., DERZE, G.R., ASMUS, H.E. (1984). *Geologia do Brasil*. Brasília, MME-DNPM, 501p.
- SCHÖLL, W. (1973). Sedimentologie und geochemie der bambuí Gruppe in SE-rond des São Francisco Becken, Minas Gerais (Brasilien). Univ. Heidelberg, Tese doutorado, 115p.
- SCHÖLL, W. & FOGAÇA, A.C.C. (1979). Estratigrafia da Serra do Espinhaço na região de Diamantina. In: *I Simp. Geol. Minas Gerais*, Anais, 55-73pgs.
- SCHRANK, A., DOURADO, B.C.V. & BIONDI, J.C. (1978). Estudo preliminar dos metavulcanitos do Grupo Macaúbas na região do Alto Jequitinhona, Minas Gerais. *XXX Cong. Bras. Geol., Anais 3*:1323-1335, Recife.
- SCHRAUDER, M. & NAVON, O. (1994). Hydrous and carbonatic mantle fluids in fibrous diamonds from Jwaneng, Botswana. *Geochem. Cosmochim. Acta*, **58**(2):761-771.
- SCHRAUDER, M., KOEBERL, C., NAVON, O. (1996). Trace element analyses of fluid in fibrous diamonds from Jwaneng, Botswana. *Geochem. Cosmochim. Acta*, **60**:4711-4724.
- SCHULZE, D.J. (1986). Calcium anomalies in the mantle and a subducted metaserpentine origin for diamonds. *Nature*, **319**:483-485.
- SELIVERSTOV, V.A., GORSHKOV, A.I., SCHEKA, S.A. & SIVTSOV, A.V. (1996). Diamonds and Carbonado from Primorya: Mineralogy, Kristallocemistry, Genesis. *Geology of ore Deposits*. **38**(6):485-499.
- SGARBI, G.N., SGARBI, P.B.A., CAMPOS, J.E.G., DARDENNE, M.A., PENHA, U.C. (2001). Bacia Sanfranciscana: o registro fanerozóico da Bacia do São Francisco. In: C.P. Pinto, M.A. Martins Neto (eds.). *Bacia do São Francisco: Geologia e recursos minerais*. SBG-MG, Belo Horizonte, 93-138.
- SHAFRANOVSKII, I.I. (1940). On the crystallography of Brazilian-type diamond. *Dokl. An SSSR*, **26**(7) (em russo).

- SHELKOV, D., VERKHOVSKY, A.B., MILLEDGE, H.J., PILLINGER, C.T. (1997). Carbonado: a comparison between Brazilian and Ubangui sources with other forms of microcrystalline diamond based on carbon and nitrogen isotopes. *Russian Geol. Geophys.*, **38**:332-340.
- SHERVAIS, J.W. (1982). Ti-V plots and the petrogenesis of modern and ophiolitic lavas. *Earth Plan. Sci. Lett.*, **59**:101-118.
- SHIBATA, K., KAMIOKA, H., KAMINSKIY, F.V., KOPTIL, V.I., SVISERO, D.P. (1993). Rare earth elements patterns of carbonados and yakutite: evidence for their crustal origin. *Mineralogical Magazine*, **57**:607-611.
- SHUMILOVA, T. (2003). Mineralogy of native carbon. *UD RAS Press*, Yekaterinburg (em russo).
- SHUMILOVA, T. (2004). Variety of carbon phases, conditions and mechanisms of diamond crystallization. *Lithos*,
- SIAL, A.N. (2004). Isótopos estáveis e espectrometria de massa de fonte gasosa. *NEG-LABISE-UFPE, Notas de aula*. 147p.
- SIGA Jr., O. (1986). *A evolução geotectônica da porção nordeste de Minas Gerais com base em informação geocronológica*, Dissertação de mestrado – IG/USP, São Paulo, 140p.
- SILAEV, V.I., PETROVSKY, V.A., SUKHAREV, A.Y., MARTINS, M. (2005). Carbonados: results of researches. *Vestnik, Revista do Inst. Geológico da República de Komi, RAS*, **11**: 7-13. Em russo.
- SILVA, R.R. (1995). *Contribution to the stratigraphy and paleogeography of the lower Espinhaço Supergroup (Mesoproterozoic), between Diamantina and Gouveia, Minas Gerais State, Brazil*. Tese de Doutorado, Univ. Freiburg, 115p.
- SIMAKOV, S.K. (1998). Redox state of earth's upper mantle peridotites under the ancient cratons and its connection with diamond genesis. *Geochimica et Cosmochimica Acta*. **62**(10):1811-1820
- SMITH, C.B. (1985). What is a Kimberlite?. In: J.E. Glover & P.G. Harris. *Kimberlite occurrence and origin: a basics for conceptual models in exploration*. **8**: 1-18
- SMITH, V.J. & DAWSON, J.B. (1985). Carbonados: diamond aggregates from early impacts of crustal rocks? *Geology*, **13**:342-343.
- SNYDER, G.A., TAYLOR, L.A., BEARD, B.L., CROZAZ, C., HALLIDAY, A.N., SOBOLEV, V.N., SOBOLEV, N.V. (1997). The origins of yakutian eclogitic xenoliths. *Jour. Petrol.*, **38**:85-113.
- SOBOLEV, N.V. (1985). Crystalline inclusions in diamonds from New South Wales, Austrália. GLOVER, J.E. & HARRIS, P.G. *Kimberlite occurrence and origin: a basics for conceptual models in exploration*. **8**:213-226.
- SOBOLEV, N.V. (1985). Kimberlites of the Siberian platform: their geological and mineralogical features. GLOVER, J.E. & HARRIS, P.G. *Kimberlite occurrence and origin: a basics for conceptual models in exploration*. **8**:275-287.
- SOBOLEV, N.V., KAMINSKY, F.V., GRIFFIN, W.L., YEFIMOVA, E.S., WIN, T.T., RYAN, C.G., BITARNOV, A.I. (1997). Mineral inclusions in diamonds from the Sputnik kimberlite pipe, Yakutia. *Lithosphere*, **39**:135-157.
- SOBOLEV, N.V., TAYLOR, L.A., SNYDER, G.A., KELLER, R.A., YEFIMOVA, E.S., SOBOLEV, V.N., SHIMIZU, N. (1998). Extreme chemical diversity in the mantle during eclogitic diamond formation: evidence from 35 garnet and 5 pyroxene inclusions in a single diamond. *Int. Geol. Rev.*, **40**:567-578.
- SPETSIUS, Z.V. (1995). Occurrence of diamond in the mantle: a case study from the Siberian Platform. *Journal of Geochemical Exploration*, **53**: 25-39.
- SPETSIUS, Z.V. & GRIFFIN, B.J. (1998). Secondary phases associated with diamonds in eclogites from the Udachnaya Pipe: implications for diamond genesis. In: *7<sup>th</sup> Int. Kimb. Conf., Cape Town*, 850-851.
- SPIX, J.B. von & MARTIUS, C.F.P. von (1823/1831). Reise im Brasilien. *Lind.*, **3**:1388p., Munchen.
- STACHEL, T. (2001). Diamonds from the asthenosphere and the transition zones. *Europena Jour. Min.*, **13**:883-892.
- STRECKEISEN, A. (1978). Classification and nomenclature of volcanic rocks, carbonatites, and melilitic rocks. IUGS Subcommission on the Systematics of Igneous Rocks. *Neues Jahrb. Mineral. Abh.*, **134**:1-14.
- SUKHAREV, A.Y., PETROVSKY, V.A., MARTINS, M. (2005). The mechanism of phase transformation of graphite into diamond. In: *II South American Symp. On Diamond Geology*, Diamantina, 162-165.
- SUNAGAWA, I. (1984). Morphology of natural and synthetic diamond crystals. In: *Materials Science of the Earth's Interior*, I. Sunagawa (ed.), p. 303-330.
- SUTHERLAND, D.G. (1982). The transport and sorting of diamonds by fluvial and marine processes. *Economic Geology*, **77**(7):1613-1620.
- SUTHERLAND, G.B.B.M., BLACKWELL, D.E., SIMERAL, W.G. (1954). The problem of the two types of diamond. *Nature*, **174**:902-904.

- SUTHERLAND, F.L., TEMBY, P., RAYNOR, L.R. & HOLLIS, J.D. (1994). A review of the east Australian diamond province. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. 2:170-185.
- SVISERO, D.P., HASUI, Y., DRUMOND, D. (1979). Geologia de kimberlitos do Alto Paranaíba, Minas Gerais. *Mineração Metalurgia*, **406**:34-38.
- SVISERO, D.P. (1995). Distribution and origin of diamonds in Brazil: An overview. *Journal of Geodynamics*, **20**(4):493-514.
- TAKAOKA, N. & OZIMA, M. (1978). Rare gas elemental abundances and isotopic compositions in diamonds. *Geological Survey*. Open-file Paper, N. 701a, 418-420.
- TAYLOR, L.A. & ANAND, M. (2004). Diamonds: time capsules from the Siberian mantle. *Chemie der Erde*, **64**(1):1-74.
- TAYLOR, W.T., JAQUES, A.L. & RIDD, M. (1990). Nitrogen-defect aggregation characteristics of some Australian diamonds: Time-temperature on the source regions of pipe and alluvial diamonds. *American Mineralogist*. **75**:1290-1310.
- TAYLOR, L.A., KELLER, R.A., SNYDER, G.A., WANG, W., CARLSON, W.D., HAURI, E.H., McCANDLESS, T., KIM, K.R., SOBOLEV, N.V., BEZBORODOV, S.M. (2000). Diamonds and their mineral inclusions and what they tell us: a detailed "pull-apart" of a diamondiferous eclogite. *Int. Geol. Rev.*, **42**:959-983.
- TAYLOR, L.A., MILLEDGE, H.J., BULANOVA, G.P., SNYDER, G.A. & KELLER, R.A. (1998). Metasomatic eclogitic diamond growth: Evidence from multiple diamond inclusions. *Intern. Geol. Rev.* **40**:663-676.
- TEIXEIRA, N.A. (1996). Assoalho oceânico no complexo ultramáfico do Quatipuru (PA): implicações geotectônicas para a faixa móvel Araguaia. In: *Cong. Bras. Geol.*, 39. Salvador, SBG, *Anais*, **6**:117-120.
- TEIXEIRA, W.; SABATÉ, P.; BARBOSA, J.; NOCE, C.M. & CARNEIRO, M.A. (2000). Archean and Paleoproterozoic tectonic evolution of the São Francisco Craton. In: CORDANI, U.G.; MILANI, E.J.; THOMAZ FILHO, A.; CAMPOS, D.A. (eds) *Tectonic evolution of South America*. Rio de Janeiro, p.101-137.
- TITKOV, S.V., GORSHKOV, A.I., VINOKUROV, S.F., BERSHOV, L.V., SODOLOV, D.I., SIVTSOV, A.V. (1998). Carbonado from Yakutian diamond deposits (Russia): microinclusions, impurities and paramagnetic centers. *7th International Kimberlite Conference, Extended Abstracts*, Cape Town, p. 914-916.
- TOLANSKY, S. & PATEL, A.R. (1975). Rectilinea etch pits in diamond. *Phil. Magazine.* **2**:1003-1005.
- TOMILENKO, A.A., CHEPUKROV, A.I., PAL'YANOV, YU. (1997). Volatile components in the upper mantle according to the data on fluid inclusions study. *Geology and geophysics*, **38**:276-285.
- TOMPKINS, L.A. (1994). Tectono-structural environments of primary diamond source rocks in Brazil: a review. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. 2:259-267.
- TORQUATO, J.R. (1980). Isótopos de carbono e oxigênio aplicados no estudo do Pré-vambriano brasileiro. Tese Prof. Titular, UFPE, 512p.
- TOULKERIDIS, T., BABINSKI, M., BUSCHWALDT, R., BRITO NEVES, B.B., TODT, W. & SANTOS, R. (1999). Are Varangian or Sturtian the glacial deposits on the São Francisco Craton? Evidence from age determination of sedimentary rocks and minerals of the neoproterozoic Una Group. In: *South America Symposium on isotope geology*, 2. Cordoba, 1999, anais, p.453-456.
- TOWIE, N.J., BUSH, M.D., MANNING, E.R., MARX, M.R. & RAMSAY, R.R. (1994). The Aries diamondiferous kimberlite pipe central kimberley Block, western Australia: Exploration, setting, and evaluation. In: *CPRM/ Núcleo Brasília, Fifth international kimberlite conference*, Araxá, Brazil. 2:319-328.
- TRAUTMAN, R.I., GRIFFIN, B.J., BULANOVA, G.P. (1998). Growth features and nitrogen aggregation properties of microdiamonds derived from kimberlitic diatremes. In: *Sixth Int. Kimberlitic Conference. Novosibirsk*. Extended abstract, 926-928.
- TROMPETTE, R. (1994). *Geology of western Gondwana 2000-500Ma. The Pan-Africano Brasileiro Amalgamation of South America and Adjacent Africa*. Rotterdam, Balkena, 366p.
- TROMPETTE, R. (2000). Gondwana evolution; its assembly at round 600 Ma. *C.R. Acad. Sci. Paris / Earth and Planetary Sciences*, **330**:305-315.
- TRUEB, F.L. & BUTTERMAN, W.C. (1969). Carbonado: A microstructural study. *American Mineralogist*, **54**:412-425.
- TRUEB, F.L. & de WYS, E.C. (1971). Carbon from Ubangi - a micro-structural study. *American Mineralogist*, **56**:1252-1268.
- UHLEIN, A. (1991). Transição Cráton-Faixa Araçuai (Ciclo Brasileiro) no estado de Minas Gerais. Aspectos estratigráficos e estruturais. *Dissertação de Doutorado*, USP, São Paulo, 259p.

- UHLEIN, A. & QUEMENEUR, J.J.G. (2000). Estrutura e deformação brasileira nos diques máficos da Serra do Espinhaço Meridional. *Geonomos*, **8**(2):19-25.
- UHLEIN, A., TROMPETTE, R.R. & ALVARENGA, C.J.S. (1994). Late proterozoic gravitational sedimentation on a continental margin under glacial influence, the Jequitai-Macaúbas sequence (Minas Gerais, Brazil). *In: Intern. Sedim. Cong.*, 14, 1994. *Abstracts...*, G, 85-86, Recife.
- UHLEIN, A., TROMPETTE, R., EGYDIO-SILVA, M.E. (1995). Rifteamentos superpostos e tectônica de inversão da borda sudeste do Cráton do São Francisco. *Geonomos*, **3**(1):99-107.
- UHLEIN, A., TROMPETTE, R.R. & ALVARENGA, C.J.S. (1999)a. Neoproterozoic glacial and gravitational sedimentation on a continental rifted margin: the Jequitai-Macaúbas sequence (Minas Gerais, Brazil). *Jour. South Am. Earth Sci.*, **4-5**/2:1-7.
- UHLEIN, A., EGYDIO-SILVA, M., TROMPETTE, R.R., VAUCHEZ, A., KATTAH, C., (1999)b. Paleogeografia e inversão tectônica da Faixa Araçuaí. SBG, Simp. Nac. Est. Tect., Lençóis, 1999. *Anais...*, Lençóis, 26-29.
- UHLEIN, A., ALVARENGA, C.J.S., TROMPETTE, R., DUPONT, H., EGYDIO-SILVA, M., CCUKROV, N., LIMA, O.N.B. (2004). Glaciação neoproterozóica sobre o cráton do São Francisco e faixas dobradas adjacentes. *In: V. Mantesso Neto, A. Bartorelli, C. Dal Ré Carneiro e B. Brito Neves (orgs.) Geologia do Continente Sul-Americano: evolução da obra de Fernando Flávio Marques Almeida*. São Paulo; Editora Beca, 539-553.
- UROVSKAYA, A.A. & ORLOV, Y.L. (1968). Nature of plastic deformation of diamond crystals. *Doklady Akademii Nauk SSSR*, **154**:112-115.
- VANCE, E.R., HARRIS, J.W., MILLEDGE, H.J. (1973). Possible origins of  $\alpha$ -damage in diamonds from kimberlite and alluvial sources. *Mineral Magazine*. **39**:349-360.
- VINOGRADOV, A.P., KROPOTOVA, O.I., ORLOV, Y.L., GRINENKO, V.A. (1996). Isotopic composition of diamond and carbonado crystals. *Geochem. Int.*, **3**:1123-1125.
- VIVEIROS, J.M.F. & WALDE, D. (1976). Geologia da Serra do Cabral, Minas Gerais, Brasil. *Munst. Forsch. Geol. Paläont*, **38/39**:15-27.
- VIVEIROS, J.M.F., SÁ, E.L., VILELA, O.V., SANTOS, O.M., MOREIRA, S. (1978). Geologia dos vales dos rios Peixe Bravo e Alto Vacaria, norte de Minas Gerais. *In:XXX Cong. Bras. Geol.*, **1**:243-254, Recife.
- VIVEIROS, J.M.F., SÁ, E.L., VILELA, O.V., SANTOS, O.M., MOREIRA, S., NETO, F.M., VIEIRA, W.S. (1979). Geologia dos vales dos rios Peixe Bravo e Alto Vacaria, norte de Minas Gerais. *In: Simp. Geol. Minas Gerais, Anais*, 75-87, Diamantina.
- XU, S., OKAY, A.I., JI, S., SENGOR, A.M.C., SU, W., LIU, Y., JIANG, L. (1992). Diamond from the dabie Shan metamorphic rocks and its implication for the tectonic setting. *Science*, **256**:80-82.
- XU, B., JIAN, P., ZHENG, H., ZOU, H., ZHANG, L., LIU, Z. (2005). U-Pb zircon geochronology and geochemistry of Neoproterozoic volcanic rocks in the Tarim Block of northwest China: implications for the breakup of Rodinia supercontinent and Neoproterozoic glaciations. *Precambrian Research*, **136**:107-123.
- ZASHU, S. & HIYAGON, H. (1995). Degassing mechanisms of noble gases from carbonado diamonds. *Geochimica et Cosmochimica Acta*. **59**(7):1321-1328.
- ZINTCHOUK, N.N. & KOPTIL', V.I. (2003). *Typomorphism of the Siberian Platform diamonds*. ALROSA Co. Ltd., Editora Moscoe Nedra, 601P. (em russo).
- YERZERSKIY, V.A. (1986). High pressure polymorphs produced by the shock transformation of coals. *Int. Geol. Rev.*, **28**:221-228.
- WALDE, D., KARFUNKEL, J., KARFUNKEL, B. (1978). O Grupo Macaúbas em Minas Gerais: estratigrafia, gênese e correlações. SBG, Centro-Oeste, Bol. **7/8**:98-113.
- WALKER, R.G. (1979). Facies models. *Geoscience Canada Reprint Series 1*, 211p.
- WALKER, R.G. (1992). Turbidites and submarine fans. *In: R.G. Walker & N. James (eds.) Facies models. Response to sea level change*. Geological Association of Canada, 239-264.
- WALKER, R.G. & JAMES, N.P. (1992). *Facies models. Response to sea level change*. Ontario, Geological Association of Canada, 454p.
- WALMSLEY, J.C. & LANG, A.R. (1992)a. On submicrometer inclusions in diamond coat. Crystallography and composition of ankerites and related rhombohedral carbonates. *Mineralogical Magazine*, **56**:533-543.
- WALMSLEY, J.C. & LANG, A.R. (1992)b. Oriented biotite inclusions in diamond coat. *Mineralogical Magazine*, **56**:108-111.
- WENTFORT, R.H. & BOVENKERK, H.P. (1961). On the origin of natural diamonds. *Astrophysics Journal*, **134**:995-1005.

WENTFORT, R.H., DEVRIES, R.C., BUNDY, F.P. (1980). Sintered superhard material. *Science*, **208**:873-880.

WHITE, S.H., de BOORDER, H. & SMITH, C.B. (1995). Structural controls of Kimberlite and Lamproite emplacement. *Journal of Geochemical Exploration*, **53**:245-264.

WILKS, J., WILKS, E. (1995). *Properties and applications of diamonds*. Butterworth Heineman, 525p.

WILLIAMS, H.E. (1925). Notas geológicas e econômicas sobre o vale do Rio São Francisco. B. SGMB, 12, Rio de Janeiro.

WILLIAMS, I.S., CLAEISSON, S. (1987). Isotope evidence for the Precambrian province and Caledonian metamorphism of high grade paragneiss from the Seve Nappes, Scandinavian Caledonides. Part II. Ion microprobe zircon U–Th–Pb. *Contrib. Mineral. Petrol.* **97**:205–217.

WIRTH, R. (2004). Focused Ion Beam (FIB): A novel technology for advanced application of micro- and nanoanalysis in geosciences and applied mineralogy. *European Journal of Mineralogy*, **16**:863-876.

WIRTH, R. (2005). Micro- and nanometre sized inclusions in diamonds and microdiamonds: a new source of information about diamond genesis and fluid composition. In: *II South American Symp. on Diamond Geology, Diamantina*, 184.

WIRTH, R. & ROCHOLL, A. (2003). Nanocrystalline diamond from the Earth's mantle underneath Hawaii. *Earth and Planetary Science Letters*, **211**:357-369.

WOODS, G.S., van WYK, J.A., COLLINS, A.T. (1990). The nitrogen content of type Ib synthetic diamond. *Phil. Magazine*, **B62**:589-595.

WOODS, G.S., & COLLINS, A.T. (1983). Infrared absorption spectra of hydrogen complexes in type I diamonds. *Jour. Phys. Chem. Solids*, **44**(5):471-475.

WOODWARD, N.B., GRAY, D.R., SPEARS, D.B. (1986). Including strain data in balanced cross-sections. *Journal of Structural Geology*, **8**:313-324.

WYLLIE, P.J. & RYABCHIKOV, I.D. (2000). Volatile components, magmas and critical fluids in upwelling mantle. *Jour. Petrology*, **41**:1195-1206.